

THE BEETLES *of* DORSET

An annotated checklist



Adrian Mylward

A note on using the checklist:

The checklist runs to over 150 pages. It has not been designed to be printed but species can be found by using Control F to open the Search box. You can then type in part/all of a species name to find a summary of Dorset records.

Version control			
1	Report prepared by:	Adrian Mylward	21/12/20
2	Updated	Adrian Mylward	05/04/22

Cover image: *Cylindera germanica* Cliff Tiger Beetle © Bryan Edwards

The Beetles of Dorset (Vice-county 9)

In his *History of Glanvilles Wootton* of 1878 C.W. Dale included records of 920 species of Coleoptera, the vast majority of which were from the neighbourhood of his home there. Then in 1926, at the age of 23, E.J. Pearce published *A List of the Coleoptera of Dorset*. Pearce pulled together the efforts of many recorders, and he subsequently added three supplements, in 1927, 1929, and 1931, 'bringing the grand total to date up to 2149 species and varieties'. Unlike Dale, who spent most of his life at Glanvilles Wootton, Pearce did not remain in the county. His 1957 Royal Entomological Society handbook on the Pselaphidae (now a subfamily of the beetle family Staphylinidae) is still much used today.

The county list below, containing 2700 species, is the first since that of Pearce. Its design is deliberately simple, to enable it to be easily updated at a time of frequent new discoveries and continual changes in nomenclature. It is published online and the hope is that it will be updated annually.

The nomenclature follows the *Checklist of Beetles of the British Isles* (Duff, A.G. 2018) but some names from the 2012 checklist have been added to help avoid confusion. Subgenera have only been included where they are needed to justify the order of the listing.

In Pearce's lists there are a few dates, but the majority of species names are followed only by a list of localities. These are fairly general: 'Wareham' or 'Upwey' for example, and the original list contained many species with just "'Dorset"; (A. Ford)' added. The second supplement did include more detailed information on localities from A. Ford for some, but not all, such species. For a variety of reasons, not all of Pearce's listed species have found their way into the list below.

This checklist is not designed to be a guide to finding beetles in Dorset, but simply an inventory of species found in the county. A 'record' here is defined by the presence of a species at a certain location on a certain date, irrespective of the number of specimens. Where the number of records exceeds 9 it is listed as 'many'.

Thanks are due to all those who have contributed records of beetles over the years, but particularly to Alison Stewart of the Dorset Environmental Records Centre for all her help and support.

Adrian F. Mylward

April 2022

1. Family SPHAERIUSIDAE Erichson, 1845

Sphaerius acaroides Walzl, 1838

Pearce: no records
First record: 1990
Number of records: 7
Latest record: 2002

2. Family GYRINIDAE Latreille, 1810

Gyrinus (Gyrinulus) minutus Fabricius, 1798

Pearce: 5 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2019

Gyrinus (Gyrinus) aeratus Stephens, 1835

Pearce: 3 localities
First subsequent record: 1977
Number of subsequent records: 2
Latest record: 1980

Gyrinus (Gyrinus) caspius Ménétries, 1832

Pearce: 1 locality
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2021

Gyrinus (Gyrinus) marinus Gyllenhal, 1808

Pearce: 2 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2019

Gyrinus (Gyrinus) substriatus Stephens, 1828

Pearce: generally common and widely distributed
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Gyrinus (Gyrinus) urinator Illiger, 1807

Pearce: 1 locality
First subsequent record: 1974
Number of subsequent records: many
Latest record: 2021

Orectochilus villosus (Müller, O.F., 1776)

Pearce: 2 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2021

3. Family HALIPLIDAE Aubé, 1836

Brychius elevatus (Panzer, 1793)

Pearce: 4 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2020

Haliphus (Haliplidius) confinis Stephens, 1828

Pearce: 2 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2019

Haliphus (Haliplidius) obliquus (Fabricius, 1787)

Pearce: 4 localities
First subsequent record: 1991
Number of subsequent records: many
Latest record: 2010

Haliphus (Haliphus) fluviatilis Aubé, 1836

Pearce: 4 localities
First subsequent record: 1977
Number of subsequent records: many
Latest record: 2019

Haliphus (Haliphus) heydeni Wehncke, 1875

Pearce: 4 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2018

Haliphus (Haliphus) immaculatus Gerhardt, 1877

Pearce: 1 locality
First subsequent record: 1937
Number of subsequent records: 8
Latest record: 2005

Haliphus (Haliphus) lineolatus Mannerheim, 1844

Pearce: no records
First record: 2009
Number of records: 1
Latest record: 2009

Haliplus (Haliplus) ruficollis (De Geer, 1774)

Pearce: 6 localities

First subsequent record: 1947

Number of subsequent records: many

Latest record: 2021

Haliplus (Haliplus) sibiricus Motschulsky, 1860

Pearce: 6 localities

First subsequent record: 1943

Number of subsequent records: 7

Latest record: 2009

Haliplus (Liaphlus) flavicollis Sturm, 1834

Pearce: 3 localities

First subsequent record: 1979

Number of subsequent records: 5

Latest record: 2014

Haliplus (Liaphlus) fulvus (Fabricius, 1801)

Pearce: 4 localities

First subsequent record: 1937

Number of subsequent records: many

Latest record: 2014

Haliplus (Liaphlus) laminatus (Schaller, 1783)

Pearce: 4 localities

First subsequent record: 2009

Number of subsequent records: 1

Latest record: 2009

Haliplus (Neohalipus) lineatocollis (Marsham, 1802)

Pearce: 7 localities

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2020

4. Family NOTERIDAE Thomson, C. G., 1860

Noterus clavicornis (De Geer, 1774)

Pearce: 6 localities

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2020

5. Family HYGROBIIDAE Régimbart, 1879 (1837)

Hygrobia hermanni (Fabricius, 1775)

Pearce: 3 localities

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2020

6. Family DYTISCIDAE Leach, 1815

Agabus (Acatodes) congener (Thunberg, 1794)

Pearce: no records

First record: 2019

Number of records: 1

Latest record: 2019

Agabus (Acatodes) sturmii (Gyllenhal in Schönherr, 1808)

Pearce: 4 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2019

Agabus (Agabus) labiatus (Brahm, 1790)

Pearce: 3 localities

First subsequent record: 1934

Number of subsequent records: 7

Latest record: 2008

Agabus (Agabus) uliginosus (Linnaeus, 1760)

Pearce: 6 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2011

Agabus (Gaurodytes) affinis (Paykull, 1798)

Pearce: no records

First record: 1947

Number of records: many

Latest record: 2019

Agabus (Gaurodytes) biguttatus (Olivier, 1795)

Pearce: 2 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2015

Agabus (Gaurodytes) bipustulatus
(Linnaeus, 1767)
Pearce: generally common and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Agabus (Gaurodytes) brunneus (Fabricius, 1798)
Pearce: 2 localities
First subsequent record: 1993
Number of subsequent records: 5
Latest record: 2007

Agabus (Gaurodytes) conspersus
(Marsham, 1802)
Pearce: 1 locality
First subsequent record: 1964
Number of subsequent records: 4
Latest record: 1981

Agabus (Gaurodytes) didymus (Olivier, 1795)
Pearce: 6 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Agabus (Gaurodytes) guttatus (Paykull, 1798)
Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2013

Agabus (Gaurodytes) melanarius Aubé, 1837
Pearce: no records
First record: 1987
Number of records: many
Latest record: 2013

Agabus (Gaurodytes) nebulosus (Forster, 1771)
Pearce: 4 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2020

Agabus (Gaurodytes) paludosus
(Fabricius, 1801)
Pearce: 6 localities
First subsequent record: 1943
Number of subsequent records: many
Latest record: 2021

Agabus (Gaurodytes) unguicularis
(Thomson, C.G., 1867)
Pearce: 1 locality
First subsequent record: 2001
Number of subsequent records: 1
Latest record: 2001

Ilybius aenescens Thomson, C.G., 1870
Pearce: 2 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2020

Ilybius ater (De Geer, 1774)
Pearce: 2 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2016

Ilybius chalconatus (Panzer, 1796)
Pearce: 4 localities (but probably including *montanus*)
First subsequent record: 1934
Number of subsequent records: 9
Latest record: 2013

Ilybius fenestratus (Fabricius, 1781)
Pearce: 1 locality
First subsequent record: 1993
Number of subsequent records: 4
Latest record: 2006

Ilybius fuliginosus (Fabricius, 1792)
Pearce: 5 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2021

Ilybius guttiger (Gyllenhal, 1808)
Pearce: no records
First record: 1979
Number of records: many
Latest record: 2017

Ilybius montanus (Stephens, 1828)
Pearce: no records (but see *chalconatus* above)
First record: 1934
Number of records: many
Latest record: 2021

Ilybius quadriguttatus (Lacordaire, 1835)
Pearce: no records
First record: 1947
Number of records: many
Latest record: 2020

Platambus maculatus (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2021

Colymbetes fuscus (Linnaeus, 1758)

Pearce: generally common and widely distributed
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2020

Nartus grapii (Gyllenhal, 1808) [2012

Rhantus grapii]

Pearce: no records
First record: 1937
Number of records: many
Latest record: 2016

Rhantus exsoletus (Forster, 1771)

Pearce: "Dorset" (A. Ford)
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2014

Rhantus frontalis (Marshall, 1802)

Pearce: 1 locality
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2006

Rhantus suturalis (MacLeay, 1825)

Pearce: 2 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2019

Rhantus suturellus (Harris, 1828)

Pearce: "Dorset" (A. Ford)
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2014

Liopterus haemorrhoidalis (Fabricius, 1787)

Pearce: 3 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2021

Laccophilus hyalinus (De Geer, 1774)

Pearce: 5 localities
First subsequent record: 1947
Number of subsequent records: 8
Latest record: 1993

Laccophilus minutus (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Dytiscus circumcinctus Ahrens, 1811

Pearce: no records
First record: 1964
Number of records: 1
Latest record: 1964

Dytiscus circumflexus Fabricius, 1801

Pearce: no records
First record: 1937
Number of records: 4
Latest record: 2007

Dytiscus marginalis Linnaeus, 1758

Pearce: 7 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2021

Dytiscus semisulcatus Müller, O.F., 1776

Pearce: 7 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2014

Hydaticus seminiger (De Geer, 1774)

Pearce: no records
First record: 1947
Number of records: many
Latest record: 2020

Acilius sulcatus (Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2018

Graphoderus cinereus (Linnaeus, 1758)

Pearce: no records
First record: 1984
Number of records: many
Latest record: 2015

Hydroporus angustatus Sturm, 1835

Pearce: 1 locality
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2020

Hydroporus discretus Fairmaire & Brisout de Barneville, C., 1859
Pearce: 2 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2019

Hydroporus dorsalis s.l. [2012
Suphrodytes dorsalis]
Pearce: 2 localities
First subsequent record: 1980
Number of subsequent records: 4
Latest record: 2005

Hydroporus erythrocephalus (Linnaeus, 1758)
Pearce: 4 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2020

Hydroporus ferrugineus Stephens, 1829
Pearce: no records
First record: 1933
Number of records: many
Latest record: 2006

Hydroporus figuratus (Gyllenhal in Sahlberg, C.R., 1826)
Pearce: no records
First record: 2005
Number of records: 2
Latest record: 2005

Hydroporus gyllenhalii Schiødte, 1841
Pearce: 8 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Hydroporus incognitus Sharp, 1869
Pearce: no records
First record: 1947
Number of records: many
Latest record: 2019

Hydroporus longicornis Sharp, 1871
Pearce: no records
First record: 2017
Number of records: 1
Latest record: 2017

Hydroporus longulus Mulsant & Rey, 1861
Pearce: no records
First record: 1966
Number of records: 3
Latest record: 2005

Hydroporus marginatus (Duftschmid, 1805)
Pearce: 1 locality
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2004

Hydroporus melanarius Sturm, 1835
Pearce: 1 locality
First subsequent record: 2005
Number of subsequent records: 3
Latest record: 2006

Hydroporus memnonius Nicolai, 1822
Pearce: 6 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Hydroporus necopinatus Fery, 1999
Pearce: no records
First record: 1937
Number of records: many
Latest record: 2019

Hydroporus neglectus Schaum, 1845
Pearce: no records
First record: 1964
Number of records: many
Latest record: 2019

Hydroporus nigrita (Fabricius, 1792)
Pearce: 1 locality
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2019

Hydroporus obscurus Sturm, 1835
Pearce: 3 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2020

Hydroporus palustris (Linnaeus, 1760)
Pearce: generally common and widely distributed
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2020

Hydroporus planus (Fabricius, 1782)

Pearce: 5 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Hydroporus pubescens (Gyllenhal, 1808)

Pearce: generally common and widely distributed
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Hydroporus scalseianus Stephens, 1828

Pearce: no records
First record: 1997
Number of records: 2
Latest record: 1998

Hydroporus striola (Gyllenhal in Sahlberg, C. R., 1826)

Pearce: 1 locality
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2013

Hydroporus tessellatus (Drapiez, 1819)

Pearce: 8 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Hydroporus tristis (Paykull, 1798)

Pearce: 2 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2019

Hydroporus umbrosus (Gyllenhal, 1808)

Pearce: 2 localities (1890, 1923)
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2019

Deronectes latus (Stephens, 1829)

Pearce: 1 locality (1892)
First subsequent record: 1968
Number of subsequent records: 4
Latest record: 2009

Nebrioporus elegans (Panzer, 1794)

Pearce: 7 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Oreodytes sanmarkii (Sahlberg, C. R., 1826)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2015

Stictotarsus duodecimpustulatus (Fabricius, 1792)

Pearce: 6 localities
First subsequent record: 1962
Number of subsequent records: many
Latest record: 2007

Graptodytes bilineatus (Sturm, 1835)

Pearce: no records
First record: 1981
Number of records: 8
Latest record: 2019

Graptodytes flavipes (Olivier, 1795)

Pearce: 2 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2021

Graptodytes granularis (Linnaeus, 1767)

Pearce: 2 localities (1892)
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2020

Graptodytes pictus (Fabricius, 1787)

Pearce: 7 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2020

Porhydrus lineatus (Fabricius, 1775)

Pearce: 2 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2014

Stictonectes lepidus (Olivier, 1795)

Pearce: 7 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2018

Hydrovatus clypealis Sharp, 1876

Pearce: 1 locality
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2014

Hygrotus (Coelambus) confluens
(Fabricius, 1787)

Pearce: 3 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2019

Hygrotus (Hygrotus) inaequalis
(Fabricius, 1777)

Pearce: 3 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2021

Hygrotus (Hygrotus) versicolor (Schaller, 1783)

Pearce: 3 localities
First subsequent record: 1966
Number of subsequent records: 3
Latest record: 2006

Hygrotus (Leptolambus) impressopunctatus (Schaller, 1783)

Pearce: no records
First record: 1981
Number of records: many
Latest record: 2020

Hygrotus (Leptolambus) novemlineatus (Stephens, 1829)

Pearce: 1 locality
No subsequent records

Hygrotus (Leptolambus) parallelogrammus (Ahrens, 1812)

Pearce: 2 localities
First subsequent record: 1997
Number of subsequent records: 1
Latest record: 1997

Hyphydrus ovatus (Linnaeus, 1760)

Pearce: 6 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2020

Bidessus unistriatus (Goeze, 1777)

Pearce: 2 localities
First subsequent record: 1942
Number of subsequent records: 1
Latest record: 1942

Hydroglyphus geminus (Fabricius, 1792)

Pearce: no records
First record: 1964
Number of records: many
Latest record: 2019

7. Family CARABIDAE Latreille, 1802

Leistus (Leistus) ferrugineus (Linnaeus, 1758)

Pearce: widely distributed
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2019

Leistus (Leistus) fulvibarbis Dejean, 1826

Pearce: 3 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Leistus (Leistus) terminatus (Panzer, 1793)

Pearce: 2 localities
First subsequent record: 1976
Number of subsequent records: 3
Latest record: 2015

Leistus (Pogonophorus) rufomarginatus (Duftschmid, 1812)

Pearce: no records
First record: 1965
Number of records: many
Latest record: 2018

Leistus (Pogonophorus) spinibarbis (Fabricius, 1775)

Pearce: common and widely distributed
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Nebria brevicollis (Fabricius, 1792)

Pearce: abundant (but probably including *salina*)
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Nebria salina Fairmaire & Laboulbène, 1854

Pearce: no records (but see *brevicollis* above)
First record: 1935
Number of records: many
Latest record: 2021

Notiophilus aquaticus (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2019

Notiophilus biguttatus (Fabricius, 1779)

Pearce: generally common and widely distributed
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Notiophilus germinyi Fauvel, 1863

Pearce: 1 locality
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Notiophilus palustris (Duftschmid, 1812)

Pearce: widely distributed
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2019

Notiophilus quadripunctatus Dejean, 1826

Pearce: 1 locality (1890)
First subsequent record: 1978
Number of subsequent records: 2
Latest record: 1982

Notiophilus rufipes Curtis, 1829

Pearce: 1 locality (1911)
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2020

Notiophilus substriatus Waterhouse, G.R., 1833

Pearce: 8 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2019

Cychrus caraboides (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2019

Calosoma sycophanta (Linnaeus, 1758)

Pearce: 1 locality
First subsequent record: 2018
Number of subsequent records: 1
Latest record: 2018

Carabus (Archicarabus) nemoralis Müller, O.F., 1764

Pearce: generally common
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2020

Carabus (Carabus) arvensis Herbst, 1784

Pearce: 4 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2020

Carabus (Carabus) granulatus Linnaeus, 1758

Pearce: very common
First subsequent record: 1946
Number of subsequent records: many
Latest record: 2021

Carabus (Hemicarabus) nitens Linnaeus, 1758

Pearce: common on swampy heaths
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2012

Carabus (Megodontus) violaceus Linnaeus, 1758

Pearce: generally common
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Carabus (Mesocarabus) problematicus Herbst, 1786

Pearce: 4 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Carabus (Morphocarabus) monilis Fabricius, 1792

Pearce: 3 localities
First subsequent record: 1980
Number of subsequent records: 8
Latest record: 2019

Cicindela campestris Linnaeus, 1758

Pearce: common and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Cicindela maritima Dejean in Latreille & Dejean, 1822

Pearce: 1 locality
First subsequent record: 1964
Number of subsequent records: 1
Latest record: 1964

Cicindela sylvatica Linnaeus, 1758

Pearce: 3 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Cylindera germanica (Linnaeus, 1758)

Pearce: widely distributed on coast
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Loricera pilicornis (Fabricius, 1775)

Pearce: abundant
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Blethisa multipunctata (Linnaeus, 1758)

Pearce: 2 localities (1896, 1925)
First subsequent record: 1977
Number of subsequent records: 7
Latest record: 2017

Elaphrus (Elaphrus) riparius (Linnaeus, 1758)

Pearce: common
First subsequent record: 1940
Number of subsequent records: many
Latest record: 2020

Elaphrus (Neoelaphrus) cupreus
Duftschmid, 1812

Pearce: 4 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2019

Elaphrus (Neoelaphrus) uliginosus
Fabricius, 1792

Pearce: 1 locality
First subsequent record: 1955
Number of subsequent records: 9
Latest record: 2018

Clivina collaris (Herbst, 1784)

Pearce: 2 localities
First subsequent record: 1967
Number of subsequent records: 2
Latest record: 1989

Clivina fossor (Linnaeus, 1758)

Pearce: not common but widely distributed
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Dyschirius (Dyschiriodes) aeneus (Dejean, 1825)

Pearce: 2 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Dyschirius (Dyschiriodes) impunctipennis
Dawson, 1854

Pearce: 2 localities
No subsequent records

Dyschirius (Dyschiriodes) nitidus (Dejean, 1825)

Pearce: no records
First record: 1964
Number of records: 1
Latest record: 1964

Dyschirius (Dyschiriodes) politus (Dejean, 1825)

Pearce: 1 locality
First subsequent record: 1968
Number of subsequent records: 7
Latest record: 2014

Dyschirius (Dyschiriodes) salinus
Schaum, 1843

Pearce: 2 localities
First subsequent record: 1989
Number of subsequent records: 5
Latest record: 2013

Dyschirius (Dyschiriodes) tristis Stephens, 1827 [2012 *Dyschirius luedersi*]

Pearce: 1 locality
First subsequent record: 1991
Number of subsequent records: many
Latest record: 2014

Dyschirius (Dyschirius) thoracicus (Rossi, 1790)

Pearce: 2 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2019

Dyschirius (Eudyschirius) globosus (Herbst, 1784)

Pearce: 4 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Broscus cephalotes (Linnaeus, 1758)

Pearce: 7 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2018

Aepopsis robinii (Laboulbène, 1849) [2012
Aepus robinii]

Pearce: 3 localities
No subsequent records

Aepus marinus (Strøm, 1783)

Pearce: 2 localities
First subsequent record: 1981
Number of subsequent records: 7
Latest record: 2010

Trechoblemus micros (Herbst, 1784)

Pearce: no records
First record: 1967
Number of records: 3
Latest record: 1989

Trechus (Epaphius) secalis (Paykull, 1790)

Pearce: 2 localities
First subsequent record: 1986
Number of subsequent records: 1
Latest record: 1986

Trechus (Trechus) fulvus Dejean, 1831

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: 2
Latest record: 1954

Trechus (Trechus) obtusus Erichson, 1837

Pearce: 2 localities
First subsequent record: 1943
Number of subsequent records: many
Latest record: 2019

Trechus (Trechus) quadristriatus (Schrank, 1781)

Pearce: generally common and widely distributed
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2021

Trechus (Trechus) rubens (Fabricius, 1792)

Pearce: 1 locality
First subsequent record: 1972
Number of subsequent records: 1
Latest record: 1972

Asaphidion curtum (Heyden, 1870)

Pearce: no records
First record: 1938
Number of records: many
Latest record: 2021

Asaphidion flavipes (Linnaeus, 1760)

Pearce: 8 localities (*sensu lato*)
First subsequent record: 1988
Number of subsequent records: many
Latest record: 2021

Asaphidion pallipes (Duftschmid, 1812)

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2010

Asaphidion stierlini (Heyden, 1880)

Pearce: no records
First record: 2002
Number of records: 5
Latest record: 2021

Bembidion (Actedium) pallidipenne (Illiger, 1802)

Pearce: 1 locality (1910)
No subsequent records

Bembidion (Bembidion) quadrimaculatum (Linnaeus, 1760)

Pearce: 6 localities
First subsequent record: 1940
Number of subsequent records: many
Latest record: 2021

Bembidion (Bembidion) quadripustulatum Audinet-Serville, 1821

Pearce: 7 localities
First subsequent record: 1993
Number of subsequent records: 3
Latest record: 2013

Bembidion (Bembidionetolitzkya)
atrocaeruleum (Stephens, 1828)

Pearce: no records
First record: 1988
Number of records: 4
Latest record: 2009

Bembidion (Bembidionetolitzkya) tibiale
(Duftschmid, 1812)

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Bembidion (Diplocampa) assimile
Gyllenhal, 1810

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Bembidion (Diplocampa) clarkii (Dawson,
1849)

Pearce: 1 locality (1852, 1896)
First subsequent record: 1948
Number of subsequent records: 5
Latest record: 2018

Bembidion (Diplocampa) fumigatum
(Duftschmid, 1812)

Pearce: no records
First record: 1963
Number of records: many
Latest record: 2020

Bembidion (Emphanes) minimum
(Fabricius, 1792)

Pearce: 3 localities
First subsequent record: 1943
Number of subsequent records: 5
Latest record: 2005

Bembidion (Emphanes) normannum
Dejean, 1831

Pearce: 3 localities
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2018

Bembidion (Eupetodromus) dentellum
(Thunberg, 1787)

Pearce: "Dorset" (A. Ford)
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2020

Bembidion (Lymnaeum) nigropiceum
(Marsham, 1802)

Pearce: 5 localities
First subsequent record: 1986
Number of subsequent records: 3
Latest record: 1999

Bembidion (Metallina) lampros (Herbst,
1784)

Pearce: generally common and widely distributed
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2021

Bembidion (Metallina) properans
(Stephens, 1828)

Pearce: no records
First record: 1935
Number of records: many
Latest record: 2021

Bembidion (Neja) nigricorne Gyllenhal,
1827

Pearce: 1 locality (1925)
First subsequent record: 1935
Number of subsequent records: many
Latest record: 1981

Bembidion (Nepha) genei Küster, 1847
[2012 *Bembidion illigeri*]

Pearce: no records
First record: 1957
Number of records: many
Latest record: 2020

Bembidion (Notaphemphanes) ephippium
(Marsham, 1802)

Pearce: 1 locality
First subsequent record: 1939
Number of subsequent records: 1
Latest record: 1939

Bembidion (Notaphus) varium (Olivier,
1795)

Pearce: 1 locality
First subsequent record: 1943
Number of subsequent records: many
Latest record: 2021

Bembidion (Ocydromus) decorum (Zenker
in Panzer, 1799)

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: 1
Latest record: 1979

Bembidion (Ocydromus) saxatile
Gyllenhal, 1827
Pearce: 7 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2018

Bembidion (Peryphanes) deletum Audinet-Serville, 1821
Pearce: 6 localities
First subsequent record: 1951
Number of subsequent records: many
Latest record: 2018

Bembidion (Peryphanes) stephensii
Crotch, 1866
Pearce: 4 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2016

Bembidion (Peryphus) bruxellense
Wesmael, 1835
Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: 5
Latest record: 2019

Bembidion (Peryphus) bualei Jacquelin du Val, 1852
Pearce: 1 locality
First subsequent record: 2000
Number of subsequent records: 1
Latest record: 2000

Bembidion (Peryphus) femoratum Sturm, 1825
Pearce: no records
First record: 1966
Number of records: 8
Latest record: 2005

Bembidion (Peryphus) maritimum (Stephens, 1835)
Pearce: "Dorset" (A. Ford)
First subsequent record: 1975
Number of subsequent records: 1
Latest record: 1975

Bembidion (Peryphus) tetracolum Say, 1823
Pearce: abundant and widely distributed
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2020

Bembidion (Philochthus) aeneum Germar, 1823
Pearce: 6 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2019

Bembidion (Philochthus) biguttatum (Fabricius, 1779)
Pearce: abundant and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Bembidion (Philochthus) guttula (Fabricius, 1792)
Pearce: common and widely distributed
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2021

Bembidion (Philochthus) iricolor Bedel, 1879
Pearce: 1 locality
First subsequent record: 2004
Number of subsequent records: 1
Latest record: 2004

Bembidion (Philochthus) lunulatum (Geoffroy in Fourcroy, 1785)
Pearce: 5 localities
First subsequent record: 1952
Number of subsequent records: many
Latest record: 2021

Bembidion (Philochthus) mannerheimii Sahlberg, C.R., 1827
Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Bembidion (Phyla) obtusum Audinet-Serville, 1821
Pearce: 8 localities
First subsequent record: 1949
Number of subsequent records: many
Latest record: 2021

Bembidion (Princidium) punctulatum Drapiez, 1820
Pearce: no records
First record: 1999
Number of records: 1
Latest record: 1999

Bembidion (Semicampa) gilvipes Sturm, 1825

Pearce: 2 localities

First subsequent record: 2006

Number of subsequent records: 3

Latest record: 2012

Bembidion (Trepanedoris) doris (Panzer, 1796)

Pearce: 2 localities

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2020

Bembidion (Trepanes) articulatum (Panzer, 1796)

Pearce: 2 localities

First subsequent record: 1943

Number of subsequent records: many

Latest record: 2020

Bracteon litorale (Olivier, 1790)

Pearce: no records

First record: 1914

Number of records: 2

Latest record: 1989

Cillenus lateralis Samouelle, 1819

Pearce: 2 localities

First subsequent record: 1954

Number of subsequent records: 5

Latest record: 2019

Ocys harpaloides (Audinet-Serville, 1821)

Pearce: 7 localities (but probably including (or all) *tachysoides*)

No subsequent *sensu stricto* records

Ocys quinquestriatus (Gyllenhal, 1810)

Pearce: 1 locality

First subsequent record: 1986

Number of subsequent records: 5

Latest record: 2004

Ocys tachysoides (Antoine, 1933)

Pearce: no records (but see *harpaloides* above)

First record: 2014

Number of records: 6

Latest record: 2021

Sinechostictus stomoides (Dejean, 1831)

[2012 *Bembidion (Sinechostictus) stomoides*]

Pearce: no records

First record: 1989

Number of records: 1

Latest record: 1989

Tachys (Paratachys) bistriatus (Duftschmid, 1812)

Pearce: 3 localities

First subsequent record: 1968

Number of subsequent records: many

Latest record: 2021

Tachys (Paratachys) micros (Fischer von Waldheim, 1828)

Pearce: 2 localities

First subsequent record: 1947

Number of subsequent records: many

Latest record: 2017

Tachys (Tachys) scutellaris Stephens, 1828

Pearce: no records

First record: 1938

Number of records: 6

Latest record: 2006

Tachyura parvula (Dejean, 1831) [2012 *Elaphropus parvulus*]

Pearce: no records

First record: 2004

Number of records: 3

Latest record: 2017

Tachyura walkeriana (Sharp, 1913) [2012 *Elaphropus walkerianus*]

Pearce: no records

First record: 2019

Number of records: 1

Latest record: 2019

Pogonus chalceus (Marsham, 1802)

Pearce: 5 localities

First subsequent record: 1938

Number of subsequent records: many

Latest record: 2021

Pogonus littoralis (Duftschmid, 1812)

Pearce: 2 localities

First subsequent record: 2013

Number of subsequent records: 3

Latest record: 2018

Pogonus luridipennis (Germar, 1822)

Pearce: 2 localities
First subsequent record: 1996
Number of subsequent records: 1
Latest record: 1996

Patrobus atrorufus (Strøm, 1768)

Pearce: 1 locality
First subsequent record: 1986
Number of subsequent records: 7
Latest record: 2013

Brachinus crepitans (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Abax parallelepipedus (Piller & Mitterpacher, 1783)

Pearce: generally common and widely distributed
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2019

Pedius longicollis (Duftschmid, 1812)
[2012 *Pterostichus* (*Pedius*) *longicollis*]

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: 7
Latest record: 2019

Poecilus cupreus (Linnaeus, 1758)

Pearce: 7 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Poecilus kugelanni (Panzer, 1796)

Pearce: 4 localities
First subsequent record: 2001
Number of subsequent records: 8
Latest record: 2003

Poecilus lepidus (Leske, 1785)

Pearce: 3 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2019

Poecilus versicolor (Sturm, 1824)

Pearce: 4 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Pterostichus (*Adelosia*) *macer* (Marsham, 1802)

Pearce: 3 localities
First subsequent record: 1939
Number of subsequent records: 8
Latest record: 2016

Pterostichus (*Argutor*) *vernalis* (Panzer, 1796)

Pearce: 3 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Pterostichus (*Bothriopterus*) *oblongopunctatus* (Fabricius, 1787)

Pearce: 4 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2016

Pterostichus (*Bothriopterus*) *quadrioveolatus* Letzner, 1852

Pearce: no records
First record: 2015
Number of records: 1
Latest record: 2015

Pterostichus (*Morphnosoma*) *melanarius* (Illiger, 1798)

Pearce: generally common
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2019

Pterostichus (*Phonias*) *diligens* (Sturm, 1824)

Pearce: 2 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Pterostichus (*Phonias*) *strenuus* (Panzer, 1796)

Pearce: 9 localities
First subsequent record: 1946
Number of subsequent records: many
Latest record: 2021

Pterostichus (*Platysma*) *niger* (Schaller, 1783)

Pearce: generally common
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Pterostichus (Pseudomaseus) anthracinus
(Illiger, 1798)
Pearce: 3 localities
First subsequent record: 1999
Number of subsequent records: many
Latest record: 2021

Pterostichus (Pseudomaseus) gracilis
(Dejean, 1828)
Pearce: 2 localities
First subsequent record: 1940
Number of subsequent records: 8
Latest record: 2011

Pterostichus (Pseudomaseus) minor
(Gyllenhal, 1827)
Pearce: 5 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2020

Pterostichus (Pseudomaseus) nigrita
(Paykull, 1790)
Pearce: 12 localities (but probably including
rhaeticus)
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2019

Pterostichus (Pseudomaseus) rhaeticus
Heer, 1837
Pearce: no records (but see *nigrita* above)
First record: 1994
Number of records: many
Latest record: 2021

Pterostichus (Steropus) madidus
(Fabricius, 1775)
Pearce: abundant
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Stomis pumicatus (Panzer, 1796)
Pearce: 5 localities
First subsequent record: 1952
Number of subsequent records: many
Latest record: 2019

Amara (Amara) aenea (De Geer, 1774)
Pearce: generally common
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Amara (Amara) communis (Panzer, 1797)
Pearce: 5 localities
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2021

Amara (Amara) convexior Stephens, 1828
Pearce: 8 localities
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2020

Amara (Amara) eurynota (Panzer, 1796)
Pearce: 1 locality
First subsequent record: 1990
Number of subsequent records: many
Latest record: 2021

Amara (Amara) famelica Zimmermann,
C., 1832
Pearce: no records
First record: 1977
Number of records: 2
Latest record: 1977

Amara (Amara) familiaris (Duftschmid,
1812)
Pearce: 4 localities
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2021

Amara (Amara) lucida (Duftschmid, 1812)
Pearce: 3 localities
First subsequent record: 1953
Number of subsequent records: 2
Latest record: 1999

Amara (Amara) lunicollis Schiødte, 1837
Pearce: 2 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2020

Amara (Amara) montivaga Sturm, 1825
Pearce: no records
First record: 1994
Number of records: 1
Latest record: 1994

Amara (Amara) nitida Sturm, 1825
Pearce: 1 locality
No subsequent records

Amara (Amara) ovata (Fabricius, 1792)

Pearce: 5 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Amara (Amara) similata (Gyllenhal, 1810)

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2021

Amara (Amara) spreta Dejean, 1831

Pearce: 1 locality
No subsequent records

Amara (Amara) tibialis (Paykull, 1798)

Pearce: 2 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Amara (Bradytus) apricaria (Paykull, 1790)

Pearce: 5 localities
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2019

Amara (Bradytus) consularis (Duftschmid, 1812)

Pearce: 1 locality
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2004

Amara (Bradytus) fulva (Müller, O.F., 1776)

Pearce: 3 localities
First subsequent record: 2012
Number of subsequent records: 1
Latest record: 2012

Amara (Celia) bifrons (Gyllenhal, 1810)

Pearce: 3 localities
First subsequent record: 2002
Number of subsequent records: 1
Latest record: 2002

Amara (Celia) infima (Duftschmid, 1812)

Pearce: 1 locality
No subsequent records

Amara (Percosia) equestris (Duftschmid, 1812)

Pearce: 2 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Amara (Zezea) plebeja (Gyllenhal, 1810)

Pearce: 2 localities
First subsequent record: 1957
Number of subsequent records: many
Latest record: 2021

Curtonotus aulicus (Panzer, 1796)

Pearce: 3 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2021

Curtonotus convexiusculus (Marsham, 1802)

Pearce: 4 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2019

Zabrus tenebrioides (Goeze, 1777)

Pearce: 5 localities
First subsequent record: 1980
Number of subsequent records: 4
Latest record: 2021

Oodes helopioides (Fabricius, 1792)

Pearce: 3 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2020

Panagaeus bipustulatus (Fabricius, 1775)

Pearce: no records
First record: 1941
Number of records: 2
Latest record: 1982

Chlaenius nigricornis (Fabricius, 1787)

Pearce: 4 localities
First subsequent record: 1940
Number of subsequent records: many
Latest record: 2017

Chlaenius nitidulus (Schränk, 1781)

Pearce: 2 localities
First subsequent record: 1930
Number of subsequent records: 2
Latest record: 1933

Chlaenius vestitus (Paykull, 1790)

Pearce: 7 localities

First subsequent record: 1964

Number of subsequent records: many

Latest record: 2020

Badister (Badister) bullatus (Schränk, 1798)

Pearce: abundant and widely distributed

First subsequent record: 1936

Number of subsequent records: many

Latest record: 2021

Badister (Badister) unipustulatus Bonelli, 1813

Pearce: 1 locality

First subsequent record: 1934

Number of subsequent records: 3

Latest record: 1958

Badister (Baudia) collaris Motschulsky, 1844

Pearce: no records

First record: 1954

Number of records: 3

Latest record: 2014

Badister (Baudia) dilatatus Chaudoir, 1837

Pearce: no records

First record: 1991

Number of records: many

Latest record: 2015

Badister (Baudia) peltatus (Panzer, 1796)

Pearce: 1 locality

No subsequent records

Badister (Trimorphus) sodalis (Duftschmid, 1812)

Pearce: 1 locality

First subsequent record: 1946

Number of subsequent records: many

Latest record: 2018

Licinus depressus (Paykull, 1790)

Pearce: no records

First record: 1963

Number of records: many

Latest record: 2011

Licinus punctatulus (Fabricius, 1792)

Pearce: 6 localities

First subsequent record: 1937

Number of subsequent records: many

Latest record: 2018

Anisodactylus (Anisodactylus) binotatus (Fabricius, 1787)

Pearce: 6 localities

First subsequent record: 1938

Number of subsequent records: many

Latest record: 2021

Anisodactylus (Anisodactylus) nemorivagus (Duftschmid, 1812)

Pearce: common on heaths

First subsequent record: 2001

Number of subsequent records: many

Latest record: 2019

Anisodactylus (Hexatrichus) poeciloides (Stephens, 1828)

Pearce: 1 locality

No subsequent records

Scybalicus oblongiusculus (Dejean, 1829)

Pearce: 3 localities

First subsequent record: 2019

Number of subsequent records: 2

Latest record: 2019

Harpalus (Cryptophonus) melancholicus Dejean, 1829

Pearce: 1 locality (1885, 1894)

No subsequent records

Harpalus (Cryptophonus) tenebrosus Dejean, 1829

Pearce: no records

First record: 1979

Number of records: 7

Latest record: 2003

Harpalus (Harpalus) affinis (Schränk, 1781)

Pearce: generally common and widely distributed

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2021

Harpalus (Harpalus) anxius (Duftschmid, 1812)

Pearce: 3 localities

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2018

Harpalus (Harpalus) attenuatus Stephens, 1828

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2019

Harpalus (Harpalus) dimidiatus (Rossi, 1790)

Pearce: 7 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2019

Harpalus (Harpalus) froelichii Sturm, 1818

Pearce: 3 localities
No subsequent records

Harpalus (Harpalus) latus (Linnaeus, 1758)

Pearce: 8 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2021

Harpalus (Harpalus) neglectus Audinet-Serville, 1821

Pearce: 10 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2018

Harpalus (Harpalus) pumilus Sturm, 1818

Pearce: 2 localities
First subsequent record: 1990
Number of subsequent records: 1
Latest record: 1990

Harpalus (Harpalus) rubripes (Duftschmid, 1812)

Pearce: widely distributed in south of county
First subsequent record: 1950
Number of subsequent records: many
Latest record: 2019

Harpalus (Harpalus) rufipalpis Sturm, 1818

Pearce: no records
First record: 1937
Number of records: many
Latest record: 2020

Harpalus (Harpalus) serripes (Quensel in Schönherr, 1806)

Pearce: 3 localities
First subsequent record: 1972
Number of subsequent records: many
Latest record: 2005

Harpalus (Harpalus) smaragdinus (Duftschmid, 1812)

Pearce: 1 locality (1914)
First subsequent record: 1954
Number of subsequent records: 3
Latest record: 2018

Harpalus (Harpalus) tardus (Panzer, 1796)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2015

Harpalus (Pseudoophonus) rufipes (De Geer, 1774)

Pearce: abundant
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2021

Ophonus (Hesperophonus) azureus (Fabricius, 1775)

Pearce: 5 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2013

Ophonus (Metophonus) cordatus (Duftschmid, 1812)

Pearce: 1 locality
First subsequent record: 1953
Number of subsequent records: 2
Latest record: 1953

Ophonus (Metophonus) laticollis Mannerheim, 1825

Pearce: 4 localities
No subsequent records

Ophonus (Metophonus) melletii (Heer, 1837)

Pearce: 3 localities
No subsequent records

Ophonus (Metophonus) parallelus
(Dejean, 1829)

Pearce: 2 localities

First subsequent record: 1953

Number of subsequent records: 1

Latest record: 1953

Ophonus (Metophonus) puncticeps
Stephens, 1828

Pearce: 2 localities

First subsequent record: 1936

Number of subsequent records: many

Latest record: 2019

Ophonus (Metophonus) puncticollis
(Paykull, 1798)

Pearce: 6 localities

First subsequent record: 1936

Number of subsequent records: 1

Latest record: 1936

Ophonus (Metophonus) rufibarbis
(Fabricius, 1792)

Pearce: 2 localities

First subsequent record: 1982

Number of subsequent records: many

Latest record: 2014

Ophonus (Metophonus) rupicola (Sturm, 1818)

Pearce: 5 localities

First subsequent record: 1953

Number of subsequent records: 7

Latest record: 2005

Ophonus (Metophonus) schaubergerianus
(Puel, 1937)

Pearce: 1 locality

First subsequent record: 2005

Number of subsequent records: 1

Latest record: 2005

Ophonus (Metophonus) subsinuatus Rey, 1886

Pearce: no records

First record: 1886

Number of records: 2

Latest record: 1886

Ophonus (Ophonus) ardosiacus (Lutshnik, 1922)

Pearce: 5 localities

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2020

Ophonus (Ophonus) sabulicola (Panzer, 1796)

Pearce: 4 localities

First subsequent record: 1947

Number of subsequent records: 1

Latest record: 1947

Acupalpus brunnipes (Sturm, 1825)

Pearce: 2 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2017

Acupalpus dubius Schilsky, 1888

Pearce: 4 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2021

Acupalpus exiguus Dejean, 1829

Pearce: 6 localities

First subsequent record: 1994

Number of subsequent records: many

Latest record: 2021

Acupalpus flavicollis (Sturm, 1825)

Pearce: 1 locality (1886)

First subsequent record: 1995

Number of subsequent records: 8

Latest record: 2018

Acupalpus meridianus (Linnaeus, 1760)

Pearce: 3 localities

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2013

Acupalpus parvulus (Sturm, 1825)

Pearce: 3 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2020

Anthracus consputus (Duftschmid, 1812)

Pearce: 3 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2020

Bradycellus caucasicus (Chaudoir, 1846)

Pearce: no records

First record: 1977

Number of records: 4

Latest record: 2009

Bradycellus distinctus (Dejean, 1829)

Pearce: 3 localities
No subsequent records

Bradycellus harpalinus (Audinet-Serville, 1821)

Pearce: 3 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Bradycellus ruficollis (Stephens, 1828)

Pearce: 4 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Bradycellus sharpi Joy, 1912

Pearce: 1 locality
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2020

Bradycellus verbasci (Duftschmid, 1812)

Pearce: 5 localities
First subsequent record: 1962
Number of subsequent records: many
Latest record: 2019

Dicheirotrichus (Dicheirotrichus) gustavii Crotch, 1871

Pearce: 5 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2015

Dicheirotrichus (Dicheirotrichus) obsoletus (Dejean, 1829)

Pearce: 5 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2019

Dicheirotrichus (Trichocellus) placidus (Gyllenhal, 1827)

Pearce: 1 locality
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Stenolophus mixtus (Herbst, 1784)

Pearce: 3 localities
First subsequent record: 1940
Number of subsequent records: many
Latest record: 2021

Stenolophus skrimshirani Stephens, 1828

Pearce: no records
First record: 1990
Number of records: many
Latest record: 2011

Stenolophus teutonus (Schränk, 1781)

Pearce: 1 locality
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2014

Platyderus depressus (Audinet-Serville, 1821)

Pearce: 4 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2019

Calathus (Amphyginus) rotundicollis Dejean, 1828

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2017

Calathus (Calathus) fuscipes (Goeze, 1777)

Pearce: abundant and widely distributed
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Calathus (Neocalathus) ambiguus (Paykull, 1790)

Pearce: 4 localities
First subsequent record: 1953
Number of subsequent records: 1
Latest record: 1953

Calathus (Neocalathus) cinctus Motschulsky, 1850

Pearce: no records
First record: 1935
Number of records: many
Latest record: 2021

Calathus (Neocalathus) erratus (Sahlberg, C.R., 1827)

Pearce: 5 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Calathus (Neocalathus) melanocephalus
(Linnaeus, 1758)

Pearce: abundant and widely distributed (but may include *cinctus*)

First subsequent record: 1982

Number of subsequent records: many

Latest record: 2019

Calathus (Neocalathus) micropterus
(Duftschmid, 1812)

Pearce: no records

First record: 1999

Number of records: 1

Latest record: 1999

Calathus (Neocalathus) mollis (Marshall, 1802)

Pearce: 4 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Laemostenus (Laemostenus) complanatus
(Dejean, 1828)

Pearce: 1 locality

No subsequent records

Laemostenus (Pristonychus) terricola
(Herbst, 1784)

Pearce: 4 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2019

Synuchus vivalis (Illiger, 1798)

Pearce: 4 localities

First subsequent record: 1943

Number of subsequent records: many

Latest record: 2011

Agonum (Agonum) marginatum (Linnaeus, 1758)

Pearce: 8 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2017

Agonum (Agonum) muelleri (Herbst, 1784)

Pearce: generally common and widely distributed

First subsequent record: 1940

Number of subsequent records: many

Latest record: 2019

Agonum (Agonum) nigrum Dejean, 1828

Pearce: 1 locality

First subsequent record: 1979

Number of subsequent records: 9

Latest record: 2018

Agonum (Europhilus) fuliginosum (Panzer, 1809)

Pearce: 5 localities

First subsequent record: 1936

Number of subsequent records: many

Latest record: 2021

Agonum (Europhilus) gracile Sturm, 1824

Pearce: 4 localities

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2021

Agonum (Europhilus) micans (Nicolai, 1822)

Pearce: 4 localities

First subsequent record: 1938

Number of subsequent records: many

Latest record: 2019

Agonum (Europhilus) piceum (Linnaeus, 1758)

Pearce: 1 locality (1919)

First subsequent record: 1968

Number of subsequent records: many

Latest record: 2010

Agonum (Europhilus) scitulum Dejean, 1828

Pearce: no records

First record: 1955

Number of records: 1

Latest record: 1955

Agonum (Europhilus) thoreyi Dejean, 1828

Pearce: 1 locality (1924)

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2021

Agonum (Olisares) emarginatum
(Gyllenhal, 1827)

Pearce: generally common

First subsequent record: 1940

Number of subsequent records: many

Latest record: 2021

Agonum (Olisares) ericeti (Panzer, 1809)

Pearce: no records
First record: 1963
Number of records: 4
Latest record: 2006

Agonum (Olisares) sexpunctatum
(Linnaeus, 1758)

Pearce: 1 locality
First subsequent record: 2002
Number of subsequent records: many
Latest record: 2017

Agonum (Olisares) versutum Sturm, 1824

Pearce: 1 locality (1919)
First subsequent record: 2013
Number of subsequent records: 1
Latest record: 2013

Agonum (Olisares) viduum (Panzer, 1796)

Pearce: 3 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2019

Anchomenus dorsalis (Pontoppidan, 1763)

Pearce: abundant and widely distributed
First subsequent record: 1940
Number of subsequent records: many
Latest record: 2019

Limodromus assimilis (Paykull, 1790)

[2012 *Platynus assimilis*]
Pearce: 1 locality (1896)
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2019

Olisthopus rotundatus (Paykull, 1790)

Pearce: 5 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2019

Oxypselaphus obscurus (Herbst, 1784)

Pearce: 5 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Paranchus albipes (Fabricius, 1796)

Pearce: generally common and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Sericoda quadripunctata (De Geer, 1774)

Pearce: 1 locality
No subsequent records

Perigona nigriceps (Dejean, 1831)

Pearce: no records
First record: 1990
Number of records: 3
Latest record: 2003

Odacantha melanura (Linnaeus, 1767)

Pearce: no records
First record: 2005
Number of records: 7
Latest record: 2019

Masoreus wetterhallii (Gyllenhal, 1813)

Pearce: 2 localities
First subsequent record: 1997
Number of subsequent records: 2
Latest record: 2014

Cymindis axillaris (Fabricius, 1794)

Pearce: 5 localities
First subsequent record: 1939
Number of subsequent records: 4
Latest record: 2018

Demetrias (Aetophorus) imperialis
(Germar, 1823)

Pearce: no records
First record: 1962
Number of records: many
Latest record: 2018

Demetrias (Demetrias) atricapillus
(Linnaeus, 1758)

Pearce: abundant and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Demetrias (Demetrias) monostigma Leach
in Samouelle, 1819

Pearce: 1 locality
No subsequent records

Calodromius spilotus (Illiger, 1798)

Pearce: 5 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Dromius agilis (Fabricius, 1787)

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: 4
Latest record: 1988

Dromius meridionalis Dejean, 1825

Pearce: 3 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Dromius quadrimaculatus (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 1943
Number of subsequent records: many
Latest record: 2021

Lionychus quadrillum (Duftschmid, 1812)

Pearce: 2 localities
First subsequent record: 1953
Number of subsequent records: 2
Latest record: 2006

Microlestes maurus (Sturm, 1827)

Pearce: 10 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2021

Microlestes minutulus (Goeze, 1777)

Pearce: no records
First record: 2013
Number of records: many
Latest record: 2021

Paradromius linearis (Olivier, 1795)

Pearce: 10 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Philorhizus melanocephalus (Dejean, 1825)

Pearce: 3 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Philorhizus notatus (Stephens, 1827)

Pearce: 4 localities
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2021

Philorhizus vectensis (Rye, 1873)

Pearce: 3 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2016

Syntomus foveatus (Geoffroy in Fourcroy, 1785)

Pearce: generally common and widely distributed
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Syntomus obscuroguttatus (Duftschmid, 1812)

Pearce: 8 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2021

Syntomus truncatellus (Linnaeus, 1760)

Pearce: 3 localities
First subsequent record: 1979
Number of subsequent records: 8
Latest record: 2016

Lebia (Lamprias) chlorocephala (Hoffmann, J., 1803)

Pearce: 2 localities
First subsequent record: 1950
Number of subsequent records: 4
Latest record: 2006

Lebia (Lamprias) cyanocephala (Linnaeus, 1758)

Pearce: 1 locality
No subsequent records

Lebia (Lebia) cruxminor (Linnaeus, 1758)

Pearce: no records
First record: 2009
Number of records: 2
Latest record: 2015

Drypta dentata (Rossi, 1790)

Pearce: 1 locality
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2011

Polistichus connexus (Geoffroy in Fourcroy, 1785)

Pearce: 3 localities
First subsequent record: 1995
Number of subsequent records: 3
Latest record: 2020

8. Family HELOPHORIDAE Leach, 1815

Helophorus (Empleurus) nubilus

Fabricius, 1777

Pearce: 8 localities

First subsequent record: 1937

Number of subsequent records: 4

Latest record: 1946

Helophorus (Empleurus) porculus Bedel, 1881

Pearce: 1 locality (1904)

No subsequent records

Helophorus (Empleurus) rufipes (Bosc d'Antic, 1791)

Pearce: 3 localities

First subsequent record: 1990

Number of subsequent records: 7

Latest record: 2021

Helophorus (Helophorus) aequalis

Thomson, C.G., 1868

Pearce: 6 localities (but probably including *grandis*)

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2021

Helophorus (Helophorus) grandis Illiger, 1798

Pearce: no records (but see *aequalis* above)

First record: 1979

Number of records: many

Latest record: 2020

Helophorus (Rhopalohelophorus) arvernicus Mulsant, 1846

Pearce: no records

First record: 1995

Number of records: 3

Latest record: 2017

Helophorus (Rhopalohelophorus) brevipalpis Bedel, 1881

Pearce: 7 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Helophorus (Rhopalohelophorus) dorsalis (Marsham, 1802)

Pearce: 2 localities

First subsequent record: 2014

Number of subsequent records: 1

Latest record: 2014

Helophorus (Rhopalohelophorus) flavipes Fabricius, 1792

Pearce: generally common

First subsequent record: 1947

Number of subsequent records: many

Latest record: 2019

Helophorus (Rhopalohelophorus) fulgidicollis Motschulsky, 1860

Pearce: 1 locality

First subsequent record: 1947

Number of subsequent records: 3

Latest record: 2008

Helophorus (Rhopalohelophorus) granularis (Linnaeus, 1760)

Pearce: 3 localities

First subsequent record: 1986

Number of subsequent records: many

Latest record: 2019

Helophorus (Rhopalohelophorus) griseus Herbst, 1793

Pearce: 3 localities

First subsequent record: 1983

Number of subsequent records: many

Latest record: 2012

Helophorus (Rhopalohelophorus) laticollis Thomson, C.G., 1853

Pearce: 1 locality (1892)

No subsequent records

Helophorus (Rhopalohelophorus) longitarsis Wollaston, 1864

Pearce: no records

First record: 2002

Number of records: 3

Latest record: 2012

Helophorus (Rhopalohelophorus) minutus Fabricius, 1775

Pearce: no records

First record: 1947

Number of records: many

Latest record: 2021

Helophorus (Rhopalohelophorus) nanus

Sturm, 1836

Pearce: no records

First record: 1933

Number of records: 3

Latest record: 2021

Helophorus (Rhopalohelophorus)

obscurus Mulsant, 1844

Pearce: no records

First record: 1979

Number of records: many

Latest record: 2021

Helophorus (Trichohelophorus) alternans

Gené, 1836

Pearce: 1 locality

First subsequent record: 1937

Number of subsequent records: 3

Latest record: 2019

9. Family GEORISSIDAE Laporte, 1840

Georissus crenulatus (Rossi, 1794)

Pearce: 4 localities

First subsequent record: 1988

Number of subsequent records: many

Latest record: 2016

10. Family HYDROCHIDAE Thomson, C.G., 1859

Hydrochus angustatus Germar, 1824

Pearce: 7 localities

First subsequent record: 1937

Number of subsequent records: many

Latest record: 2020

Hydrochus elongatus (Schaller, 1783)

Pearce: 1 locality

No subsequent records

12. Family HYDROPHILIDAE Latreille, 1802

Berosus (Berosus) affinis Brullé, 1835

Pearce: 1 locality

First subsequent record: 1964

Number of subsequent records: many

Latest record: 2020

Berosus (Berosus) luridus (Linnaeus, 1760)

Pearce: 3 localities

First subsequent record: 1937

Number of subsequent records: many

Latest record: 2014

Berosus (Berosus) signaticollis (Charpentier, 1825)

Pearce: 1 locality (1909, 1925)

First subsequent record: 1937

Number of subsequent records: many

Latest record: 2020

Berosus (Enoplurus) fulvus Kuwert, 1888

Pearce: 1 locality (1896)

No subsequent records

Laccobius (Dimorpholaccobius) atratus

Rottenberg, 1874

Pearce: no records

First record: 1947

Number of records: many

Latest record: 2019

Laccobius (Dimorpholaccobius)

bipunctatus (Fabricius, 1775)

Pearce: 5 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Laccobius (Dimorpholaccobius) sinuatus

Motschulsky, 1849

Pearce: 3 localities

First subsequent record: 1980

Number of subsequent records: many

Latest record: 2011

Laccobius (Dimorpholaccobius) striatulus

(Fabricius, 1801)

Pearce: 1 locality

First subsequent record: 1933

Number of subsequent records: many

Latest record: 2019

Laccobius (Dimorpholaccobius) ytenensis

Sharp, 1910

Pearce: 1 locality

First subsequent record: 1952

Number of subsequent records: many

Latest record: 2011

Laccobius (Laccobius) colon (Stephens, 1829)

Pearce: 1 locality
First subsequent record: 1993
Number of subsequent records: 6
Latest record: 2019

Laccobius (Laccobius) minutus (Linnaeus, 1758)

Pearce: 1 locality
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2020

Paracymus scutellaris (Rosenhauer, 1856)

Pearce: 1 locality
First subsequent record: 2006
Number of subsequent records: many
Latest record: 2019

Hydrophilus piceus (Linnaeus, 1758)

Pearce: 1 locality (1821)
No subsequent records

Hydrobius fuscipes (Linnaeus, 1758)

Pearce: generally common
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Limnoxenus niger (Gmelin, 1790)

Pearce: 1 locality
First subsequent record: 1964
Number of subsequent records: 1
Latest record: 1964

Chaetarthria seminulum (Herbst, 1797)

Pearce: 1 locality (*sensu lato*)
First subsequent record: 2005
Number of subsequent records: many
Latest record: 2017

Chaetarthria simillima Vorst & Cuppen, 2003

Pearce: no records
First record: 1997
Number of records: many
Latest record: 2019

Anacaena bipustulata (Marshall, 1802)

Pearce: 3 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2019

Anacaena globulus (Paykull, 1798)

Pearce: 6 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Anacaena limbata (Fabricius, 1792)

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Anacaena lutescens (Stephens, 1829)

Pearce: generally common
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2019

Cymbiodyta marginella (Fabricius, 1792)

Pearce: 2 localities (1899, 1922)
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2020

Enochrus (Enochrus) melanocephalus (Olivier, 1793)

Pearce: 1 locality
First subsequent record: 1989
Number of subsequent records: 3
Latest record: 2020

Enochrus (Lumetus) bicolor (Fabricius, 1792)

Pearce: 2 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2014

Enochrus (Lumetus) fuscipennis (Thomson, C.G., 1884)

Pearce: no records
First record: 2002
Number of records: 4
Latest record: 2009

Enochrus (Lumetus) halophilus (Bedel, 1878)

Pearce: no records
First record: 2000
Number of records: 7
Latest record: 2006

Enochrus (Lumetus) ochropterus
(Marsham, 1802)
Pearce: 2 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2019

Enochrus (Lumetus) quadripunctatus
(Herbst, 1797)
Pearce: no records
First record: 1963
Number of records: 9
Latest record: 2007

Enochrus (Lumetus) testaceus (Fabricius, 1801)
Pearce: 1 locality
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2019

Enochrus (Methydrus) affinis (Thunberg, 1794)
Pearce: 2 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2019

Enochrus (Methydrus) coarctatus
(Gredler, 1863)
Pearce: 2 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2015

Enochrus (Methydrus) nigritus (Sharp, 1873)
Pearce: no records
First record: 2009
Number of records: 1
Latest record: 2009

Helochares lividus (Forster, 1771)
Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Helochares punctatus Sharp, 1869
Pearce: 10 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Coelostoma orbiculare (Fabricius, 1775)
Pearce: 4 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2021

Dactylosternum abdominale (Fabricius, 1792)
Pearce: no records
First record: 2003
Number of records: 4
Latest record: 2006

Sphaeridium bipustulatum Fabricius, 1781
Pearce: 7 localities
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2020

Sphaeridium lunatum Fabricius, 1792
Pearce: no records
First record: 1969
Number of records: many
Latest record: 2021

Sphaeridium marginatum Fabricius, 1787
Pearce: no records
First record: 1997
Number of records: 8
Latest record: 2018

Sphaeridium scarabaeoides (Linnaeus, 1758)
Pearce: generally common and widely distributed
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Cercyon (Cercyon) bifenestratus Küster, 1851
Pearce: no records
First record: 2019
Number of records: 2
Latest record: 2020

Cercyon (Cercyon) convexiusculus
Stephens, 1829
Pearce: 2 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2020

Cercyon (Cercyon) depressus Stephens, 1829

Pearce: 7 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2019

Cercyon (Cercyon) granarius Erichson, 1837

Pearce: 2 localities
First subsequent record: 1998
Number of subsequent records: 2
Latest record: 2013

Cercyon (Cercyon) haemorrhoidalis (Fabricius, 1775)

Pearce: 2 localities
First subsequent record: 1987
Number of subsequent records: many
Latest record: 2019

Cercyon (Cercyon) impressus (Sturm, 1807)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2018

Cercyon (Cercyon) lateralis (Marshall, 1802)

Pearce: 3 localities
First subsequent record: 1987
Number of subsequent records: many
Latest record: 2018

Cercyon (Cercyon) littoralis (Gyllenhal, 1808)

Pearce: 7 localities
First subsequent record: 1947
Number of subsequent records: many
Latest record: 2019

Cercyon (Cercyon) marinus Thomson, C.G., 1853

Pearce: 3 localities
First subsequent record: 2009
Number of subsequent records: 6
Latest record: 2017

Cercyon (Cercyon) melanocephalus (Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Cercyon (Cercyon) nigriceps (Marshall, 1802)

Pearce: 1 locality (1910)
First subsequent record: 1993
Number of subsequent records: 4
Latest record: 2006

Cercyon (Cercyon) obsoletus (Gyllenhal, 1808)

Pearce: 2 localities
First subsequent record: 1997
Number of subsequent records: 7
Latest record: 2021

Cercyon (Cercyon) pygmaeus (Illiger, 1801)

Pearce: 2 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Cercyon (Cercyon) quisquilius (Linnaeus, 1760)

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2018

Cercyon (Cercyon) sternalis Sharp, 1918

Pearce: no records
First record: 1998
Number of records: many
Latest record: 2014

Cercyon (Cercyon) terminatus (Marshall, 1802)

Pearce: 1 locality
First subsequent record: 1992
Number of subsequent records: 2
Latest record: 2006

Cercyon (Cercyon) tristis (Illiger, 1801)

Pearce: 1 locality
First subsequent record: 1987
Number of subsequent records: many
Latest record: 2019

Cercyon (Cercyon) unipunctatus (Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: 6
Latest record: 2018

Cercyon (Dicyrtocercyon) ustulatus
(Preyssier, 1790)

Pearce: 1 locality (1892)
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2021

Cercyon (Paracercyon) analis (Paykull, 1798)

Pearce: 2 localities
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2021

Cercyon (Paracycreon) laminatus Sharp, 1873

Pearce: no records
First record: 1997
Number of records: 5
Latest record: 2020

Cryptopleurum crenatum (Kugelann, 1794)

Pearce: no records
First record: 1994
Number of records: 5
Latest record: 2014

Cryptopleurum minutum (Fabricius, 1775)

Pearce: 7 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Cryptopleurum subtile Sharp, 1884

Pearce: no records
First record: 2003
Number of records: 1
Latest record: 2003

Megasternum concinnum (Marshall, 1802)

Pearce: 5 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

13. Family SPHAERITIDAE Shuckard, 1839

Sphaerites glabratus (Fabricius, 1792)

Pearce: no records
First record: 2000
Number of records: 1
Latest record: 2000

14. Family HISTERIDAE Gyllenhal, 1808

Abraeus (Abraeus) perpusillus (Marshall, 1802)

Pearce: 1 locality
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2021

Abraeus (Postabraeus) granulum

Erichson, 1839
Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Plegaderus dissectus Erichson, 1839

Pearce: no records
First record: 2000
Number of records: many
Latest record: 2020

Plegaderus vulneratus (Panzer, 1797)

Pearce: no records
First record: 1992
Number of records: 3
Latest record: 2016

Acritus (Acritus) nigricornis (Hoffmann, J., 1803)

Pearce: no records
First record: 1969
Number of records: many
Latest record: 2020

Acritus (Pyncacritus) homeopathicus

Wollaston, 1857
Pearce: no records
First record: 1937
Number of records: 2
Latest record: 1941

Aeletes atomarius (Aubé, 1842)

Pearce: no records
First record: 2004
Number of records: 2
Latest record: 2019

Halacritus punctum (Aubé, 1842)

Pearce: 2 localities
First subsequent record: 2016
Number of subsequent records: 1
Latest record: 2016

Saprinus aeneus (Fabricius, 1775)

Pearce: 2 localities
No subsequent records

Saprinus semistriatus (Scriba, 1790)

Pearce: 4 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2019

Saprinus virescens (Paykull, 1798)

Pearce: 3 localities
First subsequent record: 1943
Number of subsequent records: 6
Latest record: 2019

Hypocaccus (Baeckmanniolus) dimidiatus (Illiger, 1807)

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: 5
Latest record: 2019

Hypocaccus (Hypocaccus) rugifrons (Paykull, 1798)

Pearce: 1 locality
No subsequent records

Gnathoncus communis (Marseul, 1862)

Pearce: no records
First record: 1987
Number of records: 1
Latest record: 1987

Gnathoncus nannetensis (Marseul, 1862)

Pearce: 3 localities
First subsequent record: 1987
Number of subsequent records: 4
Latest record: 2020

Gnathoncus rotundatus (Kugelann, 1792)

Pearce: no records
First record: 1954
Number of records: 1
Latest record: 1954

Myrmeces paykulli Kanaar, 1979

Pearce: 2 localities
No subsequent records

Dendrophilus punctatus (Herbst, 1791)

Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Kissister minimus (Laporte, 1840)

Pearce: 4 localities
First subsequent record: 1952
Number of subsequent records: many
Latest record: 2021

Carcinops pumilio (Erichson, 1834)

Pearce: no records
First record: 2000
Number of records: 4
Latest record: 2008

Paromalus flavicornis (Herbst, 1791)

Pearce: no records
First record: 1953
Number of records: many
Latest record: 2021

Onthophilus striatus (Forster, 1771)

Pearce: 5 localities
First subsequent record: 1971
Number of subsequent records: 4
Latest record: 2007

Margarinotus (Paralister) neglectus (Germar, 1813)

Pearce: no records
First record: 1993
Number of records: 4
Latest record: 2018

Margarinotus (Paralister) purpurascens (Herbst, 1791)

Pearce: 1 locality
First subsequent record: 2001
Number of subsequent records: 7
Latest record: 2019

Margarinotus (Paralister) ventralis (Marseul, 1854)

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Margarinotus (Promethister) marginatus (Erichson, 1834)

Pearce: 2 localities
First subsequent record: 2002
Number of subsequent records: 1
Latest record: 2002

Margarinotus (Ptomister) brunneus
(Fabricius, 1775)

Pearce: generally common and widely distributed
First subsequent record: 1943
Number of subsequent records: 6
Latest record: 2014

Margarinotus (Ptomister) merdarius
(Hoffmann, J., 1803)

Pearce: no records
First record: 1993
Number of records: 6
Latest record: 2020

Margarinotus (Ptomister) striola
(Sahlberg, C.R., 1819)

Pearce: 2 localities
First subsequent record: 1991
Number of subsequent records: 4
Latest record: 2019

Hister quadrimaculatus Linnaeus, 1758

Pearce: 1 locality (1901)
No subsequent records

Hister unicolor Linnaeus, 1758

Pearce: 3 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2018

Atholus bimaculatus (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1971
Number of subsequent records: 9
Latest record: 2013

Atholus duodecimstriatus (Schränk, 1781)

Pearce: 4 localities
First subsequent record: 2003
Number of subsequent records: 7
Latest record: 2021

Haeterius ferrugineus (Olivier, 1789)

Pearce: no records
First record: 2008
Number of records: 1
Latest record: 2008

15. Family HYDRAENIDAE Mulsant, 1844

Hydraena britteni Joy, 1907

Pearce: no records
First record: 1998
Number of records: 2
Latest record: 2017

Hydraena gracilis Germar, 1824

Pearce: no records
First record: 1987
Number of records: 8
Latest record: 2011

Hydraena nigrita Germar, 1824

Pearce: 1 locality
First subsequent record: 1981
Number of subsequent records: 5
Latest record: 2019

Hydraena pulchella Germar, 1824

Pearce: no records
First record: 1830
Number of records: 2
Latest record: 1861

Hydraena riparia Kugelann, 1794

Pearce: 5 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Hydraena rufipes Curtis, 1830

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 2
Latest record: 1998

Hydraena testacea Curtis, 1830

Pearce: 1 locality (1815)
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2018

Limnebius nitidus (Marsham, 1802)

Pearce: 2 localities
First subsequent record: 1990
Number of subsequent records: many
Latest record: 2017

Limnebius papposus Mulsant, 1844

Pearce: 5 localities
First subsequent record: 1938
Number of subsequent records: 3
Latest record: 2006

Limnebius truncatellus (Thunberg, 1794)

Pearce: 8 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2019

Ochthebius (Asiobates) auriculatus Rey, 1886

Pearce: no records
First record: 2005
Number of records: 3
Latest record: 2011

Ochthebius (Asiobates) bicolon Germar, 1824

Pearce: 2 localities
First subsequent record: 2002
Number of subsequent records: 5
Latest record: 2011

Ochthebius (Asiobates) dilatatus Stephens, 1829

Pearce: 4 localities
First subsequent record: 1962
Number of subsequent records: many
Latest record: 2021

Ochthebius (Asiobates) minimus (Fabricius, 1792)

Pearce: 2 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Ochthebius (Enicocerus) exsculptus Germar, 1824

Pearce: 1 locality
No subsequent records

Ochthebius (Ochthebius) lejolisii Mulsant & Rey, 1861

Pearce: no records
First record: 2008
Number of records: 6
Latest record: 2021

Ochthebius (Ochthebius) marinus (Paykull, 1798)

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2020

Ochthebius (Ochthebius) nanus Stephens, 1829

Pearce: 2 localities
First subsequent record: 2005
Number of subsequent records: 3
Latest record: 2011

Ochthebius (Ochthebius) poweri Rye, 1869

Pearce: 1 locality
First subsequent record: 1971
Number of subsequent records: many
Latest record: 2019

Ochthebius (Ochthebius) punctatus Stephens, 1829

Pearce: no records
First record: 1935
Number of records: many
Latest record: 2016

Ochthebius (Ochthebius) pusillus Stephens, 1835

Pearce: 1 locality
First subsequent record: 2005
Number of subsequent records: 6
Latest record: 2006

Ochthebius (Ochthebius) viridis Peyron, 1858

Pearce: 2 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2014

16. Family PTILIIDAE Erichson, 1845

Ptenidium (Gillmeisterium) nitidum (Heer, 1841)

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: 4
Latest record: 2019

Ptenidium (Gressnerium) gressneri Erichson, 1845

Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Ptenidium (Matthewsium) laevigatum

Erichson, 1845

Pearce: no records

First record: 1995

Number of records: 4

Latest record: 2019

Ptenidium (Ptenidium) fuscicorne

Erichson, 1845

Pearce: no records

First record: 1990

Number of records: 2

Latest record: 1995

Ptenidium (Ptenidium) longicorne Fuss,
1868

Pearce: no records

First record: 2004

Number of records: 4

Latest record: 2017

Ptenidium (Ptenidium) myrmicophilum
(Motschulsky, 1845) [2012: *Ptenidium*
formicetorum]

Pearce: 2 localities

First subsequent record: 2003

Number of subsequent records: 2

Latest record: 2006

Ptenidium (Ptenidium) punctatum
(Gyllenhal, 1827)

Pearce: 5 localities

First subsequent record: 2004

Number of subsequent records: 7

Latest record: 2019

Ptenidium (Ptenidium) pusillum
(Gyllenhal, 1808)

Pearce: 4 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2020

Ptenidium (Wankowizium) intermedium
Wankowicz, 1869

Pearce: no records

First record: 1995

Number of records: 3

Latest record: 2008

Ptiliolium spencei (Allibert, 1844)

Pearce: 1 locality

No subsequent records

Oligella foveolata (Allibert, 1844)

Pearce: no records

First record: 1997

Number of records: 1

Latest record: 1997

Ptilium exaratum (Allibert, 1844)

Pearce: no records

First record: 2000

Number of records: 1

Latest record: 2000

Ptilium horioni Rosskothén, 1934

Pearce: no records

First record: 1997

Number of records: 1

Latest record: 1997

Nephanes titan (Newman, 1834)

Pearce: 1 locality

First subsequent record: 2003

Number of subsequent records: 3

Latest record: 2003

Actinopteryx fucicola (Allibert, 1844)

Pearce: 1 locality

No subsequent records

Acrotrichis (Acrotrichis) atomaria (De
Geer, 1774)

Pearce: 4 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2014

Acrotrichis (Acrotrichis) brevipennis
(Erichson, 1845)

Pearce: 1 locality

No subsequent records

Acrotrichis (Acrotrichis) cognata
(Matthews, A., 1877)

Pearce: no records

First record: 1975

Number of records: 1

Latest record: 1975

Acrotrichis (Acrotrichis) danica Sundt,
1958

Pearce: no records

First record: 2010

Number of records: 5

Latest record: 2011

Acrotrichis (Acrotrichis) fascicularis
(Herbst, 1793)
Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2013

Acrotrichis (Acrotrichis) henrici
(Matthews, A., 1872)
Pearce: no records
First record: 1994
Number of records: 8
Latest record: 2004

Acrotrichis (Acrotrichis) insularis
(Mäklin, 1852)
Pearce: no records
First record: 2013
Number of records: 1
Latest record: 2013

Acrotrichis (Acrotrichis) intermedia
(Gillmeister, 1845)
Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: 7
Latest record: 2020

Acrotrichis (Acrotrichis) montandonii
(Allibert, 1844)
Pearce: no records
First record: 1992
Number of records: 1
Latest record: 1992

Acrotrichis (Acrotrichis) rosskotheni
Sundt, 1971
Pearce: no records
First record: 1979
Number of records: 3
Latest record: 1994

Acrotrichis (Acrotrichis) rugulosa
Rosskothén, 1935
Pearce: no records
First record: 1975
Number of records: 2
Latest record: 2015

Acrotrichis (Acrotrichis) sericans (Heer,
1841)
Pearce: 3 localities
No subsequent records

Acrotrichis (Acrotrichis) sitkaensis
(Motschulsky, 1845)
Pearce: no records
First record: 2017
Number of records: 2
Latest record: 2021

Acrotrichis (Acrotrichis) thoracica (Waltl,
1838)
Pearce: no records
First record: 2003
Number of records: 1
Latest record: 2003

Acrotrichis (Ctenopteryx) grandicollis
(Mannerheim, 1844)
Pearce: no records
First record: 1975
Number of records: 4
Latest record: 2006

Acrotrichis (Ctenopteryx) sanctaehelenae
Johnson, 1972
Pearce: no records
First record: 2003
Number of records: 3
Latest record: 2003

17. Family LEIODIDAE Fleming, 1821

Hydnobius punctatus (Sturm, 1807)
Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: 1
Latest record: 1975

Triarthron maerkelii Schmidt, W.L.E.,
1840
Pearce: 1 locality (1925)
No subsequent records

Leiodes badia (Sturm, 1807)
Pearce: 1 locality
First subsequent record: 1998
Number of subsequent records: 2
Latest record: 2008

Leiodes calcarata (Erichson, 1845)
Pearce: 3 localities
First subsequent record: 1993
Number of subsequent records: 5
Latest record: 2016

Leiodes ferruginea (Fabricius, 1787)

Pearce: 1 locality (1892)
First subsequent record: 2018
Number of subsequent records: 1
Latest record: 2018

Leiodes gyllenhalii Stephens, 1829

Pearce: 1 locality
No subsequent records

Leiodes litura Stephens, 1835

Pearce: 2 localities
First subsequent record: 1976
Number of subsequent records: 2
Latest record: 2019

Leiodes lunicollis (Rye, 1872)

Pearce: 1 locality
No subsequent records

Leiodes nigrita (Schmidt, W.L.E., 1841)

Pearce: 3 localities
No subsequent records

Leiodes rufipennis (Paykull, 1798)

Pearce: 3 localities
First subsequent record: 2008
Number of subsequent records: 6
Latest record: 2019

Liocyrtusa vittata (Curtis, 1840)

Pearce: 2 localities
No subsequent records

Colenis immunda (Sturm, 1807)

Pearce: 1 locality
First subsequent record: 1987
Number of subsequent records: 1
Latest record: 1987

Anisotoma humeralis (Herbst, 1791)

Pearce: 4 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Anisotoma orbicularis (Herbst, 1791)

Pearce: 1 locality (1892)
First subsequent record: 2000
Number of subsequent records: 2
Latest record: 2004

Amphicyllis globus (Fabricius, 1792)

Pearce: 2 localities
First subsequent record: 1977
Number of subsequent records: many
Latest record: 2015

Agathidium (Agathidium) atrum (Paykull, 1798)

Pearce: 3 localities
First subsequent record: 1953
Number of subsequent records: 1
Latest record: 1953

Agathidium (Agathidium) laevigatum
Erichson, 1845

Pearce: 1 locality
First subsequent record: 1987
Number of subsequent records: many
Latest record: 2019

Agathidium (Agathidium) seminulum
(Linnaeus, 1758)

Pearce: 1 locality
First subsequent record: 2013
Number of subsequent records: 3
Latest record: 2019

Agathidium (Cyphocheble) nigrinum Sturm, 1807

Pearce: no records
First record: 2015
Number of records: 3
Latest record: 2017

Agathidium (Neocheble) confusum Brisout de Barneville, C., 1863

Pearce: no records
First record: 2013
Number of records: 2
Latest record: 2013

Agathidium (Neocheble) marginatum
Sturm, 1807

Pearce: no records
First record: 1979
Number of records: 3
Latest record: 2017

Agathidium (Neocheble) nigripenne
(Fabricius, 1792)

Pearce: 1 locality
First subsequent record: 2007
Number of subsequent records: 2
Latest record: 2021

Agathidium (Neoceble) rotundatum
(Gyllenhal, 1827)

Pearce: no records
First record: 2008
Number of records: 4
Latest record: 2008

Agathidium (Neoceble) varians Beck,
1817

Pearce: 2 localities
First subsequent record: 2014
Number of subsequent records: 1
Latest record: 2014

Colon (Eurycolon) latum Kraatz, 1850

Pearce: 2 localities (1927, 1929)
No subsequent records

Colon (Myloechus) brunneum (Latreille,
1806)

Pearce: 2 localities
First subsequent record: 1993
Number of subsequent records: 3
Latest record: 1993

Nemadus colonoides (Kraatz, 1851)

Pearce: no records
First record: 2014
Number of records: 4
Latest record: 2020

Nargus (Demochrus) anisotomoides
(Spence, 1813)

Pearce: 3 localities
First subsequent record: 2015
Number of subsequent records: many
Latest record: 2015

Nargus (Demochrus) wilkini (Spence,
1813)

Pearce: 4 localities
First subsequent record: 1969
Number of subsequent records: 3
Latest record: 2016

Nargus (Nargus) velox (Spence, 1813)

Pearce: 2 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2021

Choleva (Choleva) agilis (Illiger, 1798)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 8
Latest record: 2015

Choleva (Choleva) angustata (Fabricius,
1781)

Pearce: 3 localities
First subsequent record: 1987
Number of subsequent records: 6
Latest record: 2015

Choleva (Choleva) cisteloides (Frölich,
1799)

Pearce: 2 localities
No subsequent records

Choleva (Choleva) fagniezi Jeannel, 1922

Pearce: no records
First record: 2017
Number of records: 2
Latest record: 2019

Choleva (Choleva) glauca Britten, 1918

Pearce: no records
First record: 2008
Number of records: 2
Latest record: 2008

Choleva (Choleva) jeanneli Britten, 1922

Pearce: no records
First record: 1979
Number of records: 8
Latest record: 2007

Choleva (Choleva) oblonga Latreille, 1806

Pearce: no records
First record: 1976
Number of records: 2
Latest record: 2016

Choleva (Cholevopsis) spadicea (Sturm,
1839)

Pearce: no records
First record: 1994
Number of records: 4
Latest record: 1999

Sciodrepoides fumatus (Spence, 1813)

Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2021

Sciodrepoides watsoni (Spence, 1813)

Pearce: 8 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Catops chrysomeloides (Panzer, 1798)

Pearce: 4 localities
No subsequent records

Catops coracinus Kellner, 1846

Pearce: no records
First record: 1952
Number of records: many
Latest record: 2016

Catops fuliginosus Erichson, 1837

Pearce: no records
First record: 1975
Number of records: many
Latest record: 2018

Catops fuscus (Panzer, 1794)

Pearce: 2 localities
First subsequent record: 1997
Number of subsequent records: 6
Latest record: 2019

Catops grandicollis Erichson, 1837

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: 4
Latest record: 2016

Catops kirbyi (Spence, 1813)

Pearce: 2 localities
First subsequent record: 1999
Number of subsequent records: 5
Latest record: 2019

Catops morio (Fabricius, 1787)

Pearce: 4 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2018

Catops nigricans (Spence, 1813)

Pearce: 4 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Catops tristis (Panzer, 1794)

Pearce: 7 localities
First subsequent record: 1986
Number of subsequent records: 8
Latest record: 2021

Apocatops nigrita (Erichson, 1837) [2012:
Catops nigrita]

Pearce: 2 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2018

Parabathyscia wollastoni (Janson, 1857)

Pearce: no records
First record: 2019
Number of records: 9
Latest record: 2021

Ptomaphagus sericatus (Chaudoir, 1845)
[2012: *Ptomaphagus medius*]

Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: 5
Latest record: 2021

Ptomaphagus subvillosus (Goeze, 1777)

Pearce: 10 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Leptinus testaceus Müller, P.W.J., 1817

Pearce: 1 locality (1893)
First subsequent record: 2000
Number of subsequent records: 1
Latest record: 2000

18. Family SILPHIDAE Latreille, 1806

Necrodes littoralis (Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1940
Number of subsequent records: many
Latest record: 2018

Thanatophilus rugosus (Linnaeus, 1758)

Pearce: generally common and widely distributed
First subsequent record: 1997
Number of subsequent records: 4
Latest record: 2019

Thanatophilus sinuatus (Fabricius, 1775)

Pearce: 5 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2003

Oiceoptoma thoracicum (Linnaeus, 1758)

Pearce: 9 localities
First subsequent record: 1992
Number of subsequent records: 9
Latest record: 2011

Aclypea opaca (Linnaeus, 1758)

Pearce: 9 localities
First subsequent record: 1975
Number of subsequent records: 2
Latest record: 1980

Dendroxena quadrimaculata (Scopoli, 1771)

Pearce: 3 localities
First subsequent record: 1934
Number of subsequent records: 2
Latest record: 1941

Phosphuga atrata (Linnaeus, 1758) [2012: *Silpha atrata*]

Pearce: generally common and widely distributed
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Ablattaria laevigata (Fabricius, 1775) [2012: *Silpha laevigata*]

Pearce: generally common
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2020

Silpha obscura Linnaeus, 1758

Pearce: 2 localities
First subsequent record: 2012
Number of subsequent records: 3
Latest record: 2016

Silpha tristis Illiger, 1798

Pearce: 5 localities
First subsequent record: 1952
Number of subsequent records: many
Latest record: 2021

Nicrophorus humator (Gleditsch, 1767)

Pearce: generally common
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Nicrophorus interruptus Stephens, 1830

Pearce: 1 locality
First subsequent record: 1952
Number of subsequent records: many
Latest record: 2021

Nicrophorus investigator Zetterstedt, 1824

Pearce: 8 localities
First subsequent record: 1936
Number of subsequent records: 8
Latest record: 2021

Nicrophorus vespillo (Linnaeus, 1758)

Pearce: generally common and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Nicrophorus vespilloides Herbst, 1783

Pearce: generally common
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2020

Nicrophorus vestigator Herschel, 1807

Pearce: 8 localities
No subsequent records

19. Family STAPHYLINIDAE Latreille, 1802

Phyllodrepa floralis (Paykull, 1789)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: 8
Latest record: 2017

Dropephylla ioptera (Stephens, 1832)

Pearce: 1 locality
First subsequent record: 2000
Number of subsequent records: many
Latest record: 2021

Dropephylla koltzei Jászay & Hlavac, 2006

Pearce: no records
First record: 2018
Number of records: 3
Latest record: 2019

Dropephylla vilis (Erichson, 1840)

Pearce: 2 localities
First subsequent record: 1970
Number of subsequent records: 6
Latest record: 2019

Hapalareae pygmaea (Paykull, 1800)

Pearce: 2 localities
First subsequent record: 2020
Number of subsequent records: 1
Latest record: 2020

Acrolocha sulcula (Stephens, 1834)

Pearce: no records
First record: 2015
Number of records: 1
Latest record: 2015

Omalium allardii Fairmaire & Brisout de Barneville, C., 1859

Pearce: 2 localities
No subsequent records

Omalium caesum Gravenhorst, 1806

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Omalium excavatum Stephens, 1834

Pearce: 2 localities
First subsequent record: 2001
Number of subsequent records: 2
Latest record: 2020

Omalium italicum Bernhauer, 1902

Pearce: no records
First record: 2003
Number of records: 9
Latest record: 2020

Omalium laeviusculum Gyllenhal, 1827

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Omalium oxyacanthae Gravenhorst, 1806

Pearce: no records
First record: 1977
Number of records: 4
Latest record: 2010

Omalium riparium Thomson, C.G., 1857

Pearce: 8 localities
First subsequent record: 2004
Number of subsequent records: many
Latest record: 2021

Omalium rivulare (Paykull, 1789)

Pearce: generally common
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2021

Omalium rugatum Mulsant & Rey, 1880

Pearce: no records
First record: 1969
Number of records: 7
Latest record: 2021

Phloeonomus punctipennis Thomson, C.G., 1867

Pearce: 1 locality
First subsequent record: 2006
Number of subsequent records: many
Latest record: 2019

Phloeonomus pusillus (Gravenhorst, 1806)

Pearce: 1 locality
First subsequent record: 1991
Number of subsequent records: 4
Latest record: 2007

Phloeostiba plana (Paykull, 1792)

Pearce: 1 locality
First subsequent record: 2001
Number of subsequent records: 2
Latest record: 2019

Xylodromus concinnus (Marshall, 1802)

Pearce: 3 localities
First subsequent record: 1969
Number of subsequent records: 6
Latest record: 2006

Xylodromus depressus (Gravenhorst, 1802)

Pearce: no records
First record: 2006
Number of records: 1
Latest record: 2006

Eusphalerum luteum (Marshall, 1802)

Pearce: 4 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2019

Eusphalerum minutum (Fabricius, 1792)

Pearce: 3 localities
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2011

Eusphalerum primulae (Stephens, 1834)

Pearce: 1 locality
First subsequent record: 1963
Number of subsequent records: 5
Latest record: 2018

Eusphalerum sorbi (Gyllenhal, 1810)

Pearce: 3 localities

First subsequent record: 1996

Number of subsequent records: 3

Latest record: 2019

Eusphalerum torquatum (Marshall, 1802)

Pearce: 3 localities

No subsequent records

Anthobium atrocephalum (Gyllenhal, 1827)

Pearce: 2 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2019

Anthobium unicolor (Marshall, 1802)

Pearce: 4 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Olophrum piceum (Gyllenhal, 1810)

Pearce: 5 localities

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2021

Acidota crenata (Fabricius, 1792)

Pearce: 1 locality

First subsequent record: 1979

Number of subsequent records: 6

Latest record: 2016

Acidota cruentata Mannerheim, 1830

Pearce: 1 locality

First subsequent record: 1975

Number of subsequent records: 2

Latest record: 1999

Lesteva (Lesteva) hansenii Lohse, 1953

Pearce: 1 locality

No subsequent records

Lesteva (Lesteva) longoelytrata (Goeze, 1777)

Pearce: 3 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2020

Lesteva (Lesteva) pubescens Mannerheim, 1830

Pearce: 1 locality

First subsequent record: 1963

Number of subsequent records: 5

Latest record: 2017

Lesteva (Lestevidia) punctata Erichson, 1839

Pearce: 2 localities

First subsequent record: 2006

Number of subsequent records: 4

Latest record: 2019

Lesteva (Lestevina) sicula Erichson, 1840

Pearce: 3 localities

First subsequent record: 1968

Number of subsequent records: many

Latest record: 2021

Micralymma marinum (Strøm, 1783)

Pearce: 2 localities (1883, 1894)

First subsequent record: 1953

Number of subsequent records: 1

Latest record: 1953

Philorinum sordidum (Stephens, 1834)

Pearce: 3 localities

First subsequent record: 1982

Number of subsequent records: many

Latest record: 2016

Coryphium angusticolle Stephens, 1834

Pearce: 2 localities

First subsequent record: 1934

Number of subsequent records: 4

Latest record: 2011

Metopsia clypeata (Müller, P.W.J., 1821)

Pearce: 3 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Megarthus bellevoeyi Saulcy, 1862

Pearce: 3 localities

First subsequent record: 1969

Number of subsequent records: 6

Latest record: 2014

Megarthus denticollis (Beck, 1817)

Pearce: no records

First record: 1972

Number of records: 4

Latest record: 2017

Megarthus depressus (Paykull, 1789)

Pearce: 1 locality
First subsequent record: 1972
Number of subsequent records: 9
Latest record: 2021

Megarthus hemipterus (Illiger, 1794)

Pearce: 1 locality
No subsequent records

Megarthus prosseni Schatzmayer, 1904

Pearce: 3 localities
First subsequent record: 1970
Number of subsequent records: 9
Latest record: 2018

Proteinus brachypterus (Fabricius, 1792)

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Proteinus laevigatus Hochhuth, 1872

Pearce: 2 localities
No subsequent records

Proteinus ovalis Stephens, 1834

Pearce: 5 localities
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2021

Micropeplus fulvus Erichson, 1840

Pearce: 1 locality
First subsequent record: 1969
Number of subsequent records: 8
Latest record: 2008

Micropeplus porcatus (Paykull, 1789)

Pearce: 1 locality
No subsequent records

Micropeplus staphylinoides (Marshall, 1802)

Pearce: 2 localities
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2021

Micropeplus tesserula Curtis 1828

Pearce: "Dorset" (A. Ford)
First subsequent record: 1986
Number of subsequent records: 1
Latest record: 1986

Bibloporus bicolor (Denny, 1825)

Pearce: no records
First record: 2018
Number of records: many
Latest record: 2019

Biblopectus ambiguus (Reichenbach, 1816)

Pearce: 1 locality (1925)
First subsequent record: 2014
Number of subsequent records: many
Latest record: 2021

Trichonyx sulcicollis (Reichenbach, 1816)

Pearce: 1 locality (1921)
No subsequent records

Amauronyx maerkelii (Aubé, 1844)

Pearce: 3 localities
No subsequent records

Euplectus karstenii (Reichenbach, 1816)

Pearce: 1 locality (1838)
First subsequent record: 2003
Number of subsequent records: many
Latest record: 2020

Euplectus kirbii Denny, 1825

Pearce: no records
First record: 2006
Number of records: 4
Latest record: 2019

Euplectus mutator Fauvel, 1895

Pearce: no records
First record: 2020
Number of records: 1
Latest record: 2020

Euplectus nanus (Reichenbach, 1816)

Pearce: 1 locality
No subsequent records

Euplectus piceus Motschulsky, 1835

Pearce: 1 locality (1890)
First subsequent record: 2018
Number of subsequent records: 8
Latest record: 2019

Euplectus sanguineus Denny, 1825

Pearce: 1 locality
No subsequent records

Euplectus signatus (Reichenbach, 1816)

Pearce: 1 locality
No subsequent records

Rybaxis longicornis (Leach, 1817)

Pearce: 5 localities
First subsequent record: 1989
Number of subsequent records: many
Latest record: 2020

Brachygluta fossulata (Reichenbach, 1816)

Pearce: 4 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2020

Brachygluta haematica (Reichenbach, 1816)

Pearce: 2 localities
First subsequent record: 1987
Number of subsequent records: 3
Latest record: 2019

Brachygluta helferi (Schmidt-Göbel, 1836)

Pearce: 3 localities
First subsequent record: 1995
Number of subsequent records: many
Latest record: 2021

Brachygluta waterhousei (Rye, 1869)

Pearce: 3 localities
First subsequent record: 1966
Number of subsequent records: 4
Latest record: 2020

Reichenbachia juncorum (Leach, 1817)

Pearce: 6 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2020

Fagniezia impressa (Panzer, 1803)

Pearce: 3 localities
First subsequent record: 1965
Number of subsequent records: 2
Latest record: 2015

Tychobythinus glabratus (Rye, 1870)

Pearce: 1 locality
No subsequent records

Bythinus burrellii Denny, 1825

Pearce: no records
First record: 2013
Number of records: 2
Latest record: 2015

Bythinus macropalpus Aubé, 1833

Pearce: 1 locality
First subsequent record: 1987
Number of subsequent records: 4
Latest record: 2016

Bryaxis bulbifer (Reichenbach, 1816)

Pearce: 3 localities
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2021

Bryaxis curtisii (Leach, 1817)

Pearce: 3 localities
First subsequent record: 1986
Number of subsequent records: 8
Latest record: 2021

Bryaxis puncticollis (Denny, 1825)

Pearce: 2 localities
No subsequent records

Tychus niger (Paykull, 1800)

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: 9
Latest record: 2019

Pselaphaulax dresdensis Reitter, 1909

Pearce: 1 locality (1925)
No subsequent records

Pselaphus heisei Herbst, 1791

Pearce: 4 localities
First subsequent record: 1979
Number of subsequent records: 8
Latest record: 2017

Claviger testaceus Preysslér, 1790

Pearce: 5 localities
First subsequent record: 1936
Number of subsequent records: 2
Latest record: 1991

Phloeocharis subtilissima Mannerheim, 1830

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Sepedophilus bipunctatus (Gravenhorst, 1802)

Pearce: 1 locality
First subsequent record: 2019
Number of subsequent records: 1
Latest record: 2019

Sepedophilus constans (Fowler, 1888)

Pearce: no records
First record: 2008
Number of records: 6
Latest record: 2021

Sepedophilus immaculatus (Stephens, 1832)

Pearce: 1 locality (1893)
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2018

Sepedophilus littoreus (Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2018

Sepedophilus lusitanicus Hammond, 1973

Pearce: no records
First record: 2002
Number of records: 7
Latest record: 2021

Sepedophilus marshami (Stephens, 1832)

Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Sepedophilus nigripennis (Stephens, 1832)

Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Sepedophilus pedicularius (Gravenhorst, 1802)

Pearce: 2 localities
First subsequent record: 1989
Number of subsequent records: 5
Latest record: 2007

Tachyporus (Palporus) nitidulus (Fabricius, 1781)

Pearce: 5 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Tachyporus (Tachyporus) atriceps Stephens, 1832

Pearce: 7 localities
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2021

Tachyporus (Tachyporus) chrysomelinus (Linnaeus, 1758)

Pearce: 5 localities (but including *dispar*)
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2021

Tachyporus (Tachyporus) dispar (Paykull, 1789)

Pearce: no records (but see *chrysomelinus*)
First record: 1996
Number of records: many
Latest record: 2021

Tachyporus (Tachyporus) formosus Matthews, A.H., 1838

Pearce: 2 localities
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2021

Tachyporus (Tachyporus) hypnorum (Fabricius, 1775)

Pearce: generally common and widely distributed
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Tachyporus (Tachyporus) obtusus (Linnaeus, 1767)

Pearce: 7 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2020

Tachyporus (Tachyporus) pallidus Sharp, 1871

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Tachyporus (Tachyporus) pusillus

Gravenhorst, 1806

Pearce: 4 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2019

Tachyporus (Tachyporus) solutus

Erichson, 1839

Pearce: generally common

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2021

Tachyporus (Tachyporus) tersus Erichson, 1839

Pearce: 2 localities

First subsequent record: 1979

Number of subsequent records: 4

Latest record: 2005

Tachyporus (Tachyporus) transversalis

Gravenhorst, 1806

Pearce: 1 locality (1925)

First subsequent record: 2014

Number of subsequent records: 5

Latest record: 2019

Lamprinodes saginatus (Gravenhorst, 1806)

Pearce: no records

First record: 2019

Number of records: 1

Latest record: 2019

Tachinus bipustulatus (Fabricius, 1792)

Pearce: 1 locality

No subsequent records

Tachinus corticinus Gravenhorst, 1802

Pearce: no records

First record: 1986

Number of records: 4

Latest record: 2011

Tachinus elongatus Gyllenhal, 1810

Pearce: 1 locality

No subsequent records

Tachinus humeralis Gravenhorst, 1802

Pearce: 2 localities

First subsequent record: 1992

Number of subsequent records: 5

Latest record: 2019

Tachinus laticollis Gravenhorst, 1802

Pearce: no records

First record: 1968

Number of records: many

Latest record: 2021

Tachinus lignorum (Linnaeus, 1758)

Pearce: 1 locality

No subsequent records

Tachinus marginellus (Fabricius, 1781)

Pearce: 3 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2018

Tachinus pallipes (Gravenhorst, 1806)

Pearce: no records

First record: 2000

Number of records: 2

Latest record: 2015

Tachinus proximus Kraatz, 1855

Pearce: no records

First record: 1994

Number of records: 2

Latest record: 1994

Tachinus rufipennis Gyllenhal, 1810

Pearce: no records

First record: 1909

Number of records: 1

Latest record: 1909

Tachinus rufipes (Linnaeus, 1758)

Pearce: generally common and widely distributed

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2021

Tachinus subterraneus (Linnaeus, 1758)

Pearce: 3 localities

First subsequent record: 1987

Number of subsequent records: 9

Latest record: 2021

Cilea silphoides (Linnaeus, 1767)

Pearce: 3 localities

First subsequent record: 1964

Number of subsequent records: 4

Latest record: 2000

Mycetoporus angularis Mulsant & Rey, 1853

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Mycetoporus clavicornis (Stephens, 1832)

Pearce: 2 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2014

Mycetoporus despectus Strand, A., 1830

Pearce: no records
First record: 2005
Number of records: 1
Latest record: 2005

Mycetoporus lepidus (Gravenhorst, 1806)

Pearce: 4 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2020

Mycetoporus longulus Mannerheim, 1830

Pearce: 2 localities
First subsequent record: 2000
Number of subsequent records: 3
Latest record: 2018

Mycetoporus nigricollis Stephens, 1835

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2007

Mycetoporus piceolus Rey, 1883

Pearce: no records
First record: 1976
Number of records: 1
Latest record: 1976

Mycetoporus punctus (Gravenhorst, 1806)

Pearce: 2 localities
First subsequent record: 2013
Number of subsequent records: 2
Latest record: 2015

Mycetoporus rufescens (Stephens, 1832)

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Ischnosoma longicorne (Mäklin, 1847)

Pearce: 1 locality (1919)
First subsequent record: 1986
Number of subsequent records: 1
Latest record: 1986

Ischnosoma splendidum (Gravenhorst, 1806)

Pearce: 3 localities
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2021

Bryoporus cernuus (Gravenhorst, 1806)

Pearce: no records
First record: 1976
Number of records: 1
Latest record: 1976

Lordithon exoletus (Erichson, 1839)

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: 9
Latest record: 2015

Lordithon lunulatus (Linnaeus, 1760)

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Lordithon thoracicus (Fabricius, 1777)

Pearce: 6 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2020

Lordithon trinotatus (Erichson, 1839)

Pearce: 3 localities
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2020

Bolitobius castaneus (Stephens, 1832)

Pearce: 8 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2009

Bolitobius cingulatus Mannerheim, 1830

Pearce: 5 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2019

Parabolitobius inclinans (Gravenhorst, 1806)

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: 2
Latest record: 2004

Trichophya pilicornis (Gyllenhal, 1810)

Pearce: no records
First record: 1979
Number of records: 2
Latest record: 2019

Habrocerus capillaricornis (Gravenhorst, 1806)

Pearce: no records
First record: 1978
Number of records: many
Latest record: 2021

Gymnusa brevicollis (Paykull, 1800)

Pearce: 4 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2014

Gymnusa variegata Kiesenwetter, 1845

Pearce: 1 locality
No subsequent records

Deinopsis erosa (Stephens, 1832)

Pearce: 1 locality
First subsequent record: 1953
Number of subsequent records: 9
Latest record: 2017

Aleochara (Aleochara) curtula (Goeze, 1777)

Pearce: generally common
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2019

Aleochara (Aleochara) lata Gravenhorst, 1802

Pearce: 10 localities
First subsequent record: 1933
Number of subsequent records: 5
Latest record: 2010

Aleochara (Baryodma) intricata Mannerheim, 1830

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Aleochara (Ceranota) ruficornis

Gravenhorst, 1802
Pearce: 1 locality (1890)
No subsequent records

Aleochara (Coprochara) bilineata

Gyllenhal, 1810
Pearce: 1 locality
First subsequent record: 2004
Number of subsequent records: 4
Latest record: 2019

Aleochara (Coprochara) bipustulata (Linnaeus, 1760)

Pearce: 8 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Aleochara (Emplenota) obscurella

Gravenhorst, 1806
Pearce: 10 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2019

Aleochara (Polystomota) grisea Kraatz, 1856

Pearce: 5 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2019

Aleochara (Polystomota) punctatella

Motschulsky, 1858
Pearce: 3 localities
First subsequent record: 2010
Number of subsequent records: 4
Latest record: 2019

Aleochara (Xenochara) brevipennis

Gravenhorst, 1806
Pearce: 3 localities
First subsequent record: 2004
Number of subsequent records: many
Latest record: 2017

Aleochara (Xenochara) cuniculorum Kraatz, 1858

Pearce: 1 locality
No subsequent records

Aleochara (Xenochara) diversa (Sahlberg, J., 1876) [2012: *Aleochara kamila*]
Pearce: no records
First record: 1975
Number of records: 1
Latest record: 1975

Aleochara (Xenochara) fumata
Gravenhorst, 1802
Pearce: 1 locality (1894)
No subsequent records

Aleochara (Xenochara) funebris
Wollaston, 1864
Pearce: no records
First record: 1975
Number of records: 3
Latest record: 1975

Aleochara (Xenochara) lanuginosa
Gravenhorst, 1802
Pearce: 8 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2019

Aleochara (Xenochara) moesta
Gravenhorst, 1802
Pearce: 2 localities
No subsequent records

Aleochara (Xenochara) sanguinea
(Linnaeus, 1758)
Pearce: no records
First record: 2004
Number of records: 1
Latest record: 2004

Aleochara (Xenochara) spadicea
(Erichson, 1837)
Pearce: 1 locality (1928)
First subsequent record: 1995
Number of subsequent records: 3
Latest record: 2015

Aleochara (Xenochara) sparsa Heer, 1839
Pearce: no records
First record: 1975
Number of records: 5
Latest record: 2021

Aleochara (Xenochara) stichai Likovský, 1965
Pearce: no records
First record: 1986
Number of records: 1
Latest record: 1986

Aleochara (Xenochara) tristis
Gravenhorst, 1806
Pearce: 2 localities
First subsequent record: 2006
Number of subsequent records: 8
Latest record: 2021

Tinotus morion (Gravenhorst, 1802)
Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: 6
Latest record: 2010

Amarochara umbrosa (Erichson, 1837)
Pearce: 1 locality
No subsequent records

Phloeopora corticalis (Gravenhorst, 1802)
Pearce: no records
First record: 2018
Number of records: 1
Latest record: 2018

Phloeopora testacea (Mannerheim, 1830)
Pearce: no records
First record: 1970
Number of records: many
Latest record: 2019

Ilyobates bennetti Donisthorpe, 1914
Pearce: no records
First record: 1994
Number of records: 1
Latest record: 1994

Ilyobates nigricollis (Paykull, 1800)
Pearce: 2 localities
No subsequent records

Calodera aethiops (Gravenhorst, 1802)
Pearce: 3 localities
No subsequent records

Calodera riparia Erichson, 1837
Pearce: 1 locality
First subsequent record: 1984
Number of subsequent records: 7
Latest record: 2000

Calodera rubens Erichson, 1837

Pearce: 1 locality
No subsequent records

Tetralaucopora longitarsis (Erichson, 1839)

Pearce: 1 locality
First subsequent record: 1988
Number of subsequent records: 2
Latest record: 2017

Tetralaucopora rubicunda (Erichson, 1837)

Pearce: 1 locality
No subsequent records

Ocalea badia Erichson, 1837

Pearce: no records
First record: 2006
Number of records: 1
Latest record: 2006

Ocalea picata (Stephens, 1832)

Pearce: 1 locality
First subsequent record: 1986
Number of subsequent records: 6
Latest record: 2019

Ocalea rivularis Miller, L., 1851

Pearce: no records
First record: 1963
Number of records: 2
Latest record: 1974

Ocyusa maura (Erichson, 1837)

Pearce: no records
First record: 2011
Number of records: 5
Latest record: 2021

Ocyusa picina (Aubé, 1850)

Pearce: 2 localities
First subsequent record: 1989
Number of subsequent records: 9
Latest record: 2021

Hygropora cunctans (Erichson, 1837)

Pearce: 1 locality (1925)
No subsequent records

Oxypoda (Atlantoxypoda) lurida

Wollaston, 1857
Pearce: 1 locality
First subsequent record: 1998
Number of subsequent records: 1
Latest record: 1998

Oxypoda (Bessopora) annularis

Mannerheim, 1830
Pearce: no records
First record: 1979
Number of records: 4
Latest record: 2020

Oxypoda (Bessopora) brachyptera (Stephens, 1832)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 4
Latest record: 1979

Oxypoda (Bessopora) formiceticola Märkel, 1841

Pearce: no records
First record: 2017
Number of records: 1
Latest record: 2017

Oxypoda (Bessopora) haemorrhoea (Mannerheim, 1830)

Pearce: 2 localities
First subsequent record: 2013
Number of subsequent records: 2
Latest record: 2013

Oxypoda (Disochara) elongatula Aubé, 1850

Pearce: 3 localities
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2021

Oxypoda (Mycetodrepa) alternans (Gravenhorst, 1802)

Pearce: 1 locality
First subsequent record: 1968
Number of subsequent records: 7
Latest record: 2020

Oxypoda (Oxypoda) acuminata (Stephens, 1832)

Pearce: 1 locality
First subsequent record: 1968
Number of subsequent records: 5
Latest record: 2021

Oxypoda (Oxypoda) opaca (Gravenhorst, 1802)

Pearce: 2 localities
First subsequent record: 1971
Number of subsequent records: many
Latest record: 2021

Oxypoda (Oxypoda) spectabilis Märkel, 1844

Pearce: 1 locality (1894)
First subsequent record: 2002
Number of subsequent records: 2
Latest record: 2002

Oxypoda (Oxypoda) vittata Märkel, 1842

Pearce: 1 locality
First subsequent record: 2010
Number of subsequent records: 2
Latest record: 2016

Oxypoda (Podoxya) brevicornis (Stephens, 1832)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 3
Latest record: 1979

Oxypoda (Podoxya) carbonaria (Heer, 1841)

Pearce: 2 localities
No subsequent records

Oxypoda (Podoxya) exoleta Erichson, 1839

Pearce: 1 locality
First subsequent record: 2021
Number of subsequent records: 2
Latest record: 2021

Ischnoglossa turcica Wunderle, 1992

Pearce: 1 locality for *Ischnoglossa prolixa*, which could be this species
First record: 2014
Number of records: 2
Latest record: 2014

Dexiogyia corticina (Erichson, 1837)

Pearce: 1 locality (1904)
First subsequent record: 2001
Number of subsequent records: 3
Latest record: 2021

Crataraea suturalis (Mannerheim, 1830)

Pearce: 5 localities
First subsequent record: 1969
Number of subsequent records: 6
Latest record: 2021

Haploglossa gentilis (Märkel, 1844)

Pearce: 1 locality (1929)
First subsequent record: 2020
Number of subsequent records: 2
Latest record: 2020

Haploglossa marginalis (Gravenhorst, 1806)

Pearce: no records
First record: 1987
Number of records: 2
Latest record: 1988

Haploglossa nidicola (Fairmaire, 1853)

Pearce: 1 locality
No subsequent records

Haploglossa picipennis (Gyllenhal, 1827)

Pearce: no records
First record: 2021
Number of records: 1
Latest record: 2021

Haploglossa villosula (Stephens, 1832)

Pearce: 1 locality
First subsequent record: 1987
Number of subsequent records: many
Latest record: 2021

Thiasophila angulata (Erichson, 1837)

Pearce: no records
First record: 2014
Number of records: 7
Latest record: 2014

Thiasophila inquilina (Märkel, 1844)

Pearce: 2 localities
No subsequent records

Dinarda dentata (Gravenhorst, 1806)

Pearce: 1 locality
No subsequent records

Dinarda maerkeli Kiesenwetter, 1843

Pearce: no records
First record: 2014
Number of records: 9
Latest record: 2015

Meotica exilis (Knoch in Gravenhorst, 1806)

Pearce: 2 localities (1890, 1927)
First subsequent record: 2021
Number of subsequent records: 1
Latest record: 2021

Meotica exillima Sharp, 1915

Pearce: 1 locality
First subsequent record: 2001
Number of subsequent records: 3
Latest record: 2006

Meotica filiformis (Motschulsky, 1860)

Pearce: no records
First record: 1998
Number of records: 1
Latest record: 1998

Tachyusa concinna Heer, 1839

Pearce: no records
First record: 1998
Number of records: 2
Latest record: 1998

Tachyusa constricta Erichson, 1837

Pearce: no records
First record: 1998
Number of records: 3
Latest record: 2017

Ischnopoda leucopus (Marsham, 1802)

Pearce: no records
First record: 1988
Number of records: 1
Latest record: 1988

Ischnopoda umbratica (Erichson, 1837)

Pearce: 1 locality
No subsequent records

Thinonoma atra (Gravenhorst, 1806)

Pearce: 2 localities
First subsequent record: 1970
Number of subsequent records: 9
Latest record: 2020

Gnypeta carbonaria (Mannerheim, 1830)

Pearce: 2 localities
First subsequent record: 2017
Number of subsequent records: 1
Latest record: 2017

Gnypeta rubrior Tottenham, 1939

Pearce: no records
First record: 2013
Number of records: 3
Latest record: 2020

Dilacra luteipes (Erichson, 1837)

Pearce: 2 localities
First subsequent record: 1997
Number of subsequent records: 7
Latest record: 2020

Dilacra vilis (Erichson, 1837)

Pearce: no records
First record: 2005
Number of records: 3
Latest record: 2020

Dacrila fallax (Kraatz, 1856)

Pearce: no records
First record: 1977
Number of records: 2
Latest record: 2005

Cypha discoidea (Erichson, 1839)

Pearce: 1 locality (1925)
First subsequent record: 1999
Number of subsequent records: 2
Latest record: 2021

Cypha laeviuscula (Mannerheim, 1830)

Pearce: 1 locality
No subsequent records

Cypha longicornis (Paykull, 1800)

Pearce: 4 localities
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2021

Cypha punctum (Motschulsky, 1857)

Pearce: no records
First record: 1975
Number of records: 2
Latest record: 1993

Cypha seminulum (Erichson, 1839)

Pearce: 3 localities
No subsequent records

Oligota apicata Erichson, 1837

Pearce: 2 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2019

Oligota granaria Erichson, 1837

Pearce: no records
First record: 2020
Number of records: 1
Latest record: 2020

Oligota inflata (Mannerheim, 1830)

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: 4
Latest record: 2021

Oligota parva Kraatz, 1862

Pearce: no records
First record: 1996
Number of records: 2
Latest record: 2020

Oligota picipes (Stephens, 1832)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Oligota pumilio Kiesenwetter, 1858

Pearce: no records
First record: 2013
Number of records: 1
Latest record: 2013

Oligota punctulata Heer, 1839

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Oligota pusillima (Gravenhorst, 1806)

Pearce: 2 localities (1896, 1919)
First subsequent record: 2000
Number of subsequent records: 3
Latest record: 2008

Myllaena brevicornis (Matthews, A.H., 1838)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Myllaena dubia (Gravenhorst, 1806)

Pearce: 2 localities
First subsequent record: 1994
Number of subsequent records: 6
Latest record: 2016

Myllaena elongata (Matthews, A.H., 1838)

Pearce: no records
First record: 2020
Number of records: 1
Latest record: 2020

Myllaena gracilis (Matthews, A.H., 1838)

Pearce: no records
First record: 2015
Number of records: 1
Latest record: 2015

Myllaena infuscata Kraatz, 1853

Pearce: 1 locality
First subsequent record: 2003
Number of subsequent records: 6
Latest record: 2019

Myllaena intermedia Erichson, 1837

Pearce: 2 localities
First subsequent record: 2000
Number of subsequent records: many
Latest record: 2019

Myllaena kraatzi Sharp, 1871

Pearce: 1 locality
First subsequent record: 2016
Number of subsequent records: 3
Latest record: 2018

Myllaena minuta (Gravenhorst, 1806)

Pearce: no records
First record: 2001
Number of records: 4
Latest record: 2019

Diglotta mersa (Haliday, 1837)

Pearce: 1 locality
First subsequent record: 2006
Number of subsequent records: 2
Latest record: 2019

Diglotta sinuaticollis (Mulsant & Rey, 1870)

Pearce: 5 localities
First subsequent record: 2000
Number of subsequent records: 1
Latest record: 2000

Autalia impressa (Olivier, 1795)

Pearce: 3 localities (but may include *longicornis*)
First subsequent record: 1975
Number of subsequent records: 9
Latest record: 2019

Autalia longicornis Scheerpeltz, 1947

Pearce: no records (but see *impressa*)
First record: 1986
Number of records: 5
Latest record: 2019

Autalia rivularis (Gravenhorst, 1802)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Pseudopasilia testacea (Brisout de Barneville, C., 1863)
Pearce: 2 localities
No subsequent records

Leptusa (Dendroleptusa) fumida (Erichson, 1839)
Pearce: 2 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2021

Leptusa (Leptusa) pulchella (Mannerheim, 1830)
Pearce: no records
First record: 2017
Number of records: 3
Latest record: 2019

Leptusa (Pachygluta) ruficollis (Erichson, 1839)
Pearce: 2 localities
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2021

Heterota plumbea (Waterhouse, G.R., 1858)
Pearce: 8 localities
First subsequent record: 1966
Number of subsequent records: 9
Latest record: 2020

Bolitochara bella Märkel, 1844
Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: 6
Latest record: 2019

Bolitochara obliqua Erichson, 1837
Pearce: no records
First record: 2018
Number of records: 2
Latest record: 2018

Bolitochara tecta Assing, 2014 [2012: *Bolitochara lucida*]
Pearce: 1 locality (1874)
First subsequent record: 2018
Number of subsequent records: 1
Latest record: 2018

Encephalus complicans Stephens, 1832
Pearce: 1 locality
First subsequent record: 1997
Number of subsequent records: many
Latest record: 2017

Gyrophæna (Gyrophæna) bihamata Thomson, C.G., 1867
Pearce: no records
First record: 2006
Number of records: 4
Latest record: 2018

Gyrophæna (Gyrophæna) fasciata (Marsham, 1802)
Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: 5
Latest record: 2020

Gyrophæna (Gyrophæna) gentilis Erichson, 1839
Pearce: 2 localities
First subsequent record: 1971
Number of subsequent records: 2
Latest record: 2018

Gyrophæna (Gyrophæna) joyi Wendeler, 1924
Pearce: no records
First record: 1975
Number of records: 3
Latest record: 2019

Gyrophæna (Gyrophæna) lucidula Erichson, 1837
Pearce: 1 locality (1890)
First subsequent record: 2020
Number of subsequent records: 1
Latest record: 2020

Gyrophæna (Gyrophæna) manca Erichson, 1839
Pearce: no records
First record: 2000
Number of records: 2
Latest record: 2019

Gyrophæna (Gyrophæna) minima Erichson, 1837
Pearce: 1 locality (1891)
First subsequent record: 2000
Number of subsequent records: 2
Latest record: 2015

Gyrophæna (Gyrophæna) nana (Paykull, 1800)

Pearce: 1 locality

First subsequent record: 2019

Number of subsequent records: 1

Latest record: 2019

Gyrophæna (Leptarthrophæna) affinis Mannerheim, 1830

Pearce: 3 localities

First subsequent record: 1994

Number of subsequent records: 6

Latest record: 2014

Agaricochara latissima (Stephens, 1832)

Pearce: no records

First record: 2000

Number of records: many

Latest record: 2021

Homalota plana (Gyllenhal, 1810)

Pearce: 2 localities

First subsequent record: 1999

Number of subsequent records: 1

Latest record: 1999

Anomognathus cuspidatus (Erichson, 1839)

Pearce: 1 locality

First subsequent record: 1991

Number of subsequent records: 5

Latest record: 2020

Silusa rubiginosa Erichson, 1837

Pearce: no records

First record: 2004

Number of records: 1

Latest record: 2004

Hygronoma dimidiata (Gravenhorst, 1806)

Pearce: 6 localities

First subsequent record: 1989

Number of subsequent records: many

Latest record: 2021

Placusa pumilio (Gravenhorst, 1802)

Pearce: no records

First record: 2004

Number of records: 1

Latest record: 2004

Placusa tachyporoides (Waltl, 1838)

Pearce: 1 locality (1892)

No subsequent records

Actocharis readingii Sharp, 1870

Pearce: no records

First record: 1930

Number of records: 1

Latest record: 1930

Phytosus (Actosus) balticus Kraatz, 1859

Pearce: 3 localities

First subsequent record: 2004

Number of subsequent records: 3

Latest record: 2016

Phytosus (Actosus) nigriventris (Chevrolat, 1843)

Pearce: 4 localities

No subsequent records

Phytosus (Phytosus) spinifer Curtis, 1838

Pearce: 4 localities

First subsequent record: 2004

Number of subsequent records: 4

Latest record: 2019

Arena tabida (Kiesenwetter, 1850)

Pearce: 2 localities

No subsequent records

Schistoglossa gemina (Erichson, 1837)

Pearce: no records

First record: 2021

Number of records: 1

Latest record: 2021

Boreophilia eremita (Rye, 1866)

Pearce: no records

First record: 1998

Number of records: 1

Latest record: 1998

Hydrosmehta longula (Heer, 1839)

Pearce: 1 locality

No subsequent records

Amischa analis (Gravenhorst, 1802)

Pearce: 6 localities

First subsequent record: 1970

Number of subsequent records: many

Latest record: 2021

Amischa bifoveolata (Mannerheim, 1830)

Pearce: no records

First record: 1975

Number of records: many

Latest record: 2006

Amischa decipiens (Sharp, 1869)

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Amischa forcipata Mulsant & Rey, 1873

Pearce: no records
First record: 1979
Number of records: 7
Latest record: 2019

Amischa nigrofusca (Stephens, 1832)

Pearce: no records
First record: 1998
Number of records: many
Latest record: 2021

Nehemitropia lividipennis (Mannerheim, 1830)

Pearce: 3 localities
First subsequent record: 1971
Number of subsequent records: many
Latest record: 2013

Notothecta flavipes (Gravenhorst, 1806)

Pearce: no records
First record: 1996
Number of records: 1
Latest record: 1996

Brundinia marina (Mulsant & Rey, 1853)

Pearce: 1 locality (1886)
First subsequent record: 2021
Number of subsequent records: 1
Latest record: 2021

Brundinia meridionalis (Mulsant & Rey, 1853)

Pearce: 1 locality
No subsequent records

Ousipalia caesula (Erichson, 1839)

Pearce: 2 localities
No subsequent records

Dinaraea aequata (Erichson, 1837)

Pearce: 2 localities
First subsequent record: 2000
Number of subsequent records: many
Latest record: 2021

Dinaraea angustula (Gyllenhal, 1810)

Pearce: 1 locality (1925)
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2021

Dinaraea linearis (Gravenhorst, 1802)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 1
Latest record: 1975

Dadobia immersa (Erichson, 1837)

Pearce: 1 locality
No subsequent records

Plataraea brunnea (Fabricius, 1798)

Pearce: 1 locality
No subsequent records

Liogluta longiuscula (Gravenhorst, 1802)

Pearce: 2 localities
First subsequent record: 2014
Number of subsequent records: 5
Latest record: 2021

Liogluta microptera Thomson, C.G., 1867

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Atheta (Alaobia) gagatina (Baudi di Selve, 1848)

Pearce: 1 locality (1891)
First subsequent record: 1993
Number of subsequent records: 2
Latest record: 1993

Atheta (Alaobia) pallidicornis (Thomson, C.G., 1856)

Pearce: no records
First record: 2017
Number of records: 1
Latest record: 2017

Atheta (Alaobia) scapularis (Sahlberg, C.R., 1831)

Pearce: 2 localities (1874, 1925)
First subsequent record: 1975
Number of subsequent records: 3
Latest record: 2017

Atheta (Alaobia) trinotata (Kraatz, 1856)

Pearce: 3 localities

First subsequent record: 1975

Number of subsequent records: 8

Latest record: 2021

Atheta (Anopleta) corvina (Thomson, C.G., 1856)

Pearce: no records

First record: 1966

Number of records: 3

Latest record: 2019

Atheta (Atheta) aeneicollis (Sharp, 1869)

Pearce: 1 locality (1892)

First subsequent record: 1975

Number of subsequent records: 6

Latest record: 2004

Atheta (Atheta) aquatica (Thomson, C.G., 1852)

Pearce: 4 localities

First subsequent record: 2004

Number of subsequent records: 5

Latest record: 2021

Atheta (Atheta) autumnalis (Erichson, 1839)

Pearce: 1 locality

No subsequent records

Atheta (Atheta) basicornis (Mulsant & Rey, 1852)

Pearce: no records

First record: 2013

Number of records: 3

Latest record: 2021

Atheta (Atheta) britanniae Bernhauer & Scheerpeltz, 1926

Pearce: no records

First record: 1993

Number of records: 3

Latest record: 2017

Atheta (Atheta) brunneipennis (Thomson, C.G., 1852)

Pearce: no records

First record: 1975

Number of records: 1

Latest record: 1975

Atheta (Atheta) castanoptera (Mannerheim, 1830)

Pearce: 1 locality

First subsequent record: 1968

Number of subsequent records: 6

Latest record: 2020

Atheta (Atheta) crassicornis (Fabricius, 1792)

Pearce: no records

First record: 1975

Number of records: many

Latest record: 2021

Atheta (Atheta) divisa (Märkel, 1844)

Pearce: 1 locality

First subsequent record: 1975

Number of subsequent records: 2

Latest record: 2021

Atheta (Atheta) fungicola (Thomson, C.G., 1852)

Pearce: 3 localities

First subsequent record: 1975

Number of subsequent records: 3

Latest record: 1998

Atheta (Atheta) graminicola (Gravenhorst, 1806)

Pearce: 4 localities

First subsequent record: 1969

Number of subsequent records: many

Latest record: 2021

Atheta (Atheta) harwoodi Williams, 1930

Pearce: no records

First record: 1989

Number of records: 1

Latest record: 1989

Atheta (Atheta) hypnorum (Kiesenwetter, 1850)

Pearce: 3 localities (1874, 1892, 1919)

First subsequent record: 1982

Number of subsequent records: 1

Latest record: 1982

Atheta (Atheta) oblita (Erichson, 1839)

Pearce: 2 localities

First subsequent record: 1975

Number of subsequent records: 4

Latest record: 2021

Atheta (Atheta) ravilla (Erichson, 1839)

Pearce: 1 locality (1890)
First subsequent record: 1975
Number of subsequent records: 5
Latest record: 2020

Atheta (Atheta) triangulum (Kraatz, 1856)

Pearce: no records
First record: 1979
Number of records: many
Latest record: 2021

Atheta (Atheta) vaga (Heer, 1839)

Pearce: no records
First record: 2013
Number of records: 3
Latest record: 2021

Atheta (Atheta) xanthopus (Thomson, C.G., 1856)

Pearce: 2 localities
First subsequent record: 2006
Number of subsequent records: 3
Latest record: 2020

Atheta (Badura) cauta (Erichson, 1837)
[2012: *Dimetrota cauta*]

Pearce: no records
First record: 1975
Number of records: 2
Latest record: 2015

Atheta (Bessobia) excellens (Kraatz, 1856)

Pearce: no records
First record: 1979
Number of records: 1
Latest record: 1979

Atheta (Bessobia) fungivora (Thomson, C.G., 1867)

Pearce: 1 locality
First subsequent record: 2000
Number of subsequent records: 4
Latest record: 2012

Atheta (Bessobia) occulta (Erichson, 1837)

Pearce: no records
First record: 2004
Number of records: 1
Latest record: 2004

Atheta (Ceritaxa) subterranea (Mulsant & Rey, 1853) [2012: *Neohilara subterranea*]

Pearce: no records
First record: 1993
Number of records: 1
Latest record: 1993

Atheta (Chaetida) longicornis
(Gravenhorst, 1802)

Pearce: 2 localities (1890, 1910)
First subsequent record: 1970
Number of subsequent records: 6
Latest record: 2008

Atheta (Coprothassa) melanaria
(Mannerheim, 1830)

Pearce: 1 locality
First subsequent record: 2011
Number of subsequent records: 3
Latest record: 2019

Atheta (Datomicra) celata (Erichson, 1837)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 5
Latest record: 2020

Atheta (Datomicra) dadopora Thomson, C.G., 1867

Pearce: no records
First record: 2004
Number of records: 2
Latest record: 2009

Atheta (Datomicra) nigra (Kraatz, 1856)

Pearce: 2 localities
First subsequent record: 1971
Number of subsequent records: many
Latest record: 2017

Atheta (Datomicra) sordidula (Erichson, 1837)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 3
Latest record: 1999

Atheta (Datomicra) zosteræ (Thomson, C.G., 1856)

Pearce: no records
First record: 2006
Number of records: 1
Latest record: 2006

Atheta (Dimetrota) atramentaria
(Gyllenhal, 1810)
Pearce: 6 localities
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2019

Atheta (Dimetrota) marcida (Erichson,
1837)
Pearce: 1 locality
First subsequent record: 2006
Number of subsequent records: 5
Latest record: 2015

Atheta (Dimetrota) nigripes (Thomson,
C.G., 1856)
Pearce: no records
First record: 1975
Number of records: 4
Latest record: 2021

Atheta (Dimetrota) setigera (Sharp, 1869)
Pearce: 1 locality
No subsequent records

Atheta (Dimetrota) laticollis (Stephens,
1832) [2012: *Mycetota laticollis*]
Pearce: 2 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Atheta (Microdota) aegra (Heer, 1841)
Pearce: no records
First record: 2017
Number of records: 1
Latest record: 2017

Atheta (Microdota) amicula (Stephens,
1832)
Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 5
Latest record: 2020

Atheta (Microdota) atricolor (Sharp, 1869)
Pearce: 1 locality
No subsequent records

Atheta (Microdota) benickiella Brundin,
1948
Pearce: no records
First record: 1993
Number of records: 1
Latest record: 1993

Atheta (Microdota) indubia (Sharp, 1869)
Pearce: no records
First record: 1993
Number of records: 4
Latest record: 2021

Atheta (Microdota) minuscula (Brisout de
Barneville, C., 1860)
Pearce: no records
First record: 1975
Number of records: 2
Latest record: 1999

Atheta (Mocyta) amplicollis (Mulsant &
Rey, 1873)
Pearce: no records
First record: 1998
Number of records: many
Latest record: 2020

Atheta (Mocyta) clientula (Erichson, 1839)
Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2021

Atheta (Mocyta) fungi (Gravenhorst, 1806)
Pearce: 8 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Atheta (Mocyta) orbata (Erichson, 1837)
Pearce: 2 localities
First subsequent record: 2021
Number of subsequent records: 2
Latest record: 2021

Atheta (Parameotica) difficilis (Brisout de
Barneville, C., 1860)
Pearce: no records
First record: 2006
Number of records: 2
Latest record: 2017

Atheta (Philhygra) elongatula
(Gravenhorst, 1802)
Pearce: 1 locality
First subsequent record: 1989
Number of subsequent records: many
Latest record: 2019

Atheta (Philhygra) fallaciosa (Sharp,
1869)
Pearce: 1 locality
No subsequent records

Atheta (Philhygra) gyllenhalii (Thomson, C.G., 1856)

Pearce: no records
First record: 2021
Number of records: 1
Latest record: 2021

Atheta (Philhygra) hygrobica (Thomson, C.G., 1856)

Pearce: no records
First record: 2004
Number of records: 1
Latest record: 2004

Atheta (Philhygra) hygrotopora (Kraatz, 1856)

Pearce: no records
First record: 2013
Number of records: 1
Latest record: 2013

Atheta (Philhygra) luridipennis (Mannerheim, 1830)

Pearce: no records
First record: 2020
Number of records: 1
Latest record: 2020

Atheta (Philhygra) malleus Joy, 1913

Pearce: 2 localities
First subsequent record: 2006
Number of subsequent records: many
Latest record: 2018

Atheta (Philhygra) melanocera (Thomson, C.G., 1856)

Pearce: 1 locality
First subsequent record: 1989
Number of subsequent records: 2
Latest record: 2006

Atheta (Philhygra) obtusangula Joy, 1913

Pearce: 1 locality
First subsequent record: 2000
Number of subsequent records: 1
Latest record: 2000

Atheta (Philhygra) palustris (Kiesenwetter, 1844)

Pearce: 3 localities
First subsequent record: 2019
Number of subsequent records: 1
Latest record: 2019

Atheta (Philhygra) parca (Mulsant & Rey, 1873)

Pearce: 1 locality (1928)
No subsequent records

Atheta (Philhygra) volans (Scriba, 1859)

Pearce: 1 locality
First subsequent record: 2006
Number of subsequent records: 2
Latest record: 2006

Atheta (Rhagocneme) subsinuata (Erichson, 1839)

Pearce: 1 locality
No subsequent records

Atheta (Tetropla) liturata (Stephens, 1832)

Pearce: 1 locality (1925)
First subsequent record: 1975
Number of subsequent records: 4
Latest record: 2020

Atheta (Tetropla) nigritula (Gravenhorst, 1802)

Pearce: 1 locality (1890)
First subsequent record: 1968
Number of subsequent records: 2
Latest record: 1996

Atheta (Thinobaena) vestita (Gravenhorst, 1806)

Pearce: 5 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2020

Atheta (Traumoecia) taxiceroides Munster, 1932

Pearce: no records
First record: 2020
Number of records: 1
Latest record: 2020

Acrotone aterrima (Gravenhorst, 1802)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 7
Latest record: 2015

Acrotone benicki Allen, 1940

Pearce: no records
First record: 1937
Number of records: 2
Latest record: 2015

Acrotona exigua (Erichson, 1837)

Pearce: 1 locality
No subsequent records

Acrotona muscorum (Brisout de Barneville, C., 1860)

Pearce: no records
First record: 2003
Number of records: 4
Latest record: 2020

Acrotona obfuscata (Gravenhorst, 1802)

Pearce: 1 locality
First subsequent record: 2021
Number of subsequent records: 1
Latest record: 2021

Acrotona parvula (Mannerheim, 1830)

Pearce: no records
First record: 1975
Number of records: 3
Latest record: 2015

Acrotona pygmaea (Gravenhorst, 1802)

Pearce: 2 localities
First subsequent record: 2003
Number of subsequent records: 4
Latest record: 2006

Adota maritima (Mannerheim, 1843)

Pearce: no records
First record: 1976
Number of records: 2
Latest record: 2004

Dalotia coriaria (Kraatz, 1856)

Pearce: no records
First record: 1969
Number of records: 5
Latest record: 2020

Eurodotina inquinula (Gravenhorst, 1802)

Pearce: no records
First record: 2021
Number of records: 1
Latest record: 2021

Trichiusa immigrata Lohse, 1984

Pearce: no records
First record: 1995
Number of records: 5
Latest record: 2006

Alianta incana (Erichson, 1837)

Pearce: 1 locality
First subsequent record: 1964
Number of subsequent records: 7
Latest record: 2021

Pachnida nigella (Erichson, 1837)

Pearce: 1 locality
First subsequent record: 1998
Number of subsequent records: 3
Latest record: 2014

Thamiaraea cinnamomea (Gravenhorst, 1802)

Pearce: no records
First record: 1998
Number of records: 3
Latest record: 2019

Thamiaraea hospita (Märkel, 1844)

Pearce: 1 locality
No subsequent records

Halobrecta algae (Hardy, 1851)

Pearce: 3 localities
No subsequent records

Halobrecta algophila (Fenyés, 1909)

Pearce: no records
First record: 2017
Number of records: 1
Latest record: 2017

Halobrecta flavipes Thomson, C.G., 1861

Pearce: 5 localities
First subsequent record: 2019
Number of subsequent records: 1
Latest record: 2019

Callicerus obscurus Gravenhorst, 1802

Pearce: 2 localities
First subsequent record: 1979
Number of subsequent records: 2
Latest record: 2015

Callicerus rigidicornis (Erichson, 1839)

Pearce: 1 locality
No subsequent records

Aloconota (Aloconota) gregaria (Erichson, 1839)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2018

Aloconota (Aloconota) sulcifrons
(Stephens, 1832)
Pearce: 3 localities
No subsequent records

Aloconota (Disopora) longicollis (Mulsant
& Rey, 1852)
Pearce: 1 locality
First subsequent record: 2004
Number of subsequent records: 1
Latest record: 2004

Geostiba circellaris Gravenhorst 1806
Pearce: 2 localities
First subsequent record: 1971
Number of subsequent records: many
Latest record: 2021

Alevonota elegantula (Brisout de
Barneville, C., 1863)
Pearce: 1 locality
No subsequent records

Alevonota gracilentia (Erichson, 1839)
Pearce: no records
First record: 2017
Number of records: 1
Latest record: 2017

Drusilla canaliculata Fabricius 1787
Pearce: generally common
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Zyras collaris (Paykull, 1800)
Pearce: 3 localities
First subsequent record: 1945
Number of subsequent records: 1
Latest record: 1945

Zyras haworthi (Stephens, 1835)
Pearce: 2 localities (1884, 1929)
First subsequent record: 1987
Number of subsequent records: 2
Latest record: 1993

Pella cognata (Märkel, 1842)
Pearce: 3 localities
First subsequent record: 1980
Number of subsequent records: 3
Latest record: 2017

Pella funesta (Gravenhorst, 1806)
Pearce: 1 locality (1925)
First subsequent record: 1975
Number of subsequent records: 1
Latest record: 1975

Pella humeralis (Gravenhorst, 1802)
Pearce: 1 locality
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2015

Pella laticollis (Märkel, 1844)
Pearce: 1 locality (1929)
First record: 1975
Number of records: 1
Latest record: 1975

Pella limbata (Paykull, 1789)
Pearce: 7 localities
First subsequent record: 1975
Number of subsequent records: 3
Latest record: 2019

Lomechusa emarginata (Paykull, 1789)
Pearce: 1 locality
First subsequent record: 1932
Number of subsequent records: many
Latest record: 2016

Lomechusa paradoxa Gravenhorst, 1806
Pearce: no records
First record: 2001
Number of records: 2
Latest record: 2019

Cordalia obscura (Gravenhorst, 1802)
Pearce: 5 localities
First subsequent record: 1989
Number of subsequent records: many
Latest record: 2021

Falagria caesa Erichson, 1837
Pearce: 1 locality
First subsequent record: 1995
Number of subsequent records: many
Latest record: 2021

Falagria sulcatula (Gravenhorst, 1806)
Pearce: 2 localities
First subsequent record: 2000
Number of subsequent records: 2
Latest record: 2007

Falagrioma thoracica (Stephens, 1832)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: 8
Latest record: 2016

Myrmecopora brevipes Butler, 1909

Pearce: 2 localities
No subsequent records

Myrmecopora oweni Assing, 1997

Pearce: no records
First record: 1996
Number of records: 4
Latest record: 1998

Myrmecopora sulcata (Kiesenwetter, 1850)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: 6
Latest record: 2005

Myrmecopora uvida (Erichson, 1840)

Pearce: 7 localities
First subsequent record: 1975
Number of subsequent records: 2
Latest record: 1991

Scaphidium quadrimaculatum Olivier, 1790

Pearce: 4 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Scaphisoma agaricinum (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 2011
Number of subsequent records: 8
Latest record: 2021

Scaphisoma boleti (Panzer, 1793)

Pearce: 1 locality
First subsequent record: 1971
Number of subsequent records: 2
Latest record: 2014

Siagonium quadricorne Kirby & Spence, 1815

Pearce: 1 locality (1921)
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2019

Deleaster dichrous (Gravenhorst, 1802)

Pearce: no records
First record: 1988
Number of records: 6
Latest record: 2013

Coprophilus striatulus (Fabricius, 1792)

Pearce: 1 locality
No subsequent records

Manda mandibularis (Gyllenhal, 1827)

Pearce: 1 locality
No subsequent records

Syntomium aeneum (Müller, P.W.J., 1821)

Pearce: 1 locality
No subsequent records

Bledius (Bargus) fuscipes Rye, 1865

Pearce: 1 locality (1921)
First subsequent record: 2000
Number of subsequent records: 1
Latest record: 2000

Bledius (Bargus) longulus Erichson, 1839

Pearce: 1 locality
First subsequent record: 1989
Number of subsequent records: 4
Latest record: 2006

Bledius (Bargus) opacus (Block, 1799)

Pearce: 1 locality
First subsequent record: 1968
Number of subsequent records: 2
Latest record: 2005

Bledius (Bargus) pallipes (Gravenhorst, 1806)

Pearce: 1 locality (1925)
First subsequent record: 1953
Number of subsequent records: 4
Latest record: 2005

Bledius (Bledius) frisius Lohse, 1978

Pearce: 3 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2015

Bledius (Bledius) spectabilis Kraatz, 1857

Pearce: no records
First record: 1979
Number of records: many
Latest record: 2016

Bledius (Bledius) tricornis (Herbst, 1784)

Pearce: 1 locality
First subsequent record: 2004
Number of subsequent records: 1
Latest record: 2004

Bledius (Bledius) unicornis (Germar, 1825)

Pearce: 1 locality
First subsequent record: 1996
Number of subsequent records: 2
Latest record: 1997

Bledius (Dicarenus) fergussoni Joy, 1912

Pearce: 5 localities
No subsequent records

Bledius (Dicarenus) subniger Schneider, O., 1898

Pearce: 2 localities (1910, 1928)
No subsequent records

Bledius (Hesperophilus) atricapillus (Germar, 1825)

Pearce: 1 locality
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2019

Bledius (Hesperophilus) femoralis (Gyllenhal, 1827)

Pearce: no records
First record: 1955
Number of records: 3
Latest record: 2001

Bledius (Hesperophilus) gallicus (Gravenhorst, 1806)

Pearce: no records
First record: 2013
Number of records: 1
Latest record: 2013

Ochtheophilus angustior (Bernhauer, 1943)

Pearce: no records
First record: 2020
Number of records: 1
Latest record: 2020

Ochtheophilus aureus (Fauvel, 1871)

Pearce: no records
First record: 1997
Number of records: 5
Latest record: 2020

Teropalpus unicolor (Sharp, 1900)

Pearce: 1 locality (1912)
First subsequent record: 1966
Number of subsequent records: 5
Latest record: 2017

Thinodromus arcuatus (Stephens, 1834)

Pearce: "Dorset" (A. Ford)
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2021

Carpelimus (Carpelimus) fuliginosus (Gravenhorst, 1802)

Pearce: no records
First record: 2013
Number of records: 1
Latest record: 2013

Carpelimus (Carpelimus) obesus (Kiesenwetter, 1844)

Pearce: no records
First record: 2006
Number of records: 1
Latest record: 2006

Carpelimus (Carpelimus) pusillus (Gravenhorst, 1802)

Pearce: 1 locality
First subsequent record: 1993
Number of subsequent records: 5
Latest record: 2016

Carpelimus (Paratrogophloeus) erichsoni (Sharp, 1871)

Pearce: 6 localities for *bilineatus* which could be either species
First subsequent record: 1949
Number of subsequent records: 9
Latest record: 2018

Carpelimus (Paratrogophloeus) rivularis (Motschulsky, 1860)

Pearce: 1 locality
First subsequent record: 1949
Number of subsequent records: many
Latest record: 2019

Carpelimus (Paratrogophloeus) similis (Smetana, 1967)

Pearce: no records
First record: 1989
Number of records: 1
Latest record: 1989

Carpelimus (Troginus) zealandicus
(Sharp, 1900)

Pearce: no records
First record: 1975
Number of records: 2
Latest record: 2005

Carpelimus (Trogophloeus) corticinus
(Gravenhorst, 1806)

Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Carpelimus (Trogophloeus) elongatulus
(Erichson, 1839)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Carpelimus (Trogophloeus) gracilis
(Mannerheim, 1830)

Pearce: no records
First record: 1993
Number of records: 5
Latest record: 2021

Carpelimus (Trogophloeus) halophilus
(Kiesenwetter, 1844)

Pearce: 4 localities
No subsequent records

Carpelimus (Trogophloeus) subtilis
(Erichson, 1839)

Pearce: no records
First record: 2017
Number of records: 1
Latest record: 2017

Thinobius brevipennis Kiesenwetter, 1850

Pearce: no records
First record: 1990
Number of records: 2
Latest record: 1997

Aploderus caelatus (Gravenhorst, 1802)

Pearce: 1 locality
No subsequent records

Platystethus (Craetopycrus) alutaceus
Thomson, C.G., 1861

Pearce: no records
First record: 2014
Number of records: 7
Latest record: 2018

Platystethus (Craetopycrus) capito Heer,
1839

Pearce: 1 locality
First subsequent record: 2002
Number of subsequent records: 2
Latest record: 2008

Platystethus (Craetopycrus) cornutus
(Gravenhorst, 1802)

Pearce: 2 localities
First subsequent record: 1966
Number of subsequent records: many
Latest record: 2021

Platystethus (Craetopycrus) nitens
(Sahlberg, C.R., 1832)

Pearce: no records
First record: 1986
Number of records: many
Latest record: 2021

Platystethus (Craetopycrus) nodifrons
Mannerheim, 1830

Pearce: 1 locality
First subsequent record: 1963
Number of subsequent records: 2
Latest record: 2008

Platystethus (Platystethus) arenarius
(Geoffroy in Fourcroy, 1785)

Pearce: 5 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Anotylus complanatus (Erichson, 1839)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: 8
Latest record: 2020

Anotylus insecatus (Gravenhorst, 1806)

Pearce: 2 localities
No subsequent records

Anotylus inustus (Gravenhorst, 1806)

Pearce: 2 localities
First subsequent record: 1995
Number of subsequent records: many
Latest record: 2021

Anotylus maritimus Thomson, C.G., 1861

Pearce: 2 localities
First subsequent record: 1977
Number of subsequent records: 3
Latest record: 2019

Anotylus nitidulus (Gravenhorst, 1802)

Pearce: 3 localities
First subsequent record: 2008
Number of subsequent records: 6
Latest record: 2020

Anotylus rugosus (Fabricius, 1775)

Pearce: 5 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Anotylus sculpturatus (Gravenhorst, 1806)

Pearce: generally common
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Anotylus tetracarinatus (Block, 1799)

Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Oxytelus (Epomotylus) sculptus

Gravenhorst, 1806
Pearce: 2 localities
First subsequent record: 1990
Number of subsequent records: many
Latest record: 2021

Oxytelus (Oxytelus) migrator Fauvel, 1904

Pearce: no records
First record: 2003
Number of records: 2
Latest record: 2003

Oxytelus (Tanykraerus) laqueatus
(Marshall, 1802)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Oxyporus rufus (Linnaeus, 1758)

Pearce: 7 localities
First subsequent record: 2011
Number of subsequent records: 1
Latest record: 2011

Eutheia schaumii Kiesenwetter, 1858

Pearce: no records
First record: 2003
Number of records: 2
Latest record: 2003

Eutheia scydmaenoides Stephens, 1830

Pearce: no records
First record: 2003
Number of records: 2
Latest record: 2003

Cephennium gallicum Ganglbauer, 1899

Pearce: 7 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2020

Neuraphes angulatus (Müller, P.W.J. &
Kunze, 1822)

Pearce: 1 locality (1925)
First subsequent record: 1979
Number of subsequent records: 9
Latest record: 2016

Neuraphes elongatulus (Müller, P.W.J. &
Kunze, 1822)

Pearce: 1 locality
First subsequent record: 1987
Number of subsequent records: 1
Latest record: 1987

Neuraphes plicicollis (Reitter, 1880)

Pearce: no records
First record: 2019
Number of records: 4
Latest record: 2019

Neuraphes praeteritus (Rye, 1872)

Pearce: no records
First record: 2018
Number of records: 1
Latest record: 2018

Stenichnus bicolor (Denny, 1825)

Pearce: no records
First record: 2018
Number of records: 2
Latest record: 2019

Stenichnus collaris (Müller, P.W.J. &
Kunze, 1822)

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Stenichnus poweri (Fowler, 1884)

Pearce: no records
First record: 1998
Number of records: 4
Latest record: 2000

Stenichnus pusillus (Müller, P.W.J. & Kunze, 1822)

Pearce: 4 localities
First subsequent record: 2002
Number of subsequent records: many
Latest record: 2020

Stenichnus scutularis (Müller, P.W.J. & Kunze, 1822)

Pearce: 2 localities (1899, 1901)
First subsequent record: 2002
Number of subsequent records: many
Latest record: 2021

Euconnus fimetarius (Chaudoir, 1845)

Pearce: 3 localities
First subsequent record: 1979
Number of subsequent records: 6
Latest record: 2014

Euconnus hirticollis (Illiger, 1798)

Pearce: 2 localities
First subsequent record: 2006
Number of subsequent records: 3
Latest record: 2017

Scydmaenus rufus Müller, P.W.J. & Kunze, 1822

Pearce: no records
First record: 2003
Number of records: 3
Latest record: 2004

Scydmaenus tarsatus Müller, P.W.J. & Kunze, 1822

Pearce: 4 localities
First subsequent record: 1954
Number of subsequent records: 8
Latest record: 2020

Dianous coerulescens (Gyllenhal, 1810)

Pearce: "Dorset" (A. Ford)
First subsequent record: 2017
Number of subsequent records: 2
Latest record: 2020

Stenus (Hemistenus) aceris Stephens, 1833

Pearce: 5 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Stenus (Hemistenus) fuscicornis Erichson, 1840

Pearce: no records
First record: 1975
Number of records: 3
Latest record: 1996

Stenus (Hemistenus) geniculatus Gravenhorst, 1806

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2020

Stenus (Hemistenus) impressus Germar, 1824

Pearce: generally common and widely distributed
First subsequent record: 1960
Number of subsequent records: many
Latest record: 2021

Stenus (Hemistenus) ochropus Kiesenwetter, 1858

Pearce: 1 locality
First subsequent record: 1999
Number of subsequent records: 3
Latest record: 2019

Stenus (Hemistenus) ossium Stephens, 1833

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Stenus (Hemistenus) pallipes Gravenhorst, 1802

Pearce: no records
First record: 2005
Number of records: 5
Latest record: 2014

Stenus (Hemistenus) subaeneus Erichson, 1840

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Stenus (Hypostenus) cicindeloides (Schaller, 1783)

Pearce: 1 locality (1898)
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Stenus (Hypostenus) fornicatus Stephens, 1833

Pearce: no records
First record: 2005
Number of records: 6
Latest record: 2019

Stenus (Hypostenus) fulvicornis Stephens, 1833

Pearce: 1 locality (1898)
First subsequent record: 1967
Number of subsequent records: many
Latest record: 2021

Stenus (Hypostenus) kiesenwetteri Rosenhauer, 1856

Pearce: 1 locality (1892)
First subsequent record: 2000
Number of subsequent records: 8
Latest record: 2019

Stenus (Hypostenus) latifrons Erichson, 1839

Pearce: 6 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2021

Stenus (Hypostenus) similis (Herbst, 1784)

Pearce: 8 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2019

Stenus (Hypostenus) solutus Erichson, 1840

Pearce: no records
First record: 1997
Number of records: many
Latest record: 2021

Stenus (Hypostenus) tarsalis Ljungh, 1810

Pearce: 6 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2006

Stenus (Metastenus) bifoveolatus Gyllenhal, 1827

Pearce: 2 localities
First subsequent record: 1997
Number of subsequent records: many
Latest record: 2021

Stenus (Metastenus) binotatus Ljungh, 1804

Pearce: 1 locality (1920)
First subsequent record: 1997
Number of subsequent records: 2
Latest record: 2016

Stenus (Metastenus) brevipennis Thomson, C.G., 1851

Pearce: no records
First record: 2006
Number of records: many
Latest record: 2019

Stenus (Metastenus) butrintensis Smetana, 1959

Pearce: no records
First record: 2009
Number of records: 4
Latest record: 2011

Stenus (Metastenus) canescens Rosenhauer, 1856

Pearce: 1 locality (1899)
First subsequent record: 1992
Number of subsequent records: 5
Latest record: 2021

Stenus (Metastenus) flavipes Stephens, 1833

Pearce: 5 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Stenus (Metastenus) nitidiusculus Stephens, 1833

Pearce: 3 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2020

Stenus (Metastenus) niveus Fauvel, 1865

Pearce: no records
First record: 2018
Number of records: 1
Latest record: 2018

Stenus (Metastenus) pallitarsis Stephens, 1833

Pearce: 5 localities
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2020

Stenus (Metastenus) picipennis Erichson, 1840

Pearce: 1 locality (1899)
First subsequent record: 1992
Number of subsequent records: many
Latest record: 2021

Stenus (Metastenus) picipes Stephens, 1833

Pearce: 1 locality
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2020

Stenus (Metastenus) pubescens Stephens, 1833

Pearce: 1 locality
First subsequent record: 1989
Number of subsequent records: 4
Latest record: 2006

Stenus (Stenus) argus Gravenhorst, 1806

Pearce: no records
First record: 2008
Number of records: 1
Latest record: 2008

Stenus (Stenus) assequens Rey, 1844

Pearce: no records
First record: 1967
Number of records: 1
Latest record: 1967

Stenus (Stenus) ater Mannerheim, 1830

Pearce: 6 localities
No subsequent records

Stenus (Stenus) atratulus Erichson, 1839

Pearce: 1 locality (1890)
First subsequent record: 2010
Number of subsequent records: 1
Latest record: 2010

Stenus (Stenus) biguttatus (Linnaeus, 1758)

Pearce: no records
First record: 1989
Number of records: 8
Latest record: 2009

Stenus (Stenus) bimaculatus Gyllenhal, 1810

Pearce: 6 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Stenus (Stenus) boops Ljungh, 1810

Pearce: 4 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2020

Stenus (Stenus) canaliculatus Gyllenhal, 1827

Pearce: 2 localities
First subsequent record: 1966
Number of subsequent records: 6
Latest record: 2021

Stenus (Stenus) circularis Gravenhorst, 1802

Pearce: 2 localities (1899, 1909, 1923)
First subsequent record: 2007
Number of subsequent records: 3
Latest record: 2017

Stenus (Stenus) clavicornis (Scopoli, 1763)

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2020

Stenus (Stenus) comma LeConte, 1863

Pearce: 1 locality
First subsequent record: 1989
Number of subsequent records: 2
Latest record: 2005

Stenus (Stenus) guttula Müller, P.W.J., 1821

Pearce: 3 localities
First subsequent record: 1988
Number of subsequent records: many
Latest record: 2021

Stenus (Stenus) incrassatus Erichson, 1839

Pearce: 1 locality
First subsequent record: 2005
Number of subsequent records: 2
Latest record: 2015

Stenus (Stenus) junco (Paykull, 1789)

Pearce: 6 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Stenus (Stenus) lustrator Erichson, 1839

Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Stenus (Stenus) melanarius Stephens, 1833

Pearce: 1 locality
First subsequent record: 1998
Number of subsequent records: many
Latest record: 2014

Stenus (Stenus) melanopus (Marshall, 1802)

Pearce: 2 localities
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2021

Stenus (Stenus) nanus Stephens, 1833

Pearce: 1 locality
First subsequent record: 1998
Number of subsequent records: many
Latest record: 2021

Stenus (Stenus) nitens Stephens, 1833

Pearce: no records
First record: 1998
Number of records: 6
Latest record: 2021

Stenus (Stenus) providus Erichson, 1839

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Stenus (Stenus) pusillus Stephens, 1833

Pearce: 8 localities
First subsequent record: 1966
Number of subsequent records: many
Latest record: 2021

Stenus (Stenus) subdepressus Mulsant & Rey, 1861

Pearce: no records
First record: 2016
Number of records: 1
Latest record: 2016

Stenus (Tesus) brunnipes Stephens, 1833

Pearce: generally common
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2021

Stenus (Tesus) crassus Stephens, 1833

Pearce: 1 locality
First subsequent record: 1990
Number of subsequent records: 2
Latest record: 1999

Stenus (Tesus) formicetorum

Mannerheim, 1843
Pearce: no records
First record: 1999
Number of records: 3
Latest record: 2019

Stenus (Tesus) nigrifrons Gyllenhal, 1827

Pearce: 2 localities
First subsequent record: 1997
Number of subsequent records: 1
Latest record: 1997

Euaesthetus bipunctatus (Ljungh, 1804)

Pearce: no records
First record: 1980
Number of records: 4
Latest record: 2009

Euaesthetus laeviusculus Mannerheim, 1844

Pearce: 1 locality (1925)
First subsequent record: 2020
Number of subsequent records: 3
Latest record: 2021

Euaesthetus ruficapillus (Lacordaire, 1835)

Pearce: 2 localities
First subsequent record: 2006
Number of subsequent records: many
Latest record: 2021

Lathrobium brunnipes (Fabricius, 1792)

Pearce: 8 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Lathrobium elongatum (Linnaeus, 1767)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2015

Lathrobium foveolum Stephens, 1833

Pearce: 1 locality
First subsequent record: 2006
Number of subsequent records: many
Latest record: 2020

Lathrobium fulvipenne (Gravenhorst, 1806)

Pearce: 6 localities
First subsequent record: 1987
Number of subsequent records: many
Latest record: 2020

Lathrobium geminum Kraatz, 1857

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Lathrobium impressum Heer, 1841

Pearce: 2 localities
First subsequent record: 2006
Number of subsequent records: 5
Latest record: 2014

Lathrobium longulum Gravenhorst, 1802

Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: 7
Latest record: 2019

Lathrobium pallidum Nordmann, 1837

Pearce: no records
First record: 1965
Number of records: 1
Latest record: 1965

Lobrathium multipunctum (Gravenhorst, 1802)

Pearce: 7 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2021

Tetartopeus angustatus (Lacordaire, 1835)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Tetartopeus quadratus (Paykull, 1789)

Pearce: 1 locality (1896)
First subsequent record: 1962
Number of subsequent records: 7
Latest record: 2017

Tetartopeus rufonitidus (Reitter, 1909)

Pearce: no records
First record: 1994
Number of records: 2
Latest record: 2006

Tetartopeus terminatus (Gravenhorst, 1802)

Pearce: 6 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Achenium depressum (Gravenhorst, 1802)

Pearce: 5 localities
First subsequent record: 1993
Number of subsequent records: 4
Latest record: 2002

Achenium humile (Nicolai, 1822)

Pearce: 2 localities
First subsequent record: 1988
Number of subsequent records: 1
Latest record: 1988

Medon apicalis (Kraatz, 1857)

Pearce: 1 locality
First subsequent record: 2008
Number of subsequent records: 1
Latest record: 2008

Medon brunneus (Erichson, 1839)

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Medon fuscus (Mannerheim, 1830)

Pearce: 1 locality (1886)
First subsequent record: 1993
Number of subsequent records: 2
Latest record: 1994

Medon pocifer (Peyron, 1858)

Pearce: 1 locality
First subsequent record: 1998
Number of subsequent records: 3
Latest record: 2017

Medon ripicola (Kraatz, 1854)

Pearce: no records
First record: 2016
Number of records: 1
Latest record: 2016

Sunius melanocephalus (Fabricius, 1792)

Pearce: 2 localities
First subsequent record: 1979
Number of subsequent records: 8
Latest record: 2019

Sunius propinquus (Brisout de Barneville, C., 1867)

Pearce: 2 localities (1898, 1901)
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Lithocharis nigriceps Kraatz, 1859

Pearce: no records
First record: 1971
Number of records: many
Latest record: 2021

Lithocharis ochracea (Gravenhorst, 1802)

Pearce: 1 locality (1885)
First subsequent record: 2003
Number of subsequent records: 6
Latest record: 2006

Pseudomedon obscurellus (Erichson, 1840)

Pearce: no records
First record: 1997
Number of records: 3
Latest record: 2003

Pseudomedon obsoletus (Nordmann, 1837)

Pearce: 1 locality
No subsequent records

Scopaeus laevigatus (Gyllenhal, 1827)

Pearce: no records
First record: 1989
Number of records: 7
Latest record: 2006

Scopaeus minutus Erichson, 1840

Pearce: 2 localities
No subsequent records

Scopaeus sulcicollis (Stephens, 1833)

Pearce: 2 localities
First subsequent record: 1991
Number of subsequent records: 4
Latest record: 2015

Rugilus angustatus (Geoffroy in Fourcroy, 1785)

Pearce: 2 localities (1873, 1891)
First subsequent record: 1999
Number of subsequent records: 1
Latest record: 1999

Rugilus erichsonii (Fauvel, 1867)

Pearce: 3 localities
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2021

Rugilus geniculatus (Erichson, 1839)

Pearce: 3 localities
First subsequent record: 1979
Number of subsequent records: 3
Latest record: 2020

Rugilus orbiculatus (Paykull, 1789)

Pearce: 8 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2021

Rugilus rufipes Germar, 1836

Pearce: 4 localities
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2020

Rugilus similis (Erichson, 1839)

Pearce: 1 locality
First subsequent record: 2000
Number of subsequent records: 1
Latest record: 2000

Rugilus subtilis (Erichson, 1840)

Pearce: 1 locality
No subsequent records

Astenus immaculatus Stephens, 1833

Pearce: 1 locality (1901)
First subsequent record: 2013
Number of subsequent records: 7
Latest record: 2021

Astenus lyonessius (Joy, 1908)

Pearce: 5 localities
First subsequent record: 1962
Number of subsequent records: many
Latest record: 2021

Astenus procerus (Gravenhorst, 1806)

Pearce: 3 localities
No subsequent records

Astenus pulchellus (Heer, 1839)

Pearce: 6 localities
First subsequent record: 1999
Number of subsequent records: 3
Latest record: 2021

Ochtheophilum fracticorne (Paykull, 1800)

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2020

Ochtheophilum jacquelinii (Boieldieu, 1859)

Pearce: no records
First record: 1981
Number of records: 3
Latest record: 2005

Paederus (Eopaederus) caligatus

Erichson, 1840
Pearce: 3 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Paederus (Heteropaederus) fuscipes
Curtis, 1826

Pearce: 7 localities
First subsequent record: 1987
Number of subsequent records: many
Latest record: 2021

Paederus (Paederus) riparius (Linnaeus, 1758)

Pearce: fairly common and widely distributed
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Paederus (Poederomorphus) littoralis
Gravenhorst, 1802

Pearce: generally common and widely distributed
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Othius angustus Stephens, 1833

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: 5
Latest record: 2021

Othius laeviusculus Stephens, 1833

Pearce: 3 localities
First subsequent record: 1990
Number of subsequent records: many
Latest record: 2020

Othius punctulatus (Goeze, 1777)

Pearce: 4 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Othius subuliformis Stephens, 1833

Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Atrecus affinis (Paykull, 1789)

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Leptacinus batychrus (Gyllenhal, 1827)

Pearce: 3 localities
First subsequent record: 2003
Number of subsequent records: 1
Latest record: 2003

Leptacinus formicetorum Märkel, 1841

Pearce: no records
First record: 2014
Number of records: 2
Latest record: 2014

Leptacinus intermedius Donisthorpe, 1936

Pearce: no records
First record: 2003
Number of records: 2
Latest record: 2011

Leptacinus pusillus (Stephens, 1833)

Pearce: 1 locality
First subsequent record: 1990
Number of subsequent records: many
Latest record: 2020

Phacophallus parumpunctatus (Gyllenhal, 1827)

Pearce: 2 localities
First subsequent record: 1995
Number of subsequent records: 4
Latest record: 2021

Nudobius lentus (Gravenhorst, 1806)

Pearce: no records
First record: 2000
Number of records: 2
Latest record: 2004

Gauropterus fulgidus (Fabricius, 1787)

Pearce: no records
First record: 1990
Number of records: 4
Latest record: 2013

Gyrophypnus angustatus Stephens, 1833

Pearce: 6 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Gyrophypnus fracticornis (Müller, O.F., 1776)

Pearce: 12 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2021

Gyrophypnus punctulatus (Paykull, 1789)

Pearce: no records
First record: 1994
Number of records: 7
Latest record: 2009

Gyrophypnus wagneri (Scheerpeltz, 1926)

Pearce: no records
First record: 2003
Number of records: 1
Latest record: 2003

Xantholinus (Polydontophallus) elegans (Olivier, 1795)

Pearce: 4 localities
First subsequent record: 2009
Number of subsequent records: 1
Latest record: 2009

Xantholinus (Purrolinus) tricolor (Fabricius, 1787)

Pearce: no records
First record: 1939
Number of records: 1
Latest record: 1939

Xantholinus (Xantholinus) gallicus

Coiffait, 1956
Pearce: no records
First record: 1986
Number of records: 8
Latest record: 2015

Xantholinus (Xantholinus) linearis (Olivier, 1795)

Pearce: 10 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Xantholinus (Xantholinus) longiventris Heer, 1839

Pearce: 7 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Megalinus glabratus (Gravenhorst, 1802)

Pearce: 6 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2019

Heterothops binotatus (Gravenhorst, 1802)

Pearce: 4 localities
First subsequent record: 2005
Number of subsequent records: 3
Latest record: 2006

Heterothops niger Kraatz, 1868

Pearce: no records
First record: 1994
Number of records: 4
Latest record: 1997

Heterothops praeivius Erichson, 1839

Pearce: no records
First record: 1969
Number of records: 2
Latest record: 1975

Euryporus picipes (Paykull, 1800)

Pearce: 1 locality
No subsequent records

Quedius (Distichalius) cinctus (Paykull, 1790)

Pearce: 4 localities
First subsequent record: 1971
Number of subsequent records: 5
Latest record: 2019

Quedius (Microsaurus) aetolicus Kraatz, 1858

Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Quedius (Microsaurus) brevicornis
(Thomson, C.G., 1860)
Pearce: no records
First record: 1998
Number of records: 3
Latest record: 2020

Quedius (Microsaurus) brevis Erichson,
1840
Pearce: 1 locality
First subsequent record: 1936
Number of subsequent records: 1
Latest record: 1936

Quedius (Microsaurus) cruentus (Olivier,
1795)
Pearce: 4 localities
First subsequent record: 1941
Number of subsequent records: many
Latest record: 2020

Quedius (Microsaurus) fulgidus
(Fabricius, 1792)
Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: 1
Latest record: 1979

Quedius (Microsaurus) invreae Gridelli,
1924
Pearce: no records
First record: 2007
Number of records: 2
Latest record: 2015

Quedius (Microsaurus) longicornis
Kraatz, 1857
Pearce: 1 locality
First subsequent record: 1995
Number of subsequent records: 4
Latest record: 2000

Quedius (Microsaurus) maurus (Sahlberg,
C.R., 1830)
Pearce: no records
First record: 1970
Number of records: 3
Latest record: 2013

Quedius (Microsaurus) mesomelinus
(Marsham, 1802)
Pearce: 5 localities
First subsequent record: 1987
Number of subsequent records: 8
Latest record: 2020

Quedius (Microsaurus) microps
Gravenhorst, 1847
Pearce: no records
First record: 1997
Number of records: 2
Latest record: 2014

Quedius (Microsaurus) nigrocaeruleus
Fauvel, 1876
Pearce: 1 locality (1928)
First subsequent record: 1997
Number of subsequent records: 1
Latest record: 1997

Quedius (Microsaurus) scitus
(Gravenhorst, 1806)
Pearce: no records
First record: 1987
Number of records: many
Latest record: 2015

Quedius (Microsaurus) truncicola
Fairmaire & Laboulbène, 1856
Pearce: no records
First record: 2018
Number of records: 1
Latest record: 2018

Quedius (Microsaurus) xanthopus
Erichson, 1839
Pearce: 2 localities
First subsequent record: 1994
Number of subsequent records: 4
Latest record: 2018

Quedius (Quedius) curtipennis Bernhauer,
1908
Pearce: no records (but see *fuliginosus*)
First record: 1935
Number of records: many
Latest record: 2021

Quedius (Quedius) fuliginosus
(Gravenhorst, 1802)
Pearce: 4 localities (but probably includes
curtipennis)
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Quedius (Quedius) levicollis Brullé, 1832
Pearce: generally common
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2020

Quedius (Quedius) molochinus
(Gravenhorst, 1806)

Pearce: fairly common and widely distributed
First subsequent record: 2000
Number of subsequent records: many
Latest record: 2021

Quedius (Quedius) simplicifrons
Fairmaire, 1862

Pearce: 2 localities
First subsequent record: 1935
Number of subsequent records: 4
Latest record: 2019

Quedius (Raphirus) boops (Gravenhorst,
1802)

Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Quedius (Raphirus) fumatus (Stephens,
1833)

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Quedius (Raphirus) humeralis Stephens,
1832

Pearce: no records
First record: 1975
Number of records: 8
Latest record: 2021

Quedius (Raphirus) lateralis (Gravenhorst,
1802)

Pearce: 1 locality (1919)
First subsequent record: 1966
Number of subsequent records: many
Latest record: 2019

Quedius (Raphirus) maurorufus
(Gravenhorst, 1806)

Pearce: 5 localities
First subsequent record: 1989
Number of subsequent records: many
Latest record: 2020

Quedius (Raphirus) nemoralis Baudi di
Selve, 1848

Pearce: no records
First record: 1993
Number of records: 5
Latest record: 2009

Quedius (Raphirus) nigriceps Kraatz, 1857

Pearce: 5 localities
First subsequent record: 1965
Number of subsequent records: many
Latest record: 2021

Quedius (Raphirus) nitipennis (Stephens,
1833)

Pearce: 3 localities
First subsequent record: 1989
Number of subsequent records: 6
Latest record: 2014

Quedius (Raphirus) persimilis Mulsant &
Rey, 1876

Pearce: no records
First record: 1979
Number of records: many
Latest record: 2021

Quedius (Raphirus) picipes (Mannerheim,
1830)

Pearce: generally common and widely distributed
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2021

Quedius (Raphirus) plancus Erichson,
1840

Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Quedius (Raphirus) schatzmayri Gridelli,
1922

Pearce: no records
First record: 1968
Number of records: many
Latest record: 2021

Quedius (Raphirus) scintillans
(Gravenhorst, 1806)

Pearce: 1 locality (1899)
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2018

Quedius (Raphirus) semiaeneus (Stephens,
1833)

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Quedius (Raphirus) semiobscurus (Marshall, 1802)

Pearce: 6 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Quedius (Raphirus) umbrinus Erichson, 1839

Pearce: no records
First record: 1986
Number of records: many
Latest record: 2017

Quedius (Velleius) dilatatus (Fabricius, 1787)

Pearce: no records
First record: 2019
Number of records: 2
Latest record: 2021

Acylophorus glaberrimus (Herbst, 1784)

Pearce: no records
First record: 2016
Number of records: 1
Latest record: 2016

Platydracus fulvipes (Scopoli, 1763)

Pearce: no records
First record: 1987
Number of records: 3
Latest record: 2014

Platydracus latebricola (Gravenhorst, 1806)

Pearce: 3 localities
First subsequent record: 1944
Number of subsequent records: 5
Latest record: 2020

Platydracus stercorarius (Olivier, 1795)

Pearce: 14 localities
First subsequent record: 1997
Number of subsequent records: many
Latest record: 2019

Dinothenarus pubescens (De Geer, 1774)

Pearce: 1 locality
No subsequent records

Tasgius (Rayacheila) globulifer (Geoffroy in Fourcroy, 1785)

Pearce: 1 locality
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Tasgius (Rayacheila) melanarius (Heer, 1839)

Pearce: no records
First record: 1979
Number of records: many
Latest record: 2021

Tasgius (Rayacheila) morsitans (Rossi, 1790)

Pearce: 8 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Tasgius (Rayacheila) winkleri (Bernhauer, 1906)

Pearce: generally common and widely distributed
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2021

Tasgius (Tasgius) ater (Gravenhorst, 1802)

Pearce: widely distributed in south of county
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2019

Tasgius (Tasgius) pedator (Gravenhorst, 1802)

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: 5
Latest record: 1979

Ocypus (Angulephallus) brunnipes (Fabricius, 1781)

Pearce: 3 localities
First subsequent record: 1995
Number of subsequent records: many
Latest record: 2019

Ocypus (Matidus) nitens (Schrank, 1781)

Pearce: 1 locality
No subsequent records

Ocypus (Ocypus) olens (Müller, O.F., 1764)

Pearce: generally common and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Ocypus (Pseudocypus) aeneocephalus (De Geer, 1774)

Pearce: 10 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Ocypus (Pseudocypus) fortunatarum Wollaston, 1871

Pearce: 1 locality
First subsequent record: 1998
Number of subsequent records: 3
Latest record: 2019

Ocypus (Pseudocypus) fuscatus (Gravenhorst, 1802)

Pearce: 3 localities
First subsequent record: 1935
Number of subsequent records: 2
Latest record: 2005

Staphylinus dimidiaticornis Gemminger, 1851

Pearce: 10 localities (listed as *caesareus*)
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2021

Staphylinus erythropterus Linnaeus, 1758

Pearce: 2 localities
First subsequent record: 1982
Number of subsequent records: 6
Latest record: 2019

Creophilus maxillosus (Linnaeus, 1758)

Pearce: generally common and widely distributed
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2019

Ontholestes murinus (Linnaeus, 1758)

Pearce: 8 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Ontholestes tessellatus (Geoffroy in Fourcroy, 1785)

Pearce: 7 localities
First subsequent record: 1986
Number of subsequent records: 4
Latest record: 2019

Emus hirtus (Linnaeus, 1758)

Pearce: 3 localities
No subsequent records

Neobisnius lathroboides (Baudi di Selve, 1848)

Pearce: no records
First record: 2017
Number of records: 1
Latest record: 2017

Neobisnius procerulus (Gravenhorst, 1806)

Pearce: 2 localities
No subsequent records

Neobisnius villosulus (Stephens, 1833)

Pearce: 1 locality
First subsequent record: 1994
Number of subsequent records: 7
Latest record: 2017

Erichsonius cinerascens (Gravenhorst, 1802)

Pearce: 2 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2021

Erichsonius signaticornis (Mulsant & Rey, 1853)

Pearce: 2 localities
First subsequent record: 1989
Number of subsequent records: 5
Latest record: 2006

Bisnius cephalotes (Gravenhorst, 1802)

Pearce: 6 localities
First subsequent record: 1997
Number of subsequent records: 4
Latest record: 2020

Bisnius fimetarius (Gravenhorst, 1802)

Pearce: 5 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2021

Bisnius nigriventris (Thomson, C.G., 1867)

Pearce: 1 locality
First subsequent record: 1993
Number of subsequent records: 1
Latest record: 1993

Bisnius puella (Nordmann, 1837)

Pearce: no records
First record: 1968
Number of records: 1
Latest record: 1968

Bisnius sordidus (Gravenhorst, 1802)

Pearce: 5 localities
First subsequent record: 1996
Number of subsequent records: many
Latest record: 2020

Bisnius subuliformis (Gravenhorst, 1802)

Pearce: 1 locality (1871)
First subsequent record: 1987
Number of subsequent records: 4
Latest record: 2020

Philonthus (Onychophilonthus) marginatus (Müller, O.F., 1764)

Pearce: 5 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Philonthus (Philonthus) addendus Sharp, 1867

Pearce: 3 localities
First subsequent record: 1979
Number of subsequent records: 6
Latest record: 2021

Philonthus (Philonthus) albipes (Gravenhorst, 1802)

Pearce: 1 locality (1909)
First subsequent record: 1971
Number of subsequent records: 7
Latest record: 2018

Philonthus (Philonthus) atratus (Gravenhorst, 1802)

Pearce: no records
First record: 2020
Number of records: 1
Latest record: 2020

Philonthus (Philonthus) carbonarius (Gravenhorst, 1802)

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Philonthus (Philonthus) cognatus Stephens, 1832

Pearce: no records
First record: 1935
Number of records: many
Latest record: 2021

Philonthus (Philonthus) concinnus (Gravenhorst, 1802)

Pearce: no records
First record: 2019
Number of records: 3
Latest record: 2019

Philonthus (Philonthus) confinis Strand, A., 1941

Pearce: no records
First record: 2021
Number of records: 1
Latest record: 2021

Philonthus (Philonthus) cruentatus (Gmelin, 1790)

Pearce: 5 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2020

Philonthus (Philonthus) debilis (Gravenhorst, 1802)

Pearce: 2 localities
First subsequent record: 1995
Number of subsequent records: 6
Latest record: 1999

Philonthus (Philonthus) decorus (Gravenhorst, 1802)

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Philonthus (Philonthus) discoideus (Gravenhorst, 1802)

Pearce: 2 localities
First subsequent record: 1990
Number of subsequent records: 6
Latest record: 2006

Philonthus (Philonthus) ebeninus (Gravenhorst, 1802)

Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Philonthus (Philonthus) fumarius (Gravenhorst, 1806)

Pearce: 2 localities
First subsequent record: 2011
Number of subsequent records: 3
Latest record: 2021

Philonthus (Philonthus) intermedius
(Lacordaire, 1835)
Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Philonthus (Philonthus) jurgans
Tottenham, 1937
Pearce: no records
First record: 1993
Number of records: 8
Latest record: 2020

Philonthus (Philonthus) laminatus
(Creutzer, 1799)
Pearce: 6 localities
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2021

Philonthus (Philonthus) longicornis
Stephens, 1832
Pearce: 6 localities
First subsequent record: 1971
Number of subsequent records: 4
Latest record: 2021

Philonthus (Philonthus) mannerheimi
Fauvel, 1869
Pearce: no records
First record: 1968
Number of records: 1
Latest record: 1968

Philonthus (Philonthus) micans
(Gravenhorst, 1802)
Pearce: 4 localities
First subsequent record: 1997
Number of subsequent records: 4
Latest record: 2020

Philonthus (Philonthus) micantoides
Benick & Lohse, 1956
Pearce: no records
First record: 2006
Number of records: 2
Latest record: 2007

Philonthus (Philonthus) nigrita
(Gravenhorst, 1806)
Pearce: 3 localities
First subsequent record: 1987
Number of subsequent records: many
Latest record: 2018

Philonthus (Philonthus) parvicornis
(Gravenhorst, 1802)
Pearce: 2 localities
First subsequent record: 2015
Number of subsequent records: 1
Latest record: 2015

Philonthus (Philonthus) politus (Linnaeus, 1758)
Pearce: generally common
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Philonthus (Philonthus) quisquiliarius
(Gyllenhal, 1810)
Pearce: 4 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Philonthus (Philonthus) rectangulus
Sharp, 1874
Pearce: no records
First record: 1995
Number of records: 2
Latest record: 2006

Philonthus (Philonthus) rubripennis
Stephens, 1832
Pearce: no records
First record: 2017
Number of records: 4
Latest record: 2020

Philonthus (Philonthus) rufipes (Stephens, 1832)
Pearce: 1 locality (1899)
No subsequent records

Philonthus (Philonthus) sanguinolentus
(Gravenhorst, 1802)
Pearce: 5 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Philonthus (Philonthus) spinipes Sharp, 1874
Pearce: no records
First record: 1997
Number of records: 5
Latest record: 2006

Philonthus (Philonthus) splendens
(Fabricius, 1792)
Pearce: 5 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2015

Philonthus (Philonthus) succicola
(Thomson, C.G., 1860)
Pearce: no records
First record: 1975
Number of records: many
Latest record: 2021

Philonthus (Philonthus) tenuicornis
(Mulsant & Rey, 1853)
Pearce: 2 localities
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2021

Philonthus (Philonthus) umbratilis
(Gravenhorst, 1802)
Pearce: no records
First record: 1989
Number of records: 5
Latest record: 2019

Philonthus (Philonthus) varians (Paykull, 1789)
Pearce: 8 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Philonthus (Philonthus) ventralis
(Gravenhorst, 1802)
Pearce: 1 locality
First subsequent record: 2003
Number of subsequent records: 1
Latest record: 2003

Gabronthus thermarum (Aubé, 1850)
Pearce: no records
First record: 2003
Number of records: 6
Latest record: 2013

Gabrius appendiculatus Sharp, 1910
Pearce: no records
First record: 1999
Number of records: 7
Latest record: 2008

Gabrius breviventer (Sperk, 1835)
Pearce: 3 localities
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2021

Gabrius keysianus Sharp, 1910
Pearce: 1 locality
First subsequent record: 2001
Number of subsequent records: 7
Latest record: 2021

Gabrius nigrutilus (Gravenhorst, 1802)
Pearce: 8 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2011

Gabrius piliger Mulsant & Rey, 1876
Pearce: no records
First record: 1975
Number of records: 4
Latest record: 2015

Gabrius splendidulus (Gravenhorst, 1802)
Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Gabrius trossulus (Nordmann, 1837)
Pearce: 4 localities
First subsequent record: 1999
Number of subsequent records: 3
Latest record: 2017

Gabrius velox Sharp, 1910
Pearce: 1 locality
First subsequent record: 1996
Number of subsequent records: 4
Latest record: 2015

Cafius (Cafius) fucicola Curtis, 1830
Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: 2
Latest record: 2006

Cafius (Cafius) xantholoma (Gravenhorst, 1806)
Pearce: very common along coast
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Cafius (Platygonium) cicatricosus
(Erichson, 1840)

Pearce: 1 locality
No subsequent records

Remus sericeus Holme, 1837

Pearce: 8 localities
First subsequent record: 1966
Number of subsequent records: 5
Latest record: 2020

20. Family GEOTRUPIDAE Latreille, 1802

Odonteus armiger (Scopoli, 1772)

Pearce: 2 localities
No subsequent records

Typhaeus typhoeus (Linnaeus, 1758)

Pearce: very common on heaths
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Anoplotrupes stercorosus (Scriba, 1791)

Pearce: generally common and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Geotrupes mutator (Marsham, 1802)

Pearce: widely distributed in coastal part of county
First subsequent record: 1951
Number of subsequent records: 8
Latest record: 2019

Geotrupes spiniger (Marsham, 1802)

Pearce: generally common
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2020

Geotrupes stercorarius (Linnaeus, 1758)

Pearce: generally common and widely distributed
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2020

Trypocopris pyrenaicus (Charpentier, 1825)

Pearce: 4 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

21. Family TROGIDAE Macleay, 1819

Trox perlatus (Goeze, 1777)

Pearce: 1 locality
No subsequent records

Trox sabulosus (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 1940
Number of subsequent records: 1
Latest record: 1940

Trox scaber (Linnaeus, 1767)

Pearce: 3 localities
First subsequent record: 1950
Number of subsequent records: 3
Latest record: 2013

22. Family LUCANIDAE Latreille, 1804

Sinodendron cylindricum (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Dorcus parallelipipedus (Linnaeus, 1758)

Pearce: 7 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Lucanus cervus (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 1945
Number of subsequent records: many
Latest record: 2021

23. Family SCARABAEIDAE Latreille, 1802

Aegialia arenaria (Fabricius, 1787)

Pearce: 3 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2020

Acrossus depressus (Kugelann, 1792)
[2012: *Aphodius depressus*]

Pearce: 4 localities (but at least one is misidentified *luridus* – DJ Mann)
First subsequent record: 2008
Number of subsequent records: 1
Latest record: 2008

Acrossus luridus (Fabricius, 1775) [2012:
Aphodius luridus]
Pearce: 9 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2015

Acrossus rufipes (Linnaeus, 1758) [2012:
Aphodius rufipes]
Pearce: generally common
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Agrilinus ater (De Geer, 1774) [2012:
Aphodius ater]
Pearce: generally common
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2019

Agrilinus constans (Duftschmid, 1805)
[2012: *Aphodius constans*]
Pearce: 2 localities
First subsequent record: 1999
Number of subsequent records: many
Latest record: 2020

Aphodius fimetarius (Linnaeus, 1758)
Pearce: generally common and widely distributed
(but many of these may be *pedellus*)
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2019

Aphodius foetidus (Herbst, 1783)
Pearce: 4 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Aphodius pedellus (De Geer, 1774)
Pearce: no records (but see *fimetarius*)
First record: 1935
Number of records: many
Latest record: 2021

Bodiloides ictericus (Laicharting, 1781)
[2012: *Aphodius ictericus*]
Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Bodilopsis rufa (Moll, 1782) [2012:
Aphodius rufus]
Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Bodilopsis sordida (Fabricius, 1775)
[2012: *Aphodius sordidus*]
Pearce: 1 locality
No subsequent records

Calamosternus granarius (Linnaeus,
1767) [2012: *Aphodius granarius*]
Pearce: generally common
First subsequent record: 1950
Number of subsequent records: many
Latest record: 2019

Chilothorax conspurcatus (Linnaeus,
1758) [2012: *Aphodius conspurcatus*]
Pearce: no records
First record: 2007
Number of records: 6
Latest record: 2017

Chilothorax distinctus (Müller, O.F.,
1776) [2012: *Aphodius distinctus*]
Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: 3
Latest record: 2015

Chilothorax paykulli (Bedel, 1908) [2012:
Aphodius paykulli]
Pearce: 1 locality
No subsequent records

Colobopterus erraticus (Linnaeus, 1758)
[2012: *Aphodius erraticus*]
Pearce: 5 localities
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2020

Esymus merdarius (Fabricius, 1775)
[2012: *Aphodius merdarius*]
Pearce: 5 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2015

Esymus pusillus (Herbst, 1789) [2012:
Aphodius pusillus]
Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2019

Euheptaulacus sus (Herbst, 1783)
Pearce: 1 locality
No subsequent records

Euheptaulacus villosus (Gyllenhal in
Schönherr, 1806)
Pearce: no records
First record: 1939
Number of records: 2
Latest record: 2008

Euorodalus coenosus (Panzer, 1798)
[2012: *Aphodius coenosus*]
Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Eupleurus subterraneus (Linnaeus, 1758)
[2012: *Aphodius subterraneus*]
Pearce: 1 locality
No subsequent records

Heptaulacus testudinarius (Fabricius,
1775)
Pearce: 2 localities
First subsequent record: 1930
Number of subsequent records: 2
Latest record: 1931

Limarus zenkeri (Germar, 1813) [2012:
Aphodius zenkeri]
Pearce: no records
First record: 2007
Number of records: 9
Latest record: 2017

Liothorax niger (Illiger, 1798) [2012:
Aphodius ater]
Pearce: no records
First record: 1913
Number of records: 2
Latest record: 1915

Liothorax plagiatus (Linnaeus, 1767)
[2012: *Aphodius plagiatus*]
Pearce: 2 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2015

Melinopterus prodromus (Brahm, 1790)
[2012: *Aphodius prodromus*]
Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Melinopterus sphacelatus (Panzer, 1798)
[2012: *Aphodius sphacelatus*]
Pearce: generally common
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2020

Nimbus contaminatus (Herbst, 1783)
[2012: *Aphodius contaminatus*]
Pearce: 4 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Nimbus obliteratedus (Panzer, 1823) [2012:
Aphodius obliteratedus]
Pearce: 2 localities (but misidentifications – DJ
Mann)
First subsequent record: 2014
Number of subsequent records: 8
Latest record: 2019

Otophorus haemorrhoidalis (Linnaeus,
1758) [2012: *Aphodius haemorrhoidalis*]
Pearce: 6 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2018

Oxyomus sylvestris (Scopoli, 1763)
Pearce: “Dorset” (A. Ford)
First subsequent record: 1945
Number of subsequent records: 3
Latest record: 2007

Plagiogonus arenarius (Olivier, 1789)
[2012: *Aphodius arenarius*]
Pearce: 1 locality (1921)
No subsequent records

Planolinoides borealis (Gyllenhal, 1827)
[2012: *Aphodius borealis*]
Pearce: no records
First record: 1926
Number of records: many
Latest record: 2017

Rhodaphodius foetens (Fabricius, 1787)
[2012: *Aphodius foetens*]
Pearce: 8 localities (but 4 are from misidentifications – DJ Mann)
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2017

Sigorus porcus (Fabricius, 1792) [2012: *Aphodius porcus*]
Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: 1
Latest record: 1975

Teuchestes fossor (Linnaeus, 1758) [2012: *Aphodius fossor*]
Pearce: generally common and widely distributed
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2019

Volinus sticticus (Panzer, 1798) [2012: *Aphodius sticticus*]
Pearce: 3 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2019

Copris lunaris (Linnaeus, 1758)
Pearce: 1 locality
First subsequent record: 1954
Number of subsequent records: 1
Latest record: 1954

Onthophagus coenobita (Herbst, 1783)
Pearce: 5 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2020

Onthophagus joannae Goljan, 1953
Pearce: 7 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2019

Onthophagus medius (Kugelann, 1792)
Pearce: 4 localities
First subsequent record: 1950
Number of subsequent records: many
Latest record: 2019

Onthophagus nuchicornis (Linnaeus, 1758)
Pearce: 7 localities (but all vouchers checked are *similis* – DJ Mann)
First record: 1864
Number of records: 5
Latest record: 1954

Onthophagus similis (Scriba, 1790)
Pearce: no records (but see *nuchicornis*)
First record: 1936
Number of records: many
Latest record: 2020

Hoplia philanthus (Fuessly, 1775)
Pearce: 8 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2021

Melolontha melolontha (Linnaeus, 1758)
Pearce: generally common and widely distributed
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Amphimallon fallenii (Gyllenhal, 1817)
Pearce: no records
First record: 1986
Number of records: 7
Latest record: 1996

Amphimallon solstitiale (Linnaeus, 1758)
Pearce: fairly common and widely distributed
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2021

Omaloelia ruricola (Fabricius, 1775)
Pearce: 5 localities
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2015

Serica brunnea (Linnaeus, 1758)
Pearce: 11 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Anomala dubia (Scopoli, 1763)

Pearce: 4 localities

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2021

Phyllopertha horticola (Linnaeus, 1758)

Pearce: generally common

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2021

Cetonia aurata (Linnaeus, 1758)

Pearce: 7 localities

First subsequent record: 1943

Number of subsequent records: many

Latest record: 2021

24. Family EUCINETIDAE Lacordaire, 1857

Nycteus meridionalis Laporte, 1838 [2012:

Eucinetus meridionalis]

Pearce: no records

First record: 1991

Number of records: 4

Latest record: 2004

25. Family CLAMBIDAE Fischer von Waldheim, 1821

Calyptomerus dubius (Marshall, 1802)

Pearce: 2 localities

First subsequent record: 1975

Number of subsequent records: 2

Latest record: 2003

Clambus armadillo (De Geer, 1774)

Pearce: no records

First record: 2000

Number of records: 1

Latest record: 2000

Clambus pubescens Redtenbacher, 1847

Pearce: 1 locality

First subsequent record: 1971

Number of subsequent records: 5

Latest record: 2018

Clambus punctulum (Beck, 1817)

Pearce: no records

First record: 2005

Number of records: 1

Latest record: 2005

26. Family SCIRTIDAE Fleming, 1821

Elodes elongatus Tournier, 1868

Pearce: no records

First subsequent record: 2002

Number of subsequent records: 4

Latest record: 2011

Elodes minutus (Linnaeus, 1767)

Pearce: 5 localities

First subsequent record: 1964

Number of subsequent records: 6

Latest record: 2017

Elodes pseudominutus Klausnitzer, 1971

Pearce: no records

First record: 2005

Number of records: many

Latest record: 2016

Odeles marginata (Fabricius, 1798)

Pearce: 3 localities

First subsequent record: 1987

Number of subsequent records: many

Latest record: 2017

Microcara testacea (Linnaeus, 1767)

Pearce: generally common

First subsequent record: 1967

Number of subsequent records: many

Latest record: 2020

Contacyphon coarctatus (Paykull, 1799)

Pearce: 4 localities

First subsequent record: 1959

Number of subsequent records: many

Latest record: 2021

Contacyphon hilaris (Nyholm, 1944)

Pearce: no records

First record: 1935

Number of records: many

Latest record: 2020

Contacyphon laevipennis (Tournier, 1868)

Pearce: no records

First record: 1992

Number of records: many

Latest record: 2019

Contacyphon ochraceus (Stephens, 1830)

Pearce: 2 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2019

Contacyphon padi (Linnaeus, 1758)

Pearce: 9 localities

First subsequent record: 1966

Number of subsequent records: many

Latest record: 2021

Contacyphon palustris (Thomson, C.G., 1855)

Pearce: 5 localities

First subsequent record: 1999

Number of subsequent records: 8

Latest record: 2011

Contacyphon pubescens (Fabricius, 1792)

Pearce: no records

First record: 2005

Number of records: 1

Latest record: 2005

Contacyphon variabilis (Thunberg, 1787)

Pearce: generally common

First subsequent record: 1990

Number of subsequent records: many

Latest record: 2021

Prionocyphon serricornis (Müller, P.W.J., 1821)

Pearce: 1 locality (1841)

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2019

Hydrocyphon deflexicollis (Müller, P.W.J., 1821)

Pearce: 1 locality

No subsequent records

Scirtes hemisphaericus (Linnaeus, 1758)

Pearce: 4 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2020

27. Family DASCILLIDAE Guérin-Méneville, 1843 (1834)

Dascillus cervinus (Linnaeus, 1758)

Pearce: 3 localities

First subsequent record: 1943

Number of subsequent records: many

Latest record: 2019

28. Family BUPRESTIDAE Leach, 1815

Melanophila acuminata (De Geer, 1774)

Pearce: 1 locality

No subsequent records

Agrilus angustulus (Illiger, 1803)

Pearce: 2 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2019

Agrilus biguttatus (Fabricius, 1777)

Pearce: no records

First record: 1998

Number of records: 1

Latest record: 1998

Agrilus laticornis (Illiger, 1803)

Pearce: 2 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2019

Agrilus sinuatus (Olivier, 1790)

Pearce: no records

First record: 1951

Number of records: many

Latest record: 2021

Agrilus viridis (Linnaeus, 1758)

Pearce: no records

First record: 1988

Number of records: 3

Latest record: 2021

Aphanisticus emarginatus (Olivier, 1790)

Pearce: no records

First record: 1953

Number of records: 3

Latest record: 2013

Aphanisticus pusillus (Olivier, 1790)

Pearce: 3 localities

First subsequent record: 1940

Number of subsequent records: many

Latest record: 2021

Trachys compressus Abeille de Perrin, 1891 [2012: *Trachys subglaber*]

Pearce: no records (but the record of *trogodytes*, and subsequent records of *trogodytes* could refer to this species)

First record: 1938

Number of records: many

Latest record: 2020

Trachys minutus (Linnaeus, 1758)

Pearce: 1 locality (1834)

No subsequent records

Trachys scrobiculatus Kiesenwetter, 1857

Pearce: 2 localities

First subsequent record: 1952

Number of subsequent records: many

Latest record: 2021

29. Family BYRRHIDAE Latreille, 1804

Simplocaria semistriata (Fabricius, 1794)

Pearce: 6 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2019

Byrrhus fasciatus (Forster, 1771)

Pearce: 2 localities

First subsequent record: 1937

Number of subsequent records: many

Latest record: 2019

Byrrhus pilula (Linnaeus, 1758)

Pearce: generally common and widely distributed

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2020

Byrrhus pustulatus (Forster, 1771)

Pearce: 3 localities

First subsequent record: 1940

Number of subsequent records: many

Latest record: 2018

Cytilus sericus (Forster, 1771)

Pearce: common

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2019

Porcinolus murinus (Fabricius, 1794)

Pearce: 2 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2019

Chaetophora spinosa (Rossi, 1794)

Pearce: 3 localities

First subsequent record: 1991

Number of subsequent records: 8

Latest record: 2017

Curimopsis maritima (Marshall, 1802)

Pearce: 1 locality

First subsequent record: 1989

Number of subsequent records: 2

Latest record: 1998

Curimopsis setigera (Illiger, 1798)

Pearce: 1 locality

First subsequent record: 1982

Number of subsequent records: 5

Latest record: 2016

30. Family ELMIDAE Curtis, 1830

Elmis aenea (Müller, P.W.J., 1806)

Pearce: 6 localities

First subsequent record: 1940

Number of subsequent records: many

Latest record: 2021

Esolus parallelepipedus (Müller, P.W.J., 1806)

Pearce: no records

First record: 1987

Number of records: 7

Latest record: 2011

Limnius volckmari (Panzer, 1793)

Pearce: 3 localities

First subsequent record: 1965

Number of subsequent records: many

Latest record: 2021

Oulimnius major (Rey, 1889)

Pearce: no records

First record: 1982

Number of records: 3

Latest record: 2006

Oulimnius troglodytes (Gyllenhal, 1827)

Pearce: no records

First record: 1963

Number of records: 9

Latest record: 2006

Oulimnius tuberculatus (Müller, P.W.J., 1806)

Pearce: 2 localities

First subsequent record: 1974

Number of subsequent records: many

Latest record: 2020

Riolus nitens (Müller, P.W.J., 1817)

[2012: *Normandia nitens*]

Pearce: 1 locality

No subsequent records

Riolus subviolaceus (Müller, P.W.J., 1817)

Pearce: 1 locality

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2020

Stenelmis canaliculata (Gyllenhal, 1808)

Pearce: no records

First record: 1991

Number of records: 4

Latest record: 2013

Macronychus quadrituberculatus Müller,
P.W.J., 1806

Pearce: 1 locality (1923)

First subsequent record: 1975

Number of subsequent records: 3

Latest record: 2010

31. Family DRYOPIDAE Bilberg, 1820 (1817)

Dryops auriculatus (Geoffroy in Fourcroy,
1785)

Pearce: 4 localities

First subsequent record: 2007

Number of subsequent records: 1

Latest record: 2007

Dryops ernesti Gozis, 1886

Pearce: 2 localities

First subsequent record: 1936

Number of subsequent records: many

Latest record: 2021

Dryops luridus (Erichson, 1847)

Pearce: 1 locality

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2019

Dryops similaris Bollow, 1936

Pearce: no records

First record: 2013

Number of records: 3

Latest record: 2020

Dryops striatellus (Fairmaire & Brisout de
Barnville, C., 1859)

Pearce: 5 localities

First subsequent record: 2000

Number of subsequent records: many

Latest record: 2020

32. Family LIMNICHIDAE Erichson, 1846

Limnichus pygmaeus (Sturm, 1807)

Pearce: 2 localities (1886, 1904)

First subsequent record: 1966

Number of subsequent records: many

Latest record: 2017

33. Family HETEROCERIDAE MacLeay, 1825

Augyles maritimus (Guérin-Méneville,
1844)

Pearce: 1 locality

No subsequent records

Heterocerus fenestratus (Thunberg, 1784)

Pearce: 2 localities

First subsequent record: 1993

Number of subsequent records: many

Latest record: 2020

Heterocerus flexuosus Stephens, 1828

Pearce: 2 localities

First subsequent record: 1939

Number of subsequent records: 7

Latest record: 2018

Heterocerus fossor Kiesenwetter, 1843

Pearce: no records

First record: 1989

Number of records: 3

Latest record: 2011

Heterocerus fuscus Kiesenwetter, 1843

Pearce: no records

First record: 1980

Number of records: 2

Latest record: 1995

Heterocerus marginatus (Fabricius, 1787)

Pearce: 3 localities

First subsequent record: 1989

Number of subsequent records: many

Latest record: 2017

Heterocerus obsoletus Curtis, 1828

Pearce: 1 locality (1896)

No subsequent records

34. Family PSEPHENIDAE Lacordaire, 1854

Eubria palustris (Germar, 1818)

Pearce: 2 localities

First subsequent record: 1989

Number of subsequent records: many

Latest record: 2017

36. Family EUCNEMIDAE Eschscholtz, 1829

Microrhagus pygmaeus (Fabricius, 1792)

Pearce: 2 localities (1839, 1843)

First subsequent record: 1986

Number of subsequent records: many

Latest record: 2019

Epiphanis cornutus Eschscholtz, 1829

Pearce: no records

First record: 2007

Number of records: 8

Latest record: 2019

Hylis cariniceps (Reitter, 1902)

Pearce: no records

First record: 1976

Number of records: many

Latest record: 2021

Hylis olexai (Palm, 1955)

Pearce: no records

First record: 2016

Number of records: 7

Latest record: 2019

Melasis buprestoides (Linnaeus, 1760)

Pearce: no records

First record: 1975

Number of records: 4

Latest record: 2008

Eucnemis capucina Ahrens, 1812

Pearce: no records

First record: 2018

Number of records: 2

Latest record: 2019

37. Family THROSCIDAE Laporte, 1840

Trixagus carinifrons (de Bonvouloir, 1859)

Pearce: 1 locality

First subsequent record: 1990

Number of subsequent records: many

Latest record: 2021

Trixagus dermestoides (Linnaeus, 1767)

Pearce: 3 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2021

Trixagus gracilis Wollaston, 1854

Pearce: no records

First record: 2019

Number of records: 1

Latest record: 2019

Trixagus obtusus (Curtis, 1827)

Pearce: 1 locality (1886)

First subsequent record: 1993

Number of subsequent records: many

Latest record: 2021

38. Family ELATERIDAE Leach, 1815

Agrypnus murinus (Linnaeus, 1758)

Pearce: generally common and widely distributed

First subsequent record: 1933

Number of subsequent records: many

Latest record: 2021

Agriotes acuminatus (Stephens, 1830)

Pearce: 3 localities

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2019

Agriotes lineatus (Linnaeus, 1767)

Pearce: 3 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Agriotes obscurus (Linnaeus, 1758)

Pearce: generally common

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2021

Agriotes pallidulus (Illiger, 1807)

Pearce: generally common
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Agriotes sordidus (Illiger, 1807)

Pearce: 3 localities
First subsequent record: 1991
Number of subsequent records: 8
Latest record: 2011

Agriotes sputator (Linnaeus, 1758)

Pearce: 8 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Dalopius marginatus (Linnaeus, 1758)

Pearce: 9 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2020

Ampedus balteatus (Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2019

Ampedus cinnabarinus (Eschscholtz, 1829)

Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Ampedus elongatulus (Fabricius, 1787)

Pearce: 1 locality
First subsequent record: 1952
Number of subsequent records: many
Latest record: 2021

Ampedus sanguinolentus (Schränk, 1776)

Pearce: 3 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2015

Sericus brunneus (Linnaeus, 1758)

Pearce: 3 localities
First subsequent record: 1998
Number of subsequent records: many
Latest record: 2019

Procræus tibialis (Lacordaire, 1835)

Pearce: no records
First record: 1993
Number of records: 1
Latest record: 1993

Adrastus pallens (Fabricius, 1792)

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2019

Melanotus castanipes (Paykull, 1800)

Pearce: no records (but see *villosus*)
First record: 1934
Number of records: many
Latest record: 2021

Melanotus villosus (Geoffroy in Fourcroy, 1785)

Pearce: common in woods (but could include *castanipes*)
First subsequent record: 1945
Number of subsequent records: many
Latest record: 2019

Hypnoidus riparius (Fabricius, 1792)

Pearce: 3 localities
First subsequent record: 1988
Number of subsequent records: many
Latest record: 2017

Athous (Athous) haemorrhoidalis (Fabricius, 1801)

Pearce: generally common
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Athous (Athous) vittatus (Fabricius, 1792)

Pearce: generally common
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Athous (Orthathous) bicolor (Goeze, 1777)

Pearce: 11 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2020

Athous (Orthathous) campyloides

Newman, 1833

Pearce: no records

First record: 1992

Number of records: 4

Latest record: 2019

Cidnopus aeruginosus (Olivier, 1790)

Pearce: 1 locality

First subsequent record: 2008

Number of subsequent records: 1

Latest record: 2008

Limonium poneli Leseigneur & Mertlik,
2007 [2012: *Kibunea minuta*]

Pearce: 4 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2020

Denticollis linearis (Linnaeus, 1758)

Pearce: 5 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2020

Hemicrepidius hirtus (Herbst, 1784)

Pearce: 5 localities

First subsequent record: 1936

Number of subsequent records: many

Latest record: 2019

Stenagostus rhombeus (Olivier, 1790)

Pearce: no records

First record: 1991

Number of records: 8

Latest record: 2013

Actenicerus sjaelandicus (Müller, O.F.,
1764)

Pearce: 5 localities

First subsequent record: 1933

Number of subsequent records: many

Latest record: 2021

Aplotarsus incanus (Gyllenhal, 1827)

Pearce: 4 localities

First subsequent record: 1992

Number of subsequent records: many

Latest record: 2013

Calambus bipustulatus (Linnaeus, 1767)

Pearce: 3 localities

First subsequent record: 1939

Number of subsequent records: 7

Latest record: 2018

Ctenicera cuprea (Fabricius, 1775)

Pearce: 1 locality

No subsequent records

Paraphotistus nigricornis (Panzer, 1799)

Pearce: 2 localities

First subsequent record: 2006

Number of subsequent records: 4

Latest record: 2018

Prosternon tessellatum (Linnaeus, 1758)

Pearce: 5 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2020

Selatosomus aeneus (Linnaeus, 1758)

Pearce: 1 locality

First subsequent record: 1982

Number of subsequent records: 2

Latest record: 2000

Oedostethus quadripustulatus (Fabricius,
1792)

Pearce: 1 locality

No subsequent records

Cardiophorus asellus Erichson, 1840

Pearce: 5 localities

First subsequent record: 1933

Number of subsequent records: many

Latest record: 2018

39. Family DRILIDAE Blanchard, 1845

Drilus flavescens (Geoffroy in Fourcroy,
1785)

Pearce: 3 localities

First subsequent record: 1940

Number of subsequent records: 6

Latest record: 2015

40. Family LYCIDAE Laporte, 1836

Platycis minutus (Fabricius, 1787)

Pearce: no records

First record: 1979

Number of records: many

Latest record: 2018

41. Family LAMPYRIDAE Rafinesque, 1815

Lampyris noctiluca (Linnaeus, 1758)

Pearce: generally common
First subsequent record: 1952
Number of subsequent records: many
Latest record: 2021

42. Family CANTHARIDAE Imhoff, 1856 (1815)

Podabrus alpinus (Paykull, 1798)

Pearce: 3 localities
First subsequent record: 1990
Number of subsequent records: 7
Latest record: 2021

Cantharis (Cantharis) cryptica Ashe, 1947

Pearce: *bicolor*, described as common, could refer to this species
First record: 1935
Number of records: many
Latest record: 2021

Cantharis (Cantharis) decipiens Baudi di Selve, 1872

Pearce: 3 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2020

Cantharis (Cantharis) figurata Mannerheim, 1843

Pearce: no records
First record: 1987
Number of records: 4
Latest record: 1987

Cantharis (Cantharis) flavilabris Fallén, 1807 [2012: *Cantharis nigra*]

Pearce: generally common and widely distributed
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2020

Cantharis (Cantharis) fusca Linnaeus, 1758

Pearce: 3 localities
First subsequent record: 1977
Number of subsequent records: many
Latest record: 2018

Cantharis (Cantharis) livida Linnaeus, 1758

Pearce: 5 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2020

Cantharis (Cantharis) nigra (De Geer, 1774) [2012: *Cantharis thoracica*]

Pearce: 5 localities
First subsequent record: 1951
Number of subsequent records: many
Latest record: 2019

Cantharis (Cantharis) nigricans (Müller, O.F., 1776)

Pearce: generally common
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Cantharis (Cantharis) pallida Goeze, 1777

Pearce: *bicolor*, described as common, could refer to this species
First record: 1939
Number of records: many
Latest record: 2019

Cantharis (Cantharis) pellucida Fabricius, 1792

Pearce: generally common
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2019

Cantharis (Cantharis) rufa Linnaeus, 1758

Pearce: 5 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2019

Cantharis (Cantharis) rustica Fallén, 1807

Pearce: generally common
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Cantharis (Cyrptomoptila) lateralis Linnaeus, 1758

Pearce: 2 localities
First subsequent record: 1967
Number of subsequent records: many
Latest record: 2021

Rhagonycha fulva (Scopoli, 1763)

Pearce: abundant
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Rhagonycha lignosa (Müller, O.F., 1764)

Pearce: generally common
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Rhagonycha lutea (Müller, O.F., 1764)

Pearce: 1 locality
No subsequent records

Rhagonycha nigriventris (Thomson, C.G., 1864) [2012: *Rhagonycha limbata*]

Pearce: generally common
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2021

Rhagonycha testacea (Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2019

Rhagonycha translucida (Krynicky, 1832)

Pearce: 1 locality
No subsequent records

Silis ruficollis (Fabricius, 1775)

Pearce: "Dorset" (A. Ford)
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2020

Malthinus balteatus Suffrian, 1851

Pearce: 2 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Malthinus flaveolus (Herbst, 1786)

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2020

Malthinus frontalis (Marsham, 1802)

Pearce: 1 locality
First subsequent record: 1954
Number of subsequent records: 4
Latest record: 2010

Malthinus seriepunctatus Kiesenwetter, 1852

Pearce: generally common
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2020

Malthodes dispar (Germar, 1824)

Pearce: 1 locality
First subsequent record: 1984
Number of subsequent records: many
Latest record: 2015

Malthodes fibulatus Kiesenwetter, 1852

Pearce: no records
First record: 1982
Number of records: 2
Latest record: 1995

Malthodes flavoguttatus Kiesenwetter, 1852

Pearce: 3 localities
First subsequent record: 2008
Number of subsequent records: 2
Latest record: 2008

Malthodes fuscus (Waltl, 1838)

Pearce: 2 localities
First subsequent record: 1989
Number of subsequent records: 5
Latest record: 2019

Malthodes guttifer Kiesenwetter, 1852

Pearce: 3 localities
First subsequent record: 1987
Number of subsequent records: 8
Latest record: 2008

Malthodes marginatus (Latreille, 1806)

Pearce: 6 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2020

Malthodes minimus (Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2020

Malthodes mysticus Kiesenwetter, 1852

Pearce: 1 locality
First subsequent record: 2010
Number of subsequent records: 2
Latest record: 2021

Malthodes pumilus (Brébisson, 1835)

Pearce: 2 localities

First subsequent record: 1992

Number of subsequent records: many

Latest record: 2011

44. Family DERMESTIDAE Latreille, 1804

Dermestes (Dermestes) lardarius

Linnaeus, 1758

Pearce: 4 localities

No subsequent records

Dermestes (Dermestinus) maculatus De Geer, 1774

Pearce: no records

First record: 1997

Number of records: 1

Latest record: 1997

Dermestes (Dermestinus) murinus

Linnaeus, 1758

Pearce: generally common and widely distributed

First subsequent record: 2011

Number of subsequent records: 3

Latest record: 2021

Dermestes (Dermestinus) undulatus

Brahm, 1790

Pearce: 4 localities

First subsequent record: 1935

Number of subsequent records: 6

Latest record: 1995

Attagenus pellio (Linnaeus, 1758)

Pearce: 7 localities

First subsequent record: 1982

Number of subsequent records: 2

Latest record: 1995

Anthrenus (Florilinus) museorum

(Linnaeus, 1760)

Pearce: 3 localities

First subsequent record: 1941

Number of subsequent records: 1

Latest record: 1941

Anthrenus (Helocerus) fuscus Olivier, 1790

Pearce: 6 localities

First subsequent record: 1993

Number of subsequent records: 9

Latest record: 2019

Anthrenus (Nathrenus) verbasci

(Linnaeus, 1767)

Pearce: 1 locality

First subsequent record: 1994

Number of subsequent records: many

Latest record: 2019

Ctesias serra (Fabricius, 1792)

Pearce: no records

First record: 1988

Number of records: 5

Latest record: 2004

45. Family BOSTRICHIDAE Latreille, 1802

Bostrichus capucinus (Linnaeus, 1758)

Pearce: 1 locality (1854)

No subsequent records

Rhyzopertha dominica (Fabricius, 1792)

Pearce: 1 locality

No subsequent records

Lyctus linearis (Goeze, 1777)

Pearce: 1 locality

No subsequent records

46. Family PTINIDAE Latreille, 1802

Ptinomorphus imperialis (Linnaeus, 1767)

[2012: *Hedobia imperialis*]

Pearce: 2 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2018

Niptus holoeucus (Faldermann, 1835)

Pearce: 4 localities

No subsequent records

Epauloecus unicolor (Piller & Mitterpacher, 1783) [2012: *Tipnus unicolor*]

Pearce: 2 localities

First subsequent record: 1937

Number of subsequent records: 1

Latest record: 1937

Ptinus (Bruchoptinus) palliatus Perris, 1847

Pearce: 3 localities

No subsequent records

Ptinus (Gynopterus) sexpunctatus Panzer, 1789

Pearce: 2 localities

First subsequent record: 1940

Number of subsequent records: many

Latest record: 2007

Ptinus (Pseudoptinus) lichenum (Marsham, 1802)

Pearce: 1 locality

First subsequent record: 1979

Number of subsequent records: 1

Latest record: 1979

Ptinus (Ptinus) fur (Linnaeus, 1758)

Pearce: 4 localities

First subsequent record: 1940

Number of subsequent records: 3

Latest record: 2004

Ptinus (Ptinus) subpilosus Sturm, 1837

Pearce: no records

First record: 1996

Number of records: 5

Latest record: 2009

Ptinus (Tectoptynus) tectus Boieldieu, 1856

Pearce: 1 locality

No subsequent records

Grynobius planus (Fabricius, 1787)

Pearce: 3 localities

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2020

Dryophilus pusillus (Gyllenhal, 1808)

Pearce: 1 locality (1895)

First subsequent record: 1999

Number of subsequent records: 4

Latest record: 2011

Ochina ptinoides (Marsham, 1802)

Pearce: 1 locality

First subsequent record: 2000

Number of subsequent records: many

Latest record: 2021

Xestobium rufovillosum (De Geer, 1774)

Pearce: 3 localities

First subsequent record: 1953

Number of subsequent records: 2

Latest record: 2007

Ernobius mollis (Linnaeus, 1758)

Pearce: 4 localities

First subsequent record: 2001

Number of subsequent records: many

Latest record: 2019

Hemicoelus fulvicornis (Sturm, 1837)

[2012: *Anobium fulvicorne*]

Pearce: 2 localities

First subsequent record: 2000

Number of subsequent records: many

Latest record: 2021

Anobium inexpectatum Lohse, 1954

Pearce: no records

First record: 1993

Number of records: 9

Latest record: 2021

Anobium punctatum (De Geer, 1774)

Pearce: generally common and widely distributed

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Ptilinus pectinicornis (Linnaeus, 1758)

Pearce: 2 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2021

Dorcatoma (Dorcatoma) dresdensis

Herbst, 1791

Pearce: no records

First record: 2010

Number of records: 5

Latest record: 2021

Dorcatoma (Pilosodorcatoma)

chrysomelina Sturm, 1837

Pearce: no records

First record: 2019

Number of records: many

Latest record: 2020

Dorcatoma (Pilosodorcatoma) substriata

Hummel, 1829

Pearce: no records

First record: 2007

Number of records: 4

Latest record: 2021

Dorcatoma (Sternitorcatoma)
flavicornis (Fabricius, 1792)

Pearce: no records
First record: 2019
Number of records: 7
Latest record: 2020

47. Family LYMEXYLIDAE Fleming, 1821

Lymexylon navale (Linnaeus, 1758)

Pearce: no records
First record: 2004
Number of records: 2
Latest record: 2019

48. Family PHLOIOPHILIDAE Kiesenwetter, 1863

Phloiophilus edwardsii Stephens, 1830

Pearce: 1 locality
First subsequent record: 2009
Number of subsequent records: 5
Latest record: 2019

49. Family TROGOSSITIDAE Latreille, 1802

Tenebroides mauritanicus (Linnaeus, 1758)

Pearce: 1 locality (1888)
No subsequent records

Nemozoma elongatum (Linnaeus, 1760)

Pearce: 1 locality
No subsequent records

Thymalus limbatus (Fabricius, 1787)

Pearce: 1 locality (1898)
No subsequent records

51. Family CLERIDAE Latreille, 1802

Tillus elongatus (Linnaeus, 1758)

Pearce: no records
First record: 1995
Number of records: 5
Latest record: 2018

Thanasimus formicarius (Linnaeus, 1758)

Pearce: 8 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2020

Korynetes caeruleus (De Geer, 1775)

Pearce: 2 localities
First subsequent record: 1953
Number of subsequent records: 1
Latest record: 1953

Necrobia ruficollis (Fabricius, 1775)

Pearce: 2 localities
No subsequent records

Necrobia rufipes (De Geer, 1775)

Pearce: 2 localities
No subsequent records

Necrobia violacea (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 1936
Number of subsequent records: 5
Latest record: 2021

52. Family MELYRIDAE Leach, 1815

Aplocnemus impressus (Marsham, 1802)

Pearce: 1 locality
First subsequent record: 2010
Number of subsequent records: 1
Latest record: 2010

Dasytes (Dasytes) niger (Linnaeus, 1760)

Pearce: no records
First record: 2017
Number of records: 1
Latest record: 2017

Dasytes (Mesodasytes) aeratus Stephens, 1830

Pearce: 7 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2020

Dasytes (Mesodasytes) plumbeus (Müller, O.F., 1776)

Pearce: 1 locality
First subsequent record: 2009
Number of subsequent records: 1
Latest record: 2009

Psilothrix viridicoerulea (Geoffroy in Fourcroy, 1785)

Pearce: common along coast
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2020

Axinotarsus marginalis (Laporte, 1840)

Pearce: no records
First record: 2000
Number of records: 2
Latest record: 2006

Cordylepherus viridis (Fabricius, 1787)

Pearce: 3 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2020

Malachius aeneus (Linnaeus, 1758)

Pearce: 3 localities
First subsequent record: 1940
Number of subsequent records: 2
Latest record: 1993

Malachius bipustulatus (Linnaeus, 1758)

Pearce: abundant and widely distributed
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Clanoptilus marginellus (Olivier, 1790)

Pearce: 1 locality
First subsequent record: 1954
Number of subsequent records: 4
Latest record: 2020

Cerapheles terminatus (Ménétries, 1832)

Pearce: 1 locality
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2019

Anthocomus fasciatus (Linnaeus, 1758)

Pearce: 1 locality
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2021

Anthocomus rufus (Herbst, 1784)

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

53. Family BYTURIDAE Gistel, 1848

Byturus ochraceus (Scriba, 1790)

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Byturus tomentosus (De Geer, 1774)

Pearce: 7 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

54. Family SPHINDIDAE Jacquelin du Val, 1860

Sphindus dubius (Gyllenhal, 1808)

Pearce: no records
First record: 2019
Number of records: 4
Latest record: 2019

Aspidiphorus orbiculatus (Gyllenhal, 1808)

Pearce: 1 locality (1840)
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2019

55. Family BIPHYLLIDAE LeConte, 1861

Biphyllus lunatus (Fabricius, 1787)

Pearce: 2 localities
First subsequent record: 1997
Number of subsequent records: many
Latest record: 2021

Diplocoelus fagi (Chevrolat in Guérin-Méneville, 1837)

Pearce: no records
First record: 2018
Number of records: 3
Latest record: 2021

56. Family EROTYLIDAE Latreille, 1802

Cryptophilus integer (Heer, 1841)

Pearce: no records
First record: 2020
Number of records: 3
Latest record: 2020

Dacne bipustulata (Thunberg, 1781)

Pearce: 1 locality (1921)
First subsequent record: 1940
Number of subsequent records: 8
Latest record: 2021

Dacne rufifrons (Fabricius, 1775)

Pearce: 3 localities
First subsequent record: 1988
Number of subsequent records: 2
Latest record: 2020

Triplax aenea (Schaller, 1783)

Pearce: no records
First record: 1994
Number of records: 6
Latest record: 2019

Triplax lacordairii Crotch, 1870

Pearce: 1 locality (1926)
First subsequent record: 1945
Number of subsequent records: 6
Latest record: 2019

Triplax russica (Linnaeus, 1758)

Pearce: 1 locality (1842)
First subsequent record: 1997
Number of subsequent records: many
Latest record: 2016

Tritoma bipustulata Fabricius, 1775

Pearce: 2 localities
First subsequent record: 2011
Number of subsequent records: 1
Latest record: 2011

57. Family MONOTOMIDAE Laporte, 1840

Rhizophagus (Anomophagus) cribratus
Gyllenhal, 1827

Pearce: no records
First record: 1980
Number of records: many
Latest record: 2021

Rhizophagus (Eurhizophagus) depressus
(Fabricius, 1792)

Pearce: 4 localities
First subsequent record: 2001
Number of subsequent records: 6
Latest record: 2018

Rhizophagus (Rhizophagus) bipustulatus
(Fabricius, 1792)

Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Rhizophagus (Rhizophagus) dispar
(Paykull, 1800)

Pearce: no records
First record: 1978
Number of records: many
Latest record: 2020

Rhizophagus (Rhizophagus) ferrugineus
(Paykull, 1800)

Pearce: 2 localities
First subsequent record: 1988
Number of subsequent records: many
Latest record: 2020

Rhizophagus (Rhizophagus) nitidulus
(Fabricius, 1798)

Pearce: no records
First record: 1997
Number of records: 3
Latest record: 2019

Rhizophagus (Rhizophagus)
parallelocollis Gyllenhal, 1827

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: 3
Latest record: 2021

Rhizophagus (Rhizophagus) perforatus
Erichson, 1845

Pearce: 4 localities
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2021

Monotoma angusticollis (Gyllenhal, 1827)

Pearce: 1 locality
No subsequent records

Monotoma bicolor Villa & Villa, 1835

Pearce: no records
First record: 2004
Number of records: 3
Latest record: 2020

Monotoma brevicollis Aubé, 1837

Pearce: 2 localities
First subsequent record: 2006
Number of subsequent records: 4
Latest record: 2021

Monotoma conicicollis Chevrolat in
Guérin-Ménéville, 1837
Pearce: 1 locality
First subsequent record: 2014
Number of subsequent records: 9
Latest record: 2017

Monotoma longicollis (Gyllenhal, 1827)
Pearce: 1 locality
First subsequent record: 1970
Number of subsequent records: 9
Latest record: 2011

Monotoma picipes Herbst, 1793
Pearce: 4 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2021

Monotoma spinicollis Aubé, 1837
Pearce: 1 locality
First subsequent record: 2006
Number of subsequent records: 1
Latest record: 2006

Monotoma testacea Motschulsky, 1845
Pearce: no records
First record: 2004
Number of records: 2
Latest record: 2009

58. Family CRYPTOPHAGIDAE Kirby, 1826

Telmatophilus caricis (Olivier, 1790)
Pearce: 3 localities
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2021

Telmatophilus typhae (Fallén, 1802)
Pearce: 1 locality
First subsequent record: 2001
Number of subsequent records: many
Latest record: 2021

Paramecosoma melanocephalum (Herbst,
1793)
Pearce: 2 localities
First subsequent record: 2017
Number of subsequent records: 1
Latest record: 2017

Henoticus serratus (Gyllenhal, 1808)
Pearce: no records
First record: 1993
Number of records: 2
Latest record: 2001

Cryptophagus acutangulus Gyllenhal,
1827
Pearce: 1 locality
First subsequent record: 2003
Number of subsequent records: 5
Latest record: 2012

Cryptophagus cellaris (Scopoli, 1763)
Pearce: 1 locality (1898)
No subsequent records

Cryptophagus dentatus (Herbst, 1793)
Pearce: 3 localities
First subsequent record: 2000
Number of subsequent records: many
Latest record: 2021

Cryptophagus denticulatus Heer, 1841
Pearce: no records
First record: 1994
Number of records: 5
Latest record: 2018

Cryptophagus distinguendus Sturm, 1845
Pearce: no records
First record: 1994
Number of records: 9
Latest record: 2018

Cryptophagus jakowlewi Reitter, 1888
Pearce: no records
First record: 2003
Number of records: 3
Latest record: 2020

Cryptophagus laticollis Lucas, 1846
Pearce: 3 localities
First subsequent record: 2000
Number of subsequent records: 4
Latest record: 2019

Cryptophagus lycoperdi (Scopoli, 1763)
Pearce: 3 localities
First subsequent record: 1998
Number of subsequent records: many
Latest record: 2021

Cryptophagus micaceus Rey, 1889

Pearce: no records
First record: 2018
Number of records: 2
Latest record: 2020

Cryptophagus populi Paykull, 1800

Pearce: 1 locality
No subsequent records

Cryptophagus pubescens Sturm, 1845

Pearce: 1 locality
First subsequent record: 2019
Number of subsequent records: 4
Latest record: 2019

Cryptophagus punctipennis Brisout de Barneville, C., 1863

Pearce: 2 localities
First subsequent record: 1999
Number of subsequent records: 7
Latest record: 2020

Cryptophagus ruficornis Stephens, 1830

Pearce: no records
First record: 1970
Number of records: 4
Latest record: 2021

Cryptophagus saginatus Sturm, 1845

Pearce: 2 localities
First subsequent record: 2003
Number of subsequent records: 2
Latest record: 2003

Cryptophagus scanicus (Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 1971
Number of subsequent records: many
Latest record: 2021

Cryptophagus scutellatus Newman, 1834

Pearce: no records
First record: 1975
Number of records: 4
Latest record: 2013

Cryptophagus setulosus Sturm, 1845

Pearce: 3 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2019

Micrambe ulicis (Stephens, 1830)

Pearce: abundant on furze-blossoms
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Micrambe woodroffei Johnson, 2007

Pearce: 2 localities
First subsequent record: 1993
Number of subsequent records: 2
Latest record: 2018

Antherophagus pallens (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2021

Antherophagus silaceus (Herbst, 1792)

Pearce: 1 locality
No subsequent records

Antherophagus similis Curtis, 1835

Pearce: 4 localities
First subsequent record: 1971
Number of subsequent records: 8
Latest record: 2009

Caenoscelis ferruginea (Sahlberg, C.R., 1792)

Pearce: 1 locality (1891)
First subsequent record: 2006
Number of subsequent records: 1
Latest record: 2006

Atomaria (Anchicera) apicalis Erichson, 1846

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Atomaria (Anchicera) atricapilla Stephens, 1830

Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Atomaria (Anchicera) basalis Erichson, 1846

Pearce: 1 locality
No subsequent records

Atomaria (Anchicera) fuscata (Schönherr, 1808)
Pearce: 4 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Atomaria (Anchicera) fuscipes (Gyllenhal, 1808)
Pearce: 1 locality
No subsequent records

Atomaria (Anchicera) gutta Newman, 1834
Pearce: 3 localities
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2021

Atomaria (Anchicera) lewisi Reitter, 1877
Pearce: no records
First record: 1975
Number of records: 9
Latest record: 2020

Atomaria (Anchicera) mesomela (Herbst, 1792)
Pearce: 1 locality
First subsequent record: 1964
Number of subsequent records: 9
Latest record: 2019

Atomaria (Anchicera) munda Erichson, 1846
Pearce: no records
First record: 2019
Number of records: 2
Latest record: 2019

Atomaria (Anchicera) nitidula (Marsham, 1802)
Pearce: no records
First record: 1999
Number of records: 6
Latest record: 2019

Atomaria (Anchicera) pusilla (Paykull, 1798)
Pearce: 2 localities
First subsequent record: 2020
Number of subsequent records: 1
Latest record: 2020

Atomaria (Anchicera) rubella Heer, 1841
Pearce: 1 locality
First subsequent record: 1977
Number of subsequent records: 6
Latest record: 2017

Atomaria (Anchicera) rubida Reitter, 1875
Pearce: no records
First record: 2000
Number of records: 2
Latest record: 2000

Atomaria (Anchicera) rubicollis Brisout de Barneville, C., 1863
Pearce: no records
First record: 1998
Number of records: 1
Latest record: 1998

Atomaria (Anchicera) testacea Stephens, 1830
Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Atomaria (Anchicera) turgida Erichson, 1846
Pearce: no records
First record: 2020
Number of records: 1
Latest record: 2020

Atomaria (Atomaria) linearis Stephens, 1830
Pearce: 3 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2019

Atomaria (Atomaria) nigrirostris Stephens, 1830
Pearce: no records
First record: 1966
Number of records: 2
Latest record: 2016

Atomaria (Atomaria) nigriventris Stephens, 1830
Pearce: 1 locality (1892)
No subsequent records

Atomaria (Atomaria) punctithorax Reitter, 1888

Pearce: no records
First record: 1999
Number of records: 1
Latest record: 1999

Atomaria (Atomaria) umbrina (Gyllenhal, 1827)

Pearce: 1 locality
First subsequent record: 2019
Number of subsequent records: 1
Latest record: 2019

Ootypus globosus (Waltl, 1838)

Pearce: 2 localities
First subsequent record: 1996
Number of subsequent records: 2
Latest record: 2019

Ephistemus globulus (Paykull, 1798)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

59. Family SILVANIDAE Kirby, 1837

Uleiota planatus (Linnaeus, 1760)

Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Psammoecus bipunctatus (Fabricius, 1792)

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Nausibius clavicornis (Kugelann, 1794)

Pearce: 2 localities
No subsequent records

Ahasverus advena (Waltl, 1834)

Pearce: no records
First record: 1994
Number of records: 9
Latest record: 2007

Oryzaephilus surinamensis (Linnaeus, 1758)

Pearce: 3 localities
First subsequent record: 2001
Number of subsequent records: 4
Latest record: 2010

Silvanus bidentatus (Fabricius, 1792)

Pearce: no records
First record: 2004
Number of records: 1
Latest record: 2004

Silvanus unidentatus (Olivier, 1790)

Pearce: no records
First record: 1992
Number of records: 7
Latest record: 2019

Silvanoprus fagi (Guérin-Méneville, 1844)

Pearce: no records
First record: 2021
Number of records: 1
Latest record: 2021

60. Family CUCUJIDAE Latreille, 1802

Pediacus dermestoides (Fabricius, 1792)

Pearce: no records
First record: 1971
Number of records: many
Latest record: 2021

61. Family PHALACRIDAE Leach, 1815

Phalacrus caricis Sturm, 1807

Pearce: 3 localities
First subsequent record: 2008
Number of subsequent records: 6
Latest record: 2020

Phalacrus championi Guillebeau, 1892

Pearce: no records
First record: 1992
Number of records: 2
Latest record: 2019

Phalacrus corruscus (Panzer, 1797)

Pearce: 4 localities
First subsequent record: 1935
Number of subsequent records: 6
Latest record: 2019

Phalacrus fimetarius (Fabricius, 1775)

Pearce: no records
First record: 1975
Number of records: 7
Latest record: 2019

Olibrus aeneus (Fabricius, 1792)

Pearce: 6 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Olibrus affinis (Sturm, 1807)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Olibrus corticalis (Panzer, 1797)

Pearce: 3 localities
First subsequent record: 1941
Number of subsequent records: many
Latest record: 2021

Olibrus liquidus Erichson, 1845

Pearce: 4 localities
First subsequent record: 1943
Number of subsequent records: 6
Latest record: 2019

Olibrus millefolii (Paykull, 1800)

Pearce: no records
First record: 1997
Number of records: 3
Latest record: 2019

Olibrus pygmaeus (Sturm, 1807)

Pearce: 1 locality (1909)
First subsequent record: 2001
Number of subsequent records: 5
Latest record: 2015

Stilbus oblongus (Erichson, 1845)

Pearce: 1 locality (1911)
First subsequent record: 1964
Number of subsequent records: 8
Latest record: 2020

Stilbus testaceus (Panzer, 1796)

Pearce: generally common and widely distributed
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2021

**62. Family LAEMOPHLOEIDAE
Ganglbauer, 1899**

Cryptolestes duplicatus (Waltl, 1839)

Pearce: 1 locality
No subsequent records

Cryptolestes ferrugineus (Stephens, 1831)

Pearce: 3 localities
First subsequent record: 1999
Number of subsequent records: 8
Latest record: 2020

**63. Family KATERETIDAE Kirby,
1837**

Kateretes pedicularius (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2020

Kateretes pusillus (Thunberg, 1794)

Pearce: 6 localities
First subsequent record: 1987
Number of subsequent records: many
Latest record: 2021

Kateretes rufilabris (Latreille, 1807)

Pearce: 6 localities
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2020

Brachypterus glaber (Newman, 1834)

Pearce: generally common and widely distributed
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2020

Brachypterus urticae (Fabricius, 1792)

Pearce: 3 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Brachypterus linariae (Stephens, 1830)

Pearce: no records
First record: 1940
Number of records: 1
Latest record: 1940

Brachypterus pulicarius (Linnaeus,
1758)

Pearce: 1 locality
First subsequent record: 2005
Number of subsequent records: 1
Latest record: 2005

Brachypterolus vestitus (Kiesenwetter, 1850)

Pearce: no records

First record: 2017

Number of records: 1

Latest record: 2017

64. Family NITIDULIDAE Latreille, 1802

Epuraea (Epuraea) aestiva (Linnaeus, 1758)

Pearce: 8 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2020

Epuraea (Epuraea) biguttata (Thunberg, 1784)

Pearce: 4 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Epuraea (Epuraea) distincta (Grimmer, 1841)

Pearce: no records

First record: 2001

Number of records: many

Latest record: 2021

Epuraea (Epuraea) fuscicollis (Stephens, 1835)

Pearce: no records

First record: 2019

Number of records: 1

Latest record: 2019

Epuraea (Epuraea) guttata (Olivier, 1811)

Pearce: no records

First record: 2006

Number of records: 4

Latest record: 2019

Epuraea (Epuraea) marseuli Reitter, 1873

Pearce: 2 localities

First subsequent record: 2000

Number of subsequent records: many

Latest record: 2020

Epuraea (Epuraea) melina Erichson, 1843

Pearce: 6 localities

First subsequent record: 1987

Number of subsequent records: 6

Latest record: 2016

Epuraea (Epuraea) pallescens (Stephens, 1835)

Pearce: 1 locality

First subsequent record: 2006

Number of subsequent records: 5

Latest record: 2017

Epuraea (Epuraea) silacea (Herbst, 1783)

Pearce: no records

First record: 1996

Number of records: 1

Latest record: 1996

Epuraea (Epuraeanella) limbata (Fabricius, 1787)

Pearce: 1 locality

No subsequent records

Epuraea (Haptoncus) ocularis Fairmaire, 1849

Pearce: no records

First record: 2020

Number of records: 1

Latest record: 2020

Epuraea (Micruria) melanocephala (Marsham, 1802)

Pearce: 2 localities

First subsequent record: 1970

Number of subsequent records: many

Latest record: 2020

Carpophilus marginellus Motschulsky, 1858

Pearce: no records

First record: 1990

Number of records: many

Latest record: 2020

Pria dulcamarae (Scopoli, 1763)

Pearce: 3 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2021

Meligethes aeneus (Fabricius, 1775)

Pearce: generally common and widely distributed

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2021

Meligethes atramentarius Förster, 1849

Pearce: no records

First record: 1987

Number of records: 1

Latest record: 1987

Meligethes atratus (Olivier, 1790)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Meligethes bidens Brisout de Barneville, C., 1863

Pearce: no records
First record: 1998
Number of records: 1
Latest record: 1998

Meligethes bidentatus Brisout de Barneville, C., 1863

Pearce: no records
First record: 1950
Number of records: 1
Latest record: 1950

Meligethes brunnicornis Sturm, 1845

Pearce: no records
First record: 2012
Number of records: 6
Latest record: 2021

Meligethes carinulatus Förster, 1849

Pearce: 4 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Meligethes difficilis (Heer, 1841)

Pearce: 1 locality (1892)
First subsequent record: 2008
Number of subsequent records: 3
Latest record: 2021

Meligethes erichsonii Brisout de Barneville, C., 1863

Pearce: no records
First record: 1998
Number of records: 7
Latest record: 2020

Meligethes exilis Sturm, 1845

Pearce: no records
First record: 1975
Number of records: 1
Latest record: 1975

Meligethes flavimanus Stephens, 1830

Pearce: 6 localities
First subsequent record: 1965
Number of subsequent records: 9
Latest record: 2021

Meligethes fulvipes Brisout de Barneville, C., 1863

Pearce: 1 locality (1904)
First subsequent record: 1978
Number of subsequent records: 1
Latest record: 1978

Meligethes kunzei Erichson, 1845

Pearce: no records
First record: 2019
Number of records: 2
Latest record: 2021

Meligethes lugubris Sturm, 1845

Pearce: 1 locality (1892)
First subsequent record: 2002
Number of subsequent records: 2
Latest record: 2015

Meligethes matronalis Audisio & Spornraft, 1990

Pearce: no records
First record: 2014
Number of records: 4
Latest record: 2020

Meligethes morosus Erichson, 1845

Pearce: 1 locality
First subsequent record: 1993
Number of subsequent records: 5
Latest record: 2021

Meligethes nigrescens Stephens, 1830

Pearce: 5 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Meligethes obscurus Erichson, 1845

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Meligethes ovatus Sturm, 1845

Pearce: 1 locality
First subsequent record: 2006
Number of subsequent records: 4
Latest record: 2021

Meligethes pedicularius (Gyllenhal, 1808)

Pearce: 1 locality
First subsequent record: 2004
Number of subsequent records: 1
Latest record: 2004

Meligethes persicus Faldermann, 1835

Pearce: 2 localities
First subsequent record: 1998
Number of subsequent records: 1
Latest record: 1998

Meligethes planiusculus (Heer, 1841)

Pearce: 5 localities
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2020

Meligethes rotundicollis Brisout de Barneville, C., 1863

Pearce: 1 locality
No subsequent records

Meligethes ruficornis (Marshall, 1802)

Pearce: 2 localities
First subsequent record: 1978
Number of subsequent records: 4
Latest record: 2021

Meligethes serripes (Gyllenhal, 1827)

Pearce: 1 locality
No subsequent records

Meligethes solidus (Kugelann, 1794)

Pearce: no records
First record: 1993
Number of records: 6
Latest record: 2021

Meligethes umbrosus Sturm, 1845

Pearce: 1 locality
No subsequent records

Meligethes viridescens (Fabricius, 1787)

Pearce: generally common
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2019

Nitidula bipunctata (Linnaeus, 1758)

Pearce: 7 localities
First subsequent record: 1993
Number of subsequent records: 3
Latest record: 2014

Nitidula carnaria (Schaller, 1783)

Pearce: 1 locality (1896)
No subsequent records

Nitidula rufipes (Linnaeus, 1767)

Pearce: 2 localities
No subsequent records

Omosita colon (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1939
Number of subsequent records: 5
Latest record: 2021

Omosita depressa (Linnaeus, 1758)

Pearce: 1 locality
No subsequent records

Omosita discoidea (Fabricius, 1775)

Pearce: 3 localities
First subsequent record: 1979
Number of subsequent records: 5
Latest record: 2021

Soronia grisea (Linnaeus, 1758)

Pearce: 1 locality
First subsequent record: 1998
Number of subsequent records: 7
Latest record: 2021

Soronia punctatissima (Illiger, 1794)

Pearce: 2 localities
First subsequent record: 1993
Number of subsequent records: 1
Latest record: 1993

Pocadius ferrugineus (Fabricius, 1775)

Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2009

Thalycra fervida (Olivier, 1790)

Pearce: no records
First record: 2001
Number of records: 3
Latest record: 2004

Cychramus luteus (Fabricius, 1787)

Pearce: 4 localities
First subsequent record: 1987
Number of subsequent records: 8
Latest record: 2006

Cryptarcha strigata (Fabricius, 1787)

Pearce: no records
First record: 1975
Number of records: 6
Latest record: 2019

Cryptarcha undata (Olivier, 1790)

Pearce: 1 locality

First subsequent record: 1975

Number of subsequent records: 4

Latest record: 2019

Pityophagus ferrugineus (Linnaeus, 1760)

Pearce: 3 localities

First subsequent record: 1964

Number of subsequent records: many

Latest record: 2014

Glischrochilus (*Glischrochilus*)
quadripunctatus (Linnaeus, 1758)

Pearce: 1 locality

No subsequent records

Glischrochilus (*Librodor*) *hortensis*
(Geoffroy in Fourcroy, 1785)

Pearce: no records

First record: 1999

Number of records: 8

Latest record: 2018

Glischrochilus (*Librodor*) *quadriguttatus*
(Fabricius, 1777)

Pearce: no records

First record: 1987

Number of records: 2

Latest record: 2004

65. Family BOTHRIDERIDAE Erichson, 1845

Anommatus duodecimstriatus (Müller,
P.W.J., 1821)

Pearce: 1 locality

First subsequent record: 2010

Number of subsequent records: 2

Latest record: 2011

66. Family CERYLONIDAE Billberg, 1820

Cerylon fagi Brisout de Barneville, C.,
1867

Pearce: 1 locality

No subsequent records

Cerylon ferrugineum Stephens, 1830

Pearce: no records

First record: 1963

Number of records: many

Latest record: 2021

Cerylon histeroides (Fabricius, 1792)

Pearce: 3 localities

First subsequent record: 1988

Number of subsequent records: many

Latest record: 2019

67. Family ALEXIIDAE Imhoff, 1856

Sphaerosoma pilosum (Panzer, 1793)

Pearce: 1 locality

First subsequent record: 1979

Number of subsequent records: 2

Latest record: 2015

68. Family ENDOMYCHIDAE Leach, 1815

Symbiotes latus Redtenbacher, 1849

Pearce: no records

First record: 2018

Number of records: 4

Latest record: 2021

Mycetaea subterranea (Fabricius, 1801)

Pearce: 3 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2013

Endomychus coccineus (Linnaeus, 1758)

Pearce: 3 localities

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2020

Lycoperdina bovistae (Fabricius, 1792)

Pearce: 1 locality

First subsequent record: 2000

Number of subsequent records: 5

Latest record: 2003

69. Family COCCINELLIDAE Latreille, 1807

Coccidula rufa (Herbst, 1783)

Pearce: generally common and widely distributed

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2020

Coccidula scutellata (Herbst, 1783)

Pearce: 1 locality

First subsequent record: 2009

Number of subsequent records: 4

Latest record: 2015

Rhyzobius chrysomeloides (Herbst, 1792)

Pearce: no records
First record: 2019
Number of records: 3
Latest record: 2021

Rhyzobius litura (Fabricius, 1787)

Pearce: generally common and widely distributed
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Rhyzobius lophanthae (Blaisdell, 1892)

Pearce: no records
First record: 2013
Number of records: 4
Latest record: 2021

Hyperaspis pseudopustulata Mulsant, 1853

Pearce: 6 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Nephus quadrimaculatus (Herbst, 1783)

Pearce: no records
First record: 1990
Number of records: 2
Latest record: 1994

Nephus redtenbacheri (Mulsant, 1846)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Scymnus (Neopullus) haemorrhoidalis Herbst, 1797

Pearce: 1 locality
First subsequent record: 1943
Number of subsequent records: many
Latest record: 2020

Scymnus (Neopullus) limbatus Stephens, 1832

Pearce: 3 localities
First subsequent record: 1994
Number of subsequent records: 3
Latest record: 2012

Scymnus (Pullus) auritus Thunberg, 1795

Pearce: no records
First record: 1930
Number of records: many
Latest record: 2019

Scymnus (Pullus) suturalis Thunberg, 1795

Pearce: 1 locality
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2021

Scymnus (Scymnus) femoralis (Gyllenhal, 1827)

Pearce: no records
First record: 1940
Number of records: 2
Latest record: 2012

Scymnus (Scymnus) frontalis (Fabricius, 1787)

Pearce: 5 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2019

Scymnus (Scymnus) nigrinus Kugelann, 1794

Pearce: 1 locality
First subsequent record: 1973
Number of subsequent records: 7
Latest record: 2020

Scymnus (Scymnus) schmidtii Fürsch, 1958

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Stethorus pusillus (Herbst, 1797)

Pearce: no records
First record: 2003
Number of records: many
Latest record: 2015

Chilocorus bipustulatus (Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1932
Number of subsequent records: many
Latest record: 2019

Chilocorus renipustulatus (Scriba, 1791)

Pearce: 4 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2020

Exochomus quadripustulatus (Linnaeus, 1758)

Pearce: 8 localities

First subsequent record: 1936

Number of subsequent records: many

Latest record: 2021

Platynaspis luteorubra (Goeze, 1777)

Pearce: 3 localities

First subsequent record: 1997

Number of subsequent records: many

Latest record: 2019

Halysia sedecimguttata (Linnaeus, 1758)

Pearce: 7 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Psyllobora vigintiduopunctata (Linnaeus, 1758)

Pearce: 7 localities

First subsequent record: 1942

Number of subsequent records: many

Latest record: 2021

Vibidia duodecimguttata (Poda von Neuhaus, 1761)

Pearce: 1 locality

No subsequent records

Anisosticta novemdecimpunctata (Linnaeus, 1758)

Pearce: 5 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2020

Tytthaspis sedecimpunctata (Linnaeus, 1760)

Pearce: 8 localities

First subsequent record: 1973

Number of subsequent records: many

Latest record: 2021

Hippodamia tredecimpunctata (Linnaeus, 1758)

Pearce: no records

First record: 2008

Number of records: 1

Latest record: 2008

Hippodamia variegata (Goeze, 1777)

Pearce: 2 localities

First subsequent record: 1980

Number of subsequent records: 7

Latest record: 2019

Aphidecta oblitterata (Linnaeus, 1758)

Pearce: 5 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2020

Adalia bipunctata (Linnaeus, 1758)

Pearce: generally common and widely distributed

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2021

Adalia decempunctata (Linnaeus, 1758)

Pearce: generally common and widely distributed

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Coccinella hieroglyphica Linnaeus, 1758

Pearce: 5 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2013

Coccinella magnifica Redtenbacher, 1843

Pearce: no records

First record: 1988

Number of records: many

Latest record: 2019

Coccinella septempunctata Linnaeus, 1758

Pearce: very common and widely distributed

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Coccinella undecimpunctata Linnaeus, 1758

Pearce: generally common and widely distributed

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Harmonia axyridis (Pallas, 1773)

Pearce: no records

First record: 2005

Number of records: many

Latest record: 2021

Harmonia quadripunctata (Pontoppidan, 1763)

Pearce: no records
First record: 1975
Number of records: many
Latest record: 2019

Propylea quattuordecimpunctata (Linnaeus, 1758)

Pearce: generally common and widely distributed
First subsequent record: 1932
Number of subsequent records: many
Latest record: 2021

Anatis ocellata (Linnaeus, 1758)

Pearce: 8 localities
First subsequent record: 1943
Number of subsequent records: many
Latest record: 2021

Myrrha octodecimguttata (Linnaeus, 1758)

Pearce: 9 localities
First subsequent record: 1942
Number of subsequent records: many
Latest record: 2019

Calvia quattuordecimguttata (Linnaeus, 1758)

Pearce: 9 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Myzia oblongoguttata (Linnaeus, 1758)

Pearce: 7 localities
First subsequent record: 1945
Number of subsequent records: many
Latest record: 2021

Subcoccinella vigintiquatuorpunctata (Linnaeus, 1758)

Pearce: 7 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

70. Family CORYLOPHIDAE LeConte, 1852

Orthoperus aequalis Sharp, 1885

Pearce: no records
First record: 2003
Number of records: 2
Latest record: 2020

Orthoperus atomus (Gyllenhal, 1808)

Pearce: no records
First record: 2011
Number of records: 2
Latest record: 2018

Orthoperus nigrescens Stephens, 1829

Pearce: no records
First record: 1993
Number of records: many
Latest record: 2020

Sericoderus brevicornis Matthews, A., 1890

Pearce: no records
First record: 2013
Number of records: many
Latest record: 2021

Sericoderus lateralis (Gyllenhal, 1827)

Pearce: 2 localities
First subsequent record: 1990
Number of subsequent records: many
Latest record: 2020

Corylophus cassidoides (Marshall, 1802)

Pearce: 3 localities
First subsequent record: 1991
Number of subsequent records: many
Latest record: 2020

Corylophus sublaevipennis Jacquelin du Val, 1859

Pearce: 2 localities
First subsequent record: 1990
Number of subsequent records: many
Latest record: 2021

Rypobius praetermissus Bowstead, 1999

Pearce: no records
First record: 1931
Number of records: 4
Latest record: 2006

71. Family LATRIDIIDAE Erichson, 1842

Stephostethus alternans (Mannerheim, 1844)

Pearce: no records
First record: 1993
Number of records: 2
Latest record: 2014

Stephostethus angusticollis (Gyllenhal, 1827)

Pearce: 1 locality
No subsequent records

Stephostethus lardarius (De Geer, 1775)

Pearce: 2 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2019

Cartodere (Aridia) bifasciata (Reitter, 1877)

Pearce: no records
First record: 1975
Number of records: many
Latest record: 2021

Cartodere (Aridia) nodifer (Westwood, 1839)

Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Cartodere (Cartodere) constricta (Gyllenhal, 1827)

Pearce: 1 locality
First subsequent record: 2018
Number of subsequent records: 1
Latest record: 2018

Latridius assimilis Mannerheim, 1844

Pearce: no records
First record: 1975
Number of records: 1
Latest record: 1975

Latridius minutus (Linnaeus, 1767)

Pearce: 3 localities (but these could refer to *assimilis* or *porcatus*)
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2019

Latridius porcatus Herbst, 1793

Pearce: no records
First record: 2000
Number of records: many
Latest record: 2021

Enicmus brevicornis (Mannerheim, 1844)

Pearce: no records
First record: 2000
Number of records: many
Latest record: 2021

Enicmus fungicola Thomson, C.G., 1868

Pearce: no records
First record: 2017
Number of records: 6
Latest record: 2019

Enicmus histrio Joy & Tomlin, 1910

Pearce: 1 locality
First subsequent record: 1990
Number of subsequent records: 9
Latest record: 2019

Enicmus rugosus (Herbst, 1793)

Pearce: no records
First record: 2018
Number of records: 2
Latest record: 2019

Enicmus testaceus (Stephens, 1830)

Pearce: no records
First record: 1993
Number of records: many
Latest record: 2021

Enicmus transversus (Olivier, 1790)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Dienerella ruficollis (Marshall, 1802)

Pearce: 1 locality
First subsequent record: 1969
Number of subsequent records: 9
Latest record: 2010

Corticaria crenulata (Gyllenhal, 1827)

Pearce: 4 localities
First subsequent record: 1972
Number of subsequent records: many
Latest record: 2021

Corticaria elongata (Gyllenhal, 1827)

Pearce: 3 localities
First subsequent record: 1972
Number of subsequent records: 5
Latest record: 2007

Corticaria ferruginea Marshall, 1802

Pearce: 3 localities
No subsequent records

Corticaria fulva (Comolli, 1837)

Pearce: 1 locality (1895)

First subsequent record: 1975

Number of subsequent records: 2

Latest record: 1975

Corticaria impressa (Olivier, 1790)

Pearce: 5 localities

First subsequent record: 1972

Number of subsequent records: many

Latest record: 2021

Corticaria punctulata Marsham, 1802

Pearce: 1 locality

First subsequent record: 1975

Number of subsequent records: 9

Latest record: 2013

Corticaria umbilicata (Beck, 1817)

Pearce: no records

First record: 2019

Number of records: 1

Latest record: 2019

Corticarina curta (Wollaston, 1854)

Pearce: 5 localities

First subsequent record: 1989

Number of subsequent records: many

Latest record: 2021

Corticarina minuta (Fabricius, 1792)

Pearce: 5 localities

First subsequent record: 1991

Number of subsequent records: many

Latest record: 2021

Corticarina similata (Gyllenhal, 1827)

Pearce: no records

First record: 1993

Number of records: 2

Latest record: 1993

Cortinicara gibbosa (Herbst, 1793)

Pearce: 8 localities

First subsequent record: 1994

Number of subsequent records: many

Latest record: 2021

Melanophthalma transversalis (Gyllenhal, 1827)

Pearce: 1 locality

First subsequent record: 1994

Number of subsequent records: 1

Latest record: 1994

**72. Family MYCETOPHAGIDAE
Latreille, 1815**

Triphyllus bicolor (Fabricius, 1777)

Pearce: 1 locality

First subsequent record: 1996

Number of subsequent records: 5

Latest record: 2004

Litargus connexus (Geoffroy in Fourcroy, 1785)

Pearce: 3 localities

First subsequent record: 1984

Number of subsequent records: many

Latest record: 2020

Mycetophagus (Ilendus) multipunctatus
Fabricius, 1792

Pearce: no records

First record: 1997

Number of records: 7

Latest record: 2020

Mycetophagus (Mycetophagus)
quadripustulatus (Linnaeus, 1760)

Pearce: 2 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2019

Mycetophagus (Parilendus) quadriguttatus
Müller, P.W.J., 1821

Pearce: no records

First record: 1990

Number of records: 2

Latest record: 2000

Mycetophagus (Philomyces) populi
Fabricius, 1798

Pearce: no records

First record: 2004

Number of records: 2

Latest record: 2004

Mycetophagus (Ulolendus) atomarius
(Fabricius, 1787)

Pearce: no records

First record: 1969

Number of records: 9

Latest record: 2021

Mycetophagus (Ulolendus) piceus
(Fabricius, 1777)
Pearce: no records
First record: 2004
Number of records: 9
Latest record: 2019

Typhaea stercorea (Linnaeus, 1758)
Pearce: 5 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

73. Family CIIDAE Leach in Samouelle, 1819

Octotemnus glabriculus (Gyllenhal, 1827)
Pearce: 5 localities
First subsequent record: 1997
Number of subsequent records: many
Latest record: 2019

Sulcacis nitidus (Fabricius, 1792)
Pearce: no records
First record: 2013
Number of records: 1
Latest record: 2013

Strigocis bicornis (Mellié, 1849)
Pearce: no records
First record: 2004
Number of records: 4
Latest record: 2019

Orthocis alni (Gyllenhal, 1813)
Pearce: no records
First record: 2008
Number of records: 2
Latest record: 2021

Cis bidentatus (Olivier, 1790)
Pearce: 2 localities
First subsequent record: 1966
Number of subsequent records: 8
Latest record: 2021

Cis bilamellatus Wood, 1884
Pearce: no records
First record: 1957
Number of records: many
Latest record: 2021

Cis boleti (Scopoli, 1763)
Pearce: 5 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Cis castaneus (Herbst, 1793)
Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Cis fagi Waltl, 1839
Pearce: no records
First record: 2019
Number of records: 1
Latest record: 2019

Cis festivus (Panzer, 1793)
Pearce: no records
First record: 2004
Number of records: 7
Latest record: 2019

Cis micans (Fabricius, 1792)
Pearce: no records
First record: 2008
Number of records: 2
Latest record: 2011

Cis punctulatus Gyllenhal, 1827
Pearce: no records
First record: 2018
Number of records: 1
Latest record: 2018

Cis pygmaeus (Marsham, 1802)
Pearce: no records
First record: 1968
Number of records: 8
Latest record: 2021

Cis submicans Abiella de Perrin, 1874
Pearce: no records
First record: 1992
Number of records: many
Latest record: 2021

Cis vestitus Mellié, 1849
Pearce: no records
First record: 2000
Number of records: many
Latest record: 2021

Cis villosulus (Marsham, 1802)

Pearce: 2 localities

First subsequent record: 1992

Number of subsequent records: 3

Latest record: 2015

Ennearthron cornutum (Gyllenhal, 1827)

Pearce: 2 localities

First subsequent record: 1992

Number of subsequent records: many

Latest record: 2019

74. Family TETRATOMIDAE Billberg, 1820

Tetratoma desmarestii Latreille, 1807

Pearce: no records

First record: 2004

Number of records: 1

Latest record: 2004

Tetratoma fungorum Fabricius, 1790

Pearce: no records

First record: 2011

Number of records: 1

Latest record: 2011

Hallomenus binotatus (Quensel, 1790)

Pearce: 1 locality

First subsequent record: 1975

Number of subsequent records: 5

Latest record: 2018

75. Family MELANDRYIDAE Leach, 1815

Abdera biflexuosa (Curtis, 1829)

Pearce: 1 locality

First subsequent record: 2000

Number of subsequent records: 7

Latest record: 2019

Abdera flexuosa (Paykull, 1799)

Pearce: no records

First record: 1995

Number of records: 3

Latest record: 2019

Anisoxya fuscula (Illiger, 1798)

Pearce: no records

First record: 2015

Number of records: 1

Latest record: 2015

Melandrya caraboides (Linnaeus 1760)

Pearce: 1 locality

First subsequent record: 2005

Number of subsequent records: 1

Latest record: 2005

Orchesia (Clinocara) minor Walker, 1837

Pearce: 2 localities

First subsequent record: 1965

Number of subsequent records: many

Latest record: 2021

Orchesia (Clinocara) undulata Kraatz, 1853

Pearce: no records

First record: 1996

Number of records: many

Latest record: 2019

Orchesia (Orchesia) micans (Panzer, 1793)

Pearce: 1 locality (1891)

First subsequent record: 2019

Number of subsequent records: 1

Latest record: 2019

Conopalpus testaceus (Olivier, 1790)

Pearce: 2 localities

First subsequent record: 1954

Number of subsequent records: 3

Latest record: 2018

76. Family MORDELLIDAE Latreille, 1802

Mordella leucaspis Küster, 1849

Pearce: 2 localities for *M. aculeata* could be this species or *M. holomelaena*

First record: 1987

Number of records: 1

Latest record: 1987

Tomoxia bucephala Costa, A., 1854

Pearce: 1 locality (1870)

First subsequent record: 1992

Number of subsequent records: 6

Latest record: 2011

Variimorda villosa (Schrank, 1781)

Pearce: 1 locality

First subsequent record: 1974

Number of subsequent records: many

Latest record: 2021

Mordellistena brevicauda (Boheman, 1849)

Pearce: 1 locality
No subsequent records

Mordellistena humeralis (Linnaeus, 1758)

Pearce: 1 locality
First subsequent record: 2007
Number of subsequent records: 3
Latest record: 2020

Mordellistena neuwaldeggiana (Panzer, 1796)

Pearce: 1 locality
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2021

Mordellistena parvula (Gyllenhal, 1827)

Pearce: 1 locality
First subsequent record: 2002
Number of subsequent records: many
Latest record: 2021

Mordellistena pumila (Gyllenhal, 1810)

Pearce: 4 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2019

Mordellistena variegata (Fabricius, 1798)

Pearce: no records
First record: 2008
Number of records: 3
Latest record: 2012

Mordellochroa abdominalis (Fabricius, 1775)

Pearce: 2 localities
First subsequent record: 1993
Number of subsequent records: many
Latest record: 2021

77. Family RIPIPHORIDAE Gemminger, 1870

Metoecus paradoxus (Linnaeus, 1760)

Pearce: 1 locality (1880)
First subsequent record: 1997
Number of subsequent records: 2
Latest record: 2019

78. Family ZOPHERIDAE Solier, 1834

Pycnomerus fuliginosus Erichson, 1842

Pearce: no records
First record: 2008
Number of records: many
Latest record: 2021

Colydium elongatum (Fabricius, 1787)

Pearce: no records
First record: 1992
Number of records: 3
Latest record: 2020

Orthocerus clavicornis (Linnaeus, 1758)

Pearce: 1 locality
No subsequent records

Synchita humeralis (Fabricius, 1792)

Pearce: no records
First record: 2009
Number of records: 1
Latest record: 2009

Synchita undata Guérin-Ménéville, 1844

Pearce: no records
First record: 2008
Number of records: 2
Latest record: 2021

Bitoma crenata (Fabricius, 1775)

Pearce: 2 localities
First subsequent record: 1991
Number of subsequent records: many
Latest record: 2019

Langelandia anophthalma Aubé, 1842

Pearce: no records
First record: 2021
Number of records: 4
Latest record: 2021

79. Family TENEBRIONIDAE Latreille, 1802

Lagria hirta (Linnaeus, 1758)

Pearce: 12 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Alphitobius diaperinus (Panzer, 1796)

Pearce: no records
First record: 2004
Number of records: 2
Latest record: 2006

Blaps mucronata Latreille, 1804

Pearce: 6 localities
No subsequent records

Eledona agricola (Herbst, 1783)

Pearce: no records
First record: 1988
Number of records: 8
Latest record: 2018

Nalassus laevioctostriatus (Goeze, 1777)

Pearce: generally common
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Helops caeruleus (Linnaeus, 1758)

Pearce: 1 locality
First subsequent record: 2011
Number of subsequent records: 4
Latest record: 2016

Melanimon tibialis (Fabricius, 1781)

Pearce: 2 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2019

Opatrum sabulosum (Linnaeus, 1760)

Pearce: 3 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2020

Palorus ratzeburgii (Wissmann, 1848)

Pearce: 2 localities
No subsequent records

Phylan gibbus (Fabricius, 1775)

Pearce: 4 localities
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2018

Tenebrio molitor Linnaeus, 1758

Pearce: 1 locality
First subsequent record: 1999
Number of subsequent records: 3
Latest record: 2019

Tenebrio obscurus Fabricius, 1792

Pearce: 3 localities
No subsequent records

Tribolium castaneum (Herbst, 1797)

Pearce: 4 localities
First subsequent record: 2018
Number of subsequent records: 1
Latest record: 2018

Crypticus quisquilius (Linnaeus, 1760)

Pearce: 2 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Alphitophagus bifasciatus (Say, 1824)

Pearce: no records
First record: 2011
Number of records: 1
Latest record: 2011

Diaperis boleti (Linnaeus, 1758)

Pearce: no records
First record: 1964
Number of records: many
Latest record: 2019

Corticeus fraxini (Kugelann, 1794)

Pearce: no records
First record: 1998
Number of records: 3
Latest record: 2016

Corticeus unicolor Piller & Mitterpacher, 1783

Pearce: no records
First record: 2016
Number of records: 4
Latest record: 2017

Myrmexixenus vaporariorum Guérin-Méneville, 1843

Pearce: no records
First record: 2008
Number of records: 4
Latest record: 2011

Phaleria cadaverina (Fabricius, 1792)

Pearce: 2 localities
First subsequent record: 1953
Number of subsequent records: 6
Latest record: 2007

Scaphidema metallica (Fabricius, 1792)

Pearce: 2 localities
First subsequent record: 1997
Number of subsequent records: 6
Latest record: 2011

Prionychus ater (Fabricius, 1775)

Pearce: no records
First record: 1998
Number of records: 3
Latest record: 2019

Gonodera luperus (Herbst, 1783)

Pearce: 1 locality
First subsequent record: 2018
Number of subsequent records: 1
Latest record: 2018

Isomira murina (Linnaeus, 1758)

Pearce: 7 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Pseudocistela ceramboides (Linnaeus, 1758)

Pearce: 2 localities (1823, 1867)
No subsequent records

Mycetochara humeralis (Fabricius, 1787)

Pearce: 1 locality
No subsequent records

Cteniopus sulphureus (Linnaeus, 1758)

Pearce: 1 locality
First subsequent record: 1969
Number of subsequent records: 5
Latest record: 2015

Omophlus pubescens (Linnaeus, 1767)

Pearce: 2 localities
First subsequent record: 1989
Number of subsequent records: many
Latest record: 2019

80. Family OEDEMERIDAE Latreille, 1810

Ischnomera cyanea (Fabricius, 1792)

Pearce: 4 localities for *caerulea/cyanea*
First subsequent record: 1994
Number of subsequent records: 6
Latest record: 2021

Ischnomera sanguinicollis (Fabricius, 1787)

Pearce: no records
First record: 2009
Number of records: 1
Latest record: 2009

Nacerdes melanura (Linnaeus, 1758)

Pearce: 1 locality (1886)
First subsequent record: 1964
Number of subsequent records: 6
Latest record: 2018

Oedemera (Oedemera) lurida (Marsham, 1802)

Pearce: 4 localities
First subsequent record: 1974
Number of subsequent records: many
Latest record: 2021

Oedemera (Oedemera) nobilis (Scopoli, 1763)

Pearce: 5 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Oedemera (Oncomera) femoralis Olivier, 1803

Pearce: 1 locality
First subsequent record: 1974
Number of subsequent records: many
Latest record: 2021

81. Family MELOIDAE Gyllenhal, 1810

Lytta vesicatoria (Linnaeus, 1758)

Pearce: 9 localities
First subsequent record: 2009
Number of subsequent records: 4
Latest record: 2020

Meloe (Eurymeloe) rugosus Marsham, 1802

Pearce: no records
First record: 2003
Number of records: many
Latest record: 2020

Meloe (Meloe) proscarabaeus Linnaeus, 1758

Pearce: generally common and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Meloe (Meloe) violaceus Marsham, 1802

Pearce: 5 localities
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2021

Meloe (Treiodous) autumnalis Olivier,
1792

Pearce: 1 locality
No subsequent records

Sitaris muralis (Forster, 1771)

Pearce: no records
First record: 2015
Number of records: 4
Latest record: 2021

84. Family PYROCHROIDAE Latreille, 1806

Pyrochroa coccinea (Linnaeus, 1760)

Pearce: no records
First record: 1993
Number of records: 4
Latest record: 2017

Pyrochroa serraticornis (Scopoli, 1763)

Pearce: 6 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2021

85. Family SALPINGIDAE Leach, 1815

Aglenus brunneus (Gyllenhal, 1813)

Pearce: no records
First record: 2008
Number of records: 1
Latest record: 2008

Lissodema denticollis (Gyllenhal, 1813)

Pearce: 2 localities
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2019

Sphaeriestes castaneus (Panzer, 1796)

Pearce: 4 localities
No subsequent records

Sphaeriestes reyi (Abeille de Perrin, 1874)

Pearce: 1 locality
No subsequent records

Sphaeriestes stockmanni (Biström, 1977)

[2012: *Sphaeriestes ater*]

Pearce: 1 locality (1925)
No subsequent records

Vincenzellus ruficollis (Panzer, 1794)

Pearce: 1 locality (1881)
First subsequent record: 1992
Number of subsequent records: 7
Latest record: 2021

Salpingus planirostris (Fabricius, 1787)

Pearce: 5 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Salpingus ruficollis (Linnaeus, 1760)

Pearce: 1 locality
First subsequent record: 2004
Number of subsequent records: many
Latest record: 2021

86. Family ANTHICIDAE Latreille, 1819

Anthicus angustatus Curtis, 1838

Pearce: 2 localities
First subsequent record: 1998
Number of subsequent records: 1
Latest record: 1998

Anthicus antherinus (Linnaeus, 1760)

Pearce: 6 localities
First subsequent record: 1962
Number of subsequent records: many
Latest record: 2021

Anthicus tristis Schmidt, W.L.E., 1842

Pearce: 5 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2021

Cordicollis instabilis (Schmidt, W.L.E.,
1842)

Pearce: 2 localities
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2020

Cyclodinus constrictus (Curtis, 1838)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: 8
Latest record: 2011

Cyclodinus salinus (Crotch, 1867)

Pearce: 2 localities

First subsequent record: 2004

Number of subsequent records: 3

Latest record: 2007

Omonadus bifasciatus (Rossi, 1792)

Pearce: no records

First record: 1995

Number of records: 6

Latest record: 2006

Omonadus floralis (Linnaeus, 1758)

Pearce: 8 localities

First subsequent record: 1978

Number of subsequent records: many

Latest record: 2021

Omonadus formicarius (Goeze, 1777)

Pearce: no records

First record: 1995

Number of records: many

Latest record: 2021

Stricticollis tobias (Marseul, 1879)

Pearce: no records

First record: 1996

Number of records: 3

Latest record: 2003

Notoxus monoceros (Linnaeus, 1760)

Pearce: 2 localities

First subsequent record: 1998

Number of subsequent records: many

Latest record: 2019

87. Family ADERIDAE Csiki, 1909

Aderus populneus (Creutzer in Panzer, 1796)

Pearce: 1 locality (1916)

First subsequent record: 1997

Number of subsequent records: 3

Latest record: 2021

Euglenes oculatus (Paykull, 1798)

Pearce: no records

First record: 1996

Number of records: many

Latest record: 2020

88. Family SCRAPTIIDAE Gistel, 1848

Scraptia dubia (Olivier, 1790)

Pearce: 1 locality (1842)

No subsequent records

Anaspis (Anaspis) fasciata (Forster, 1771)

Pearce: 2 localities

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2020

Anaspis (Anaspis) frontalis (Linnaeus, 1758)

Pearce: generally common

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2019

Anaspis (Anaspis) garneysi Fowler, 1889

Pearce: 1 locality

First subsequent record: 1987

Number of subsequent records: many

Latest record: 2021

Anaspis (Anaspis) lurida Stephens, 1832

Pearce: 1 locality

First subsequent record: 1986

Number of subsequent records: many

Latest record: 2020

Anaspis (Anaspis) maculata (Geoffroy in Fourcroy, 1785)

Pearce: generally common

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2021

Anaspis (Anaspis) pulicaria Costa, A., 1854

Pearce: 9 localities

First subsequent record: 1969

Number of subsequent records: many

Latest record: 2021

Anaspis (Anaspis) regimbarti Schilsky, 1895

Pearce: 9 localities

First subsequent record: 1969

Number of subsequent records: many

Latest record: 2020

Anaspis (Anaspis) thoracica (Linnaeus, 1758)

Pearce: no records

First record: 1989

Number of records: many

Latest record: 2018

Anaspis (Nassipa) costai Emery, 1876

Pearce: no records
First record: 1996
Number of records: 9
Latest record: 2013

Anaspis (Nassipa) rufilabris (Gyllenhal, 1827)

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2020

89. Family CERAMBYCIDAE Latreille, 1802

Prionus coriarius (Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 1981
Number of subsequent records: 5
Latest record: 2021

Rhagium (Hagrium) bifasciatum Fabricius, 1775

Pearce: generally common and widely distributed
First subsequent record: 1959
Number of subsequent records: many
Latest record: 2021

Rhagium (Megarhagium) mordax (De Geer, 1775)

Pearce: 4 localities
First subsequent record: 1958
Number of subsequent records: many
Latest record: 2020

Rhagium (Rhagium) inquisitor (Linnaeus, 1758)

Pearce: 1 locality
No subsequent records

Stenocorus meridianus (Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Grammoptera ruficornis (Fabricius, 1781)

Pearce: generally common and widely distributed
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Leptura aurulenta Fabricius, 1792

Pearce: no records
First record: 2008
Number of records: 1
Latest record: 2008

Leptura quadrifasciata Linnaeus, 1758

Pearce: 3 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Stictoleptura rubra (Linnaeus, 1758)

Pearce: no records
First record: 2020
Number of records: 3
Latest record: 2020

Stictoleptura scutellata (Fabricius, 1781)

Pearce: no records
First record: 2016
Number of records: 1
Latest record: 2016

Paracorymbia fulva (De Geer, 1775)

Pearce: no records
First record: 1987
Number of records: 7
Latest record: 2021

Pachytodes cerambyciformis (Schränk, 1781)

Pearce: no records
First record: 1945
Number of records: many
Latest record: 2008

Alosterna tabacicolor (De Geer, 1775)

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Pseudovadonia livida (Fabricius, 1777)

Pearce: 9 localities
First subsequent record: 1951
Number of subsequent records: many
Latest record: 2021

Rutpela maculata (Poda von Neuhaus, 1761)

Pearce: generally common
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Stenurella (Nigrostenurella) nigra
(Linnaeus, 1758)
Pearce: 2 localities
No subsequent records

Stenurella (Stenurella) melanura
(Linnaeus, 1758)
Pearce: 4 localities
First subsequent record: 1940
Number of subsequent records: many
Latest record: 2021

Asemum striatum (Linnaeus, 1758)
Pearce: 5 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2020

Tetropium gabrieli Weise, 1905
Pearce: no records
First record: 1958
Number of records: 1
Latest record: 1958

Arhopalus ferus (Mulsant, 1839)
Pearce: 1 locality
First subsequent record: 1936
Number of subsequent records: 8
Latest record: 1975

Arhopalus rusticus (Linnaeus, 1758)
Pearce: no records
First record: 1958
Number of records: many
Latest record: 2021

Gracilia minuta (Fabricius, 1781)
Pearce: 2 localities
No subsequent records

Obrium brunneum (Fabricius, 1792)
Pearce: no records
First record: 1936
Number of records: many
Latest record: 2019

Nathrius brevipennis (Mulsant, 1839)
Pearce: no records
First record: 1904
Number of records: 1
Latest record: 1904

Molorchus minor (Linnaeus, 1758)
Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: 6
Latest record: 2008

Glaphyra umbellatarum (Schreber, 1759)
Pearce: 1 locality
No subsequent records

Aromia moschata (Linnaeus, 1758)
Pearce: 3 localities
First subsequent record: 1934
Number of subsequent records: 4
Latest record: 1998

Callidium violaceum (Fabricius, 1775)
Pearce: 1 locality
No subsequent records

Phymatodes testaceus (Linnaeus, 1758)
Pearce: 1 locality
First subsequent record: 1988
Number of subsequent records: 6
Latest record: 2021

Poecilium alni (Linnaeus, 1767)
Pearce: 4 localities
No subsequent records

Clytus arietis (Linnaeus, 1758)
Pearce: 6 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Anaglyptus mysticus (Linnaeus, 1758)
Pearce: 1 locality
First subsequent record: 1996
Number of subsequent records: 4
Latest record: 2011

Agapanthia villosoviridescens (De Geer, 1775)
Pearce: no records
First record: 2009
Number of records: many
Latest record: 2021

Pogonocherus hispidulus (Piller & Mitterpacher, 1783)
Pearce: 3 localities
First subsequent record: 1940
Number of subsequent records: many
Latest record: 2021

Pogonocherus hispidus (Linnaeus, 1758)

Pearce: 3 localities

First subsequent record: 1964

Number of subsequent records: many

Latest record: 2021

Acanthocinus aedilis (Linnaeus, 1758)

Pearce: 1 locality (1912)

No subsequent records

Leiopus linnei Wallin, Nylander &
Kvamme, 2009

Pearce: no records (but see *Leiopus nebulosus*)

First record: 2019

Number of records: 1

Latest record: 2019

Leiopus nebulosus (Linnaeus, 1758)

Pearce: 8 localities (but may be this species or
Leiopus linnei)

First subsequent record: 2010

Number of subsequent records: 8

Latest record: 2019

Saperda populnea (Linnaeus, 1758)

Pearce: 1 locality

First subsequent record: 1980

Number of subsequent records: 2

Latest record: 1997

Phytoecia cylindrica (Linnaeus, 1758)

Pearce: 1 locality

First subsequent record: 1947

Number of subsequent records: many

Latest record: 2021

Tetrops praeustus (Linnaeus, 1758)

Pearce: 4 localities

First subsequent record: 1942

Number of subsequent records: many

Latest record: 2021

90. Family MEGALOPODIDAE Latreille, 1802

Zeugophora subspinosus (Fabricius, 1781)

Pearce: 1 locality (1866)

No subsequent records

91. Family ORSCODACNIDAE Thomson, C. G., 1859

Orsodacne cerasi (Linnaeus, 1758)

Pearce: no records

First record: 1987

Number of records: 4

Latest record: 2018

Orsodacne humeralis Latreille, 1804

Pearce: no records

First record: 2019

Number of records: 1

Latest record: 2019

92. Family CHRYSOMELIDAE Latreille, 1802

Donacia aquatica (Linnaeus, 1758)

Pearce: 2 localities

No subsequent records

Donacia bicolora Zschach, 1788

Pearce: 7 localities

First subsequent record: 1980

Number of subsequent records: many

Latest record: 2020

Donacia cinerea Herbst, 1784

Pearce: 1 locality

No subsequent records

Donacia clavipes Fabricius, 1792

Pearce: 3 localities

First subsequent record: 1935

Number of subsequent records: 8

Latest record: 2014

Donacia crassipes Fabricius, 1775

Pearce: 1 locality

No subsequent records

Donacia dentata Hoppe, 1795

Pearce: 1 locality

No subsequent records

Donacia impressa Paykull, 1799

Pearce: 2 localities

First subsequent record: 1996

Number of subsequent records: 6

Latest record: 2008

Donacia marginata Hoppe, 1795

Pearce: 2 localities
First subsequent record: 1992
Number of subsequent records: many
Latest record: 2021

Donacia obscura Gyllenhal, 1813

Pearce: 1 locality
No subsequent records

Donacia semicuprea Panzer, 1796

Pearce: 2 localities
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2013

Donacia simplex Fabricius, 1775

Pearce: 5 localities
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2020

Donacia sparganii Ahrens, 1810

Pearce: 1 locality
No subsequent records

Donacia thalassina Germar, 1811

Pearce: 2 localities (1895)
First subsequent record: 1934
Number of subsequent records: 3
Latest record: 2013

Donacia versicolore (Brahm, 1790)

Pearce: 1 locality
First subsequent record: 1997
Number of subsequent records: many
Latest record: 2021

Donacia vulgaris Zschach, 1788

Pearce: 1 locality
First subsequent record: 1962
Number of subsequent records: many
Latest record: 2021

Plateumaris (Euplateumaris) discolor
(Panzer, 1795)

Pearce: 6 localities
First subsequent record: 1932
Number of subsequent records: many
Latest record: 2021

Plateumaris (Euplateumaris) sericea
(Linnaeus, 1758)

Pearce: generally common
First subsequent record: 1969
Number of subsequent records: many
Latest record: 2019

Plateumaris (Plateumaris) rustica (Kunze,
1818)

Pearce: 8 localities
First subsequent record: 1992
Number of subsequent records: 1
Latest record: 1992

Bruchidius cisti (Fabricius, 1775)

Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Bruchidius varius (Olivier, 1795)

Pearce: no records
First record: 2008
Number of records: 8
Latest record: 2021

Bruchidius villosus (Fabricius, 1792)

Pearce: 1 locality
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2017

Bruchus atomarius (Linnaeus, 1760)

Pearce: 3 localities
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2019

Bruchus loti Paykull, 1800

Pearce: 2 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Bruchus pisorum (Linnaeus, 1758)

Pearce: 2 localities
No subsequent records

Bruchus rufimanus Boheman, 1833

Pearce: 5 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2019

Bruchus rufipes Herbst, 1783

Pearce: no records
First record: 1954
Number of records: many
Latest record: 2021

Crioceris asparagi (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 2001
Number of subsequent records: 3
Latest record: 2021

Lema cyanella (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2017

Lilioceris lili (Scopoli, 1763)

Pearce: no records
First record: 1977
Number of records: many
Latest record: 2021

Oulema duftschmidi (Redtenbacher, 1874)
[2012: *Oulema rufocyanea*]

Pearce: no records (but see *melanopus*)
First record: 1978
Number of records: many
Latest record: 2016

Oulema melanopus (Linnaeus, 1758)

Pearce: generally common and widely distributed
(but probably includes *duftschmidi*)
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Oulema obscura (Stephens, 1831)

Pearce: 2 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Cryptocephalus (Burlinius) bilineatus
(Linnaeus, 1767)

Pearce: 1 locality
First subsequent record: 2005
Number of subsequent records: 2
Latest record: 2020

Cryptocephalus (Burlinius) fulvus (Goeze, 1777)

Pearce: 7 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2020

Cryptocephalus (Burlinius) labiatus
(Linnaeus, 1760)

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Cryptocephalus (Burlinius) punctiger
Paykull, 1799

Pearce: 1 locality
No subsequent records

Cryptocephalus (Burlinius) pusillus
Fabricius, 1777

Pearce: 5 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Cryptocephalus (Cryptocephalus)
aureolus Suffrian, 1847

Pearce: 3 localities
First subsequent record: 1977
Number of subsequent records: many
Latest record: 2020

Cryptocephalus (Cryptocephalus)
biguttatus (Scopoli, 1763)

Pearce: 1 locality
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Cryptocephalus (Cryptocephalus)
bipunctatus (Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2020

Cryptocephalus (Cryptocephalus)
hypochaeridis (Linnaeus, 1758)

Pearce: 1 locality (1924)
First subsequent record: 1943
Number of subsequent records: many
Latest record: 2015

Cryptocephalus (Cryptocephalus) moraei
(Linnaeus, 1758)
Pearce: 5 localities
First subsequent record: 1997
Number of subsequent records: many
Latest record: 2021

Cryptocephalus (Cryptocephalus)
parvulus Müller, O.F., 1776
Pearce: 3 localities
First subsequent record: 2001
Number of subsequent records: 3
Latest record: 2011

Cryptocephalus (Cryptocephalus)
primarius Harold, 1872
Pearce: no records
First record: 1949
Number of records: many
Latest record: 2019

Cryptocephalus (Cryptocephalus)
sexpunctatus (Linnaeus, 1758)
Pearce: 1 locality
No subsequent records

Clytra quadripunctata (Linnaeus, 1758)
Pearce: 2 localities
First subsequent record: 1980
Number of subsequent records: 8
Latest record: 2018

Oomorphus concolor (Sturm, 1807)
Pearce: 4 localities
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2019

Cassida denticollis Suffrian, 1844
Pearce: no records
First record: 1982
Number of records: 2
Latest record: 2021

Cassida flaveola Thunberg, 1794
Pearce: 6 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2019

Cassida hemisphaerica Herbst, 1799
Pearce: 2 localities
First subsequent record: 2016
Number of subsequent records: 2
Latest record: 2016

Cassida murraea Linnaeus, 1767
Pearce: 5 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2021

Cassida nebulosa Linnaeus, 1758
Pearce: no records
First record: 1979
Number of records: many
Latest record: 2013

Cassida nobilis Linnaeus, 1758
Pearce: 6 localities
First subsequent record: 1935
Number of subsequent records: 6
Latest record: 2021

Cassida prasina Illiger, 1798
Pearce: 1 locality (1841)
First subsequent record: 1998
Number of subsequent records: many
Latest record: 2020

Cassida rubiginosa Müller, O.F., 1776
Pearce: 9 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Cassida sanguinosa Suffrian, 1844
Pearce: no records
First record: 2004
Number of records: 6
Latest record: 2014

Cassida vibex Linnaeus, 1767
Pearce: 1 locality
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Cassida viridis Linnaeus, 1758
Pearce: 3 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2019

Cassida vittata de Villers, 1789
Pearce: 4 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2020

Pilemostoma fastuosum (Schaller, 1783)

Pearce: 1 locality (1877)
First subsequent record: 1991
Number of subsequent records: 1
Latest record: 1991

Timarcha goettingensis (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2021

Timarcha tenebricosa (Fabricius, 1775)

Pearce: generally common and widely distributed
First subsequent record: 1932
Number of subsequent records: many
Latest record: 2021

Gastrophysa polygoni (Linnaeus, 1758)

Pearce: generally common and widely distributed
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Gastrophysa viridula De Geer 1775

Pearce: 2 localities
First subsequent record: 1977
Number of subsequent records: many
Latest record: 2021

Phaedon (Paraphaedon) tumidulus
(Germar, 1824)

Pearce: generally common and widely distributed
First subsequent record: 1966
Number of subsequent records: many
Latest record: 2021

Phaedon (Phaedon) armoraciae
(Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2021

Phaedon (Phaedon) cochleariae
(Fabricius, 1792)

Pearce: 4 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2021

Prasocuris (Hydrothassa) glabra (Herbst, 1783)

Pearce: 4 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2013

Prasocuris (Hydrothassa) marginella
(Linnaeus, 1758)

Pearce: 8 localities
First subsequent record: 1967
Number of subsequent records: many
Latest record: 2021

Prasocuris (Prasocuris) junci (Brahm, 1790)

Pearce: generally common and widely distributed
First subsequent record: 1927
Number of subsequent records: many
Latest record: 2021

Prasocuris (Prasocuris) phellandrii
(Linnaeus, 1758)

Pearce: generally common and widely distributed
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Chrysomela populi Linnaeus, 1758

Pearce: 7 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Plagiodera versicolora (Laicharting, 1781)

Pearce: no records
First record: 1975
Number of records: 7
Latest record: 2019

Plagiosterna aenea (Linnaeus, 1758)
[2012: *Chrysomela aenea*]

Pearce: no records
First record: 1994
Number of records: 2
Latest record: 2007

Phratora (Phratora) vulgatissima
(Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Phratora (Phyllodecta) laticollis (Suffrian, 1851)

Pearce: 5 localities

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2021

Phratora (Phyllodecta) vitellinae (Linnaeus, 1758)

Pearce: generally common

First subsequent record: 1982

Number of subsequent records: 7

Latest record: 2018

Chrysolina (Chrysolina) banksii (Fabricius, 1775)

Pearce: 8 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Chrysolina (Chrysolina) staphylaea (Linnaeus, 1758)

Pearce: 4 localities

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2019

Chrysolina (Chrysolinopsis) americana (Linnaeus, 1758)

Pearce: no records

First record: 2007

Number of records: many

Latest record: 2021

Chrysolina (Colaphodes) haemoptera (Linnaeus, 1758)

Pearce: 11 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2020

Chrysolina (Colaphosoma) sturmi (Westhoff, 1882)

Pearce: 8 localities

First subsequent record: 2008

Number of subsequent records: 4

Latest record: 2016

Chrysolina (Erythrochrysa) polita (Linnaeus, 1758)

Pearce: generally common

First subsequent record: 1937

Number of subsequent records: many

Latest record: 2021

Chrysolina (Hypericia) brunsvicensis (Gravenhorst, 1807)

Pearce: 3 localities

First subsequent record: 2006

Number of subsequent records: 2

Latest record: 2008

Chrysolina (Hypericia) hyperici (Forster, 1771)

Pearce: 8 localities

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2019

Chrysolina (Sphaeromela) varians (Schaller, 1783)

Pearce: 1 locality (1893)

First subsequent record: 1963

Number of subsequent records: 4

Latest record: 2008

Chrysolina (Stichoptera) sanguinolenta (Linnaeus, 1758)

Pearce: 5 localities

No subsequent records

Chrysolina (Sulcicollis) oricalcia (Müller, O.F., 1776)

Pearce: 2 localities

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2021

Chrysolina (Synerga) herbacea (Duftschmid, 1825)

Pearce: 2 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Gonioctena (Gonioctena) viminalis (Linnaeus, 1758)

Pearce: 3 localities

First subsequent record: 1978

Number of subsequent records: 9

Latest record: 1996

Gonioctena (Goniomena) pallida (Linnaeus, 1758)

Pearce: 1 locality (1894)

First subsequent record: 1978

Number of subsequent records: 7

Latest record: 2017

Gonioctena (Spartophila) olivacea
(Forster, 1771)
Pearce: 2 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2020

Luperomorpha xanthodera (Fairmaire,
1888)
Pearce: no records
First record: 2015
Number of records: 1
Latest record: 2015

Agelastica alni (Linnaeus, 1758)
Pearce: no records
First record: 2019
Number of records: many
Latest record: 2021

Sermylassa halensis (Linnaeus, 1767)
Pearce: 8 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Calomicrus circumfusus (Marsham, 1802)
Pearce: 4 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2020

Luperus flavipes (Linnaeus, 1767)
Pearce: 4 localities
First subsequent record: 1980
Number of subsequent records: 6
Latest record: 2012

Luperus longicornis (Fabricius, 1781)
Pearce: generally common
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2020

Phyllobrotica quadrimaculata (Linnaeus,
1758)
Pearce: 4 localities
First subsequent record: 1998
Number of subsequent records: 8
Latest record: 2020

Galeruca laticollis Sahlberg, C.R., 1838
Pearce: 2 localities (1893, 1919)
First subsequent record: 1965
Number of subsequent records: 1
Latest record: 1965

Galeruca tanaceti (Linnaeus, 1758)
Pearce: 3 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2020

Galerucella (Galerucella) nymphaeae
(Linnaeus, 1758)
Pearce: 2 localities
First subsequent record: 1994
Number of subsequent records: 3
Latest record: 2017

Galerucella (Galerucella) sagittariae
(Gyllenhal, 1813)
Pearce: 5 localities
First subsequent record: 1992
Number of subsequent records: many
Latest record: 2018

Galerucella (Neogalerucella) calmariensis
(Linnaeus, 1767)
Pearce: 4 localities
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2016

Galerucella (Neogalerucella) lineola
(Fabricius, 1781)
Pearce: 6 localities
First subsequent record: 1962
Number of subsequent records: many
Latest record: 2021

Galerucella (Neogalerucella) pusilla
(Duftschmid, 1825)
Pearce: no records
First record: 2002
Number of records: 7
Latest record: 2021

Galerucella (Neogalerucella) tenella
(Linnaeus, 1760)
Pearce: 5 localities
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2020

Lochmaea caprea (Linnaeus, 1758)
Pearce: 7 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Lochmaea crataegi (Forster, 1771)

Pearce: generally common
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2021

Lochmaea suturalis (Thomson, C.G., 1866)

Pearce: generally common
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2020

Pyrrhalta viburni (Paykull, 1799)

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: 8
Latest record: 2019

Mniophila muscorum (Koch, J.D.W., 1803)

Pearce: 1 locality
First subsequent record: 1995
Number of subsequent records: 1
Latest record: 1995

Sphaeroderma rubidum (Graëlls, 1858)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Sphaeroderma testaceum (Fabricius, 1775)

Pearce: 7 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2019

Apteropeda globosa (Illiger, 1794)

Pearce: no records
First record: 1982
Number of records: 7
Latest record: 2016

Apteropeda orbiculata (Marshall, 1802)

Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Podagrica fuscicornis (Linnaeus, 1767)

Pearce: no records
First record: 2005
Number of records: 5
Latest record: 2007

Podagrica fuscipes (Fabricius, 1775)

Pearce: 2 localities
First subsequent record: 2006
Number of subsequent records: 7
Latest record: 2018

Mantura matthewsii (Curtis, 1833)

Pearce: 1 locality
First subsequent record: 1955
Number of subsequent records: 9
Latest record: 2020

Mantura obtusata (Gyllenhal, 1813)

Pearce: 1 locality
First subsequent record: 2005
Number of subsequent records: 2
Latest record: 2007

Mantura rustica (Linnaeus, 1767)

Pearce: 3 localities
First subsequent record: 1982
Number of subsequent records: 2
Latest record: 1983

Crepidodera aurata (Marshall, 1802)

Pearce: 9 localities
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Crepidodera aurea (Geoffroy in Fourcroy, 1785)

Pearce: 5 localities
First subsequent record: 1938
Number of subsequent records: many
Latest record: 2021

Crepidodera fulvicornis (Fabricius, 1792)

Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Crepidodera nitidula (Linnaeus, 1758)

Pearce: no records
First record: 1981
Number of records: 2
Latest record: 1983

Crepidodera plutus (Latreille, 1804)

Pearce: 1 locality
First subsequent record: 1994
Number of subsequent records: 9
Latest record: 2021

Chaetocnema (Chaetocnema) arida

Foudras, 1860

Pearce: no records

First record: 1963

Number of records: many

Latest record: 2020

Chaetocnema (Chaetocnema) confusa

(Boheman, 1851)

Pearce: 1 locality

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2021

Chaetocnema (Chaetocnema) hortensis

(Geoffroy in Fourcroy, 1785)

Pearce: 9 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2021

Chaetocnema (Chaetocnema) sahlbergii

(Gyllenhal, 1827)

Pearce: 1 locality (1887)

First subsequent record: 2015

Number of subsequent records: 3

Latest record: 2015

Chaetocnema (Chaetocnema) subcoerulea

(Kutschera, 1864)

Pearce: 3 localities

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2019

Chaetocnema (Tlanoma) concinna

(Marsham, 1802)

Pearce: generally common and widely distributed
(but probably includes *picipes*)

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2021

Chaetocnema (Tlanoma) picipes Stephens,

1831

Pearce: no records (but see *concinna* above)

First record: 2010

Number of records: many

Latest record: 2021

Psylliodes affinis (Paykull, 1799)

Pearce: 3 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2021

Psylliodes chalconera (Illiger, 1807)

Pearce: 1 locality (1904)

First subsequent record: 1980

Number of subsequent records: 4

Latest record: 2018

Psylliodes chrysocephala (Linnaeus, 1758)

Pearce: 1 locality (1897)

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2021

Psylliodes cuprea (Koch, J.D.W., 1803)

Pearce: 3 localities

First subsequent record: 1995

Number of subsequent records: 4

Latest record: 2014

Psylliodes dulcamarae (Koch, J.D.W., 1803)

Pearce: 1 locality

First subsequent record: 1953

Number of subsequent records: 8

Latest record: 2008

Psylliodes laticollis Kutschera, 1860

Pearce: no records

First record: 1987

Number of records: 5

Latest record: 2019

Psylliodes luteola (Müller, O.F., 1776)

Pearce: 2 localities

First subsequent record: 1993

Number of subsequent records: many

Latest record: 2021

Psylliodes marcida (Illiger, 1807)

Pearce: 1 locality

No subsequent records

Psylliodes napi (Fabricius, 1792)

Pearce: 4 localities

First subsequent record: 1978

Number of subsequent records: many

Latest record: 2021

Psylliodes picina (Marsham, 1802)

Pearce: 2 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Altica brevicollis Foudras, 1860

Pearce: generally common
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2021

Altica helianthemis (Allard, 1859)

Pearce: 3 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2021

Altica longicollis (Allard, 1860)

Pearce: 3 localities
First subsequent record: 1999
Number of subsequent records: 9
Latest record: 2019

Altica lythri Aubé, 1843

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Altica oleracea (Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Altica palustris Weise, 1888

Pearce: 1 locality
First subsequent record: 1933
Number of subsequent records: many
Latest record: 2021

Aphthona atratula Allard, 1859

Pearce: 1 locality
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2016

Aphthona atrocaerulea (Stephens, 1831)

Pearce: 3 localities
First subsequent record: 1994
Number of subsequent records: many
Latest record: 2018

Aphthona euphorbiae (Schränk, 1781)

Pearce: 3 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2020

Aphthona herbigrada (Curtis, 1837)

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Aphthona lutescens (Gyllenhal, 1813)

Pearce: 6 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Aphthona melancholica Weise, 1888

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Aphthona nigriceps (Redtenbacher, W., 1842)

Pearce: 1 locality
First subsequent record: 2020
Number of subsequent records: 1
Latest record: 2020

Aphthona nonstriata (Goeze, 1777)

Pearce: 4 localities
First subsequent record: 1948
Number of subsequent records: many
Latest record: 2021

Batophila aerata (Marsham, 1802)

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Batophila rubi (Paykull, 1799)

Pearce: 3 localities
First subsequent record: 2011
Number of subsequent records: 4
Latest record: 2015

Hermaphysalis mercurialis (Fabricius, 1792)

Pearce: 4 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2021

Lythraea salicariae (Paykull, 1800)

Pearce: no records
First record: 1981
Number of records: 3
Latest record: 2009

Ochrosis ventralis (Illiger, 1807)

Pearce: 2 localities

First subsequent record: 2017

Number of subsequent records: 2

Latest record: 2020

Derocrepis rufipes (Linnaeus, 1758)

Pearce: 6 localities

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2021

Neocrepidodera ferruginea (Scopoli, 1763)

Pearce: generally common and widely distributed

First subsequent record: 1968

Number of subsequent records: many

Latest record: 2020

Neocrepidodera impressa (Fabricius, 1801)

Pearce: 1 locality

First subsequent record: 1937

Number of subsequent records: many

Latest record: 2018

Neocrepidodera transversa (Marsham, 1802)

Pearce: generally common and widely distributed

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2021

Hippuriphila modeeri (Linnaeus, 1760)

Pearce: 5 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2020

Longitarsus (Longitarsus) aeneicollis (Faldermann, 1837)

Pearce: 2 localities

First subsequent record: 1995

Number of subsequent records: 2

Latest record: 2008

Longitarsus (Longitarsus) aeruginosus (Foudras, 1860)

Pearce: no records

First record: 1925

Number of records: 1

Latest record: 1925

Longitarsus (Longitarsus) agilis (Rye, 1868)

Pearce: no records

First record: 2009

Number of records: 1

Latest record: 2009

Longitarsus (Longitarsus) atricillus (Linnaeus, 1760)

Pearce: 7 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2020

Longitarsus (Longitarsus) ballotae (Marsham, 1802)

Pearce: 2 localities

First subsequent record: 1951

Number of subsequent records: 4

Latest record: 2020

Longitarsus (Longitarsus) brunneus (Duftschmid, 1825)

Pearce: 3 localities

No subsequent records

Longitarsus (Longitarsus) dorsalis (Fabricius, 1781)

Pearce: 2 localities

First subsequent record: 1982

Number of subsequent records: many

Latest record: 2019

Longitarsus (Longitarsus) exsoletus (Linnaeus, 1758)

Pearce: 5 localities

First subsequent record: 1966

Number of subsequent records: many

Latest record: 2021

Longitarsus (Longitarsus) ferrugineus (Foudras, 1860)

Pearce: 1 locality

No subsequent records

Longitarsus (Longitarsus) flavicornis (Stephens, 1831)

Pearce: 1 locality (but see *jacobaeae*)

First subsequent record: 1933

Number of subsequent records: many

Latest record: 2020

Longitarsus (Longitarsus) gracilis

(Kutschera, 1864)

Pearce: 6 localities

First subsequent record: 1982

Number of subsequent records: many

Latest record: 2020

Longitarsus (Longitarsus) holsaticus

(Linnaeus, 1758)

Pearce: 1 locality (1916)

First subsequent record: 1987

Number of subsequent records: 1

Latest record: 1987

Longitarsus (Longitarsus) jacobaeae

(Waterhouse, G.R., 1858)

Pearce: 6 localities (but probably includes *flavicornis*)

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2019

Longitarsus (Longitarsus) kutscherae

(Rye, 1872)

Pearce: 1 locality

No subsequent records

Longitarsus (Longitarsus) luridus

(Scopoli, 1763)

Pearce: generally common

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2020

Longitarsus (Longitarsus) lycopi (Foudras, 1860)

Pearce: 3 localities

First subsequent record: 1995

Number of subsequent records: 6

Latest record: 2020

Longitarsus (Longitarsus) melanocephalus

(De Geer, 1775)

Pearce: 8 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2021

Longitarsus (Longitarsus) membranaceus

(Foudras, 1860)

Pearce: 4 localities

First subsequent record: 2001

Number of subsequent records: 8

Latest record: 2020

Longitarsus (Longitarsus) minusculus

(Foudras, 1860)

Pearce: no records

First record: 2002

Number of records: 6

Latest record: 2020

Longitarsus (Longitarsus) nasturtii

(Fabricius, 1792)

Pearce: 2 localities

First subsequent record: 2019

Number of subsequent records: 1

Latest record: 2019

Longitarsus (Longitarsus) nigerrimus

(Gyllenhal, 1827)

Pearce: 1 locality

No subsequent records

Longitarsus (Longitarsus) nigrofasciatus

(Goeze, 1777)

Pearce: no records

First record: 2006

Number of records: 3

Latest record: 2019

Longitarsus (Longitarsus) obliteratedus

(Rosenhauer, 1847)

Pearce: no records

First record: 1983

Number of records: many

Latest record: 2021

Longitarsus (Longitarsus) ochroleucus

(Marsham, 1802)

Pearce: 4 localities

First record: 1954

Number of records: 1

Latest record: 1954

Longitarsus (Longitarsus) parvulus

(Paykull, 1799)

Pearce: no records

First record: 1993

Number of records: many

Latest record: 2021

Longitarsus (Longitarsus) pellucidus

(Foudras, 1860)

Pearce: no records

First record: 1978

Number of records: 3

Latest record: 2018

Longitarsus (Longitarsus) pratensis
(Panzer, 1794)
Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Longitarsus (Longitarsus) quadriguttatus
(Pontoppidan, 1763)
Pearce: no records
First record: 1987
Number of records: 1
Latest record: 1987

Longitarsus (Longitarsus) rubiginosus
(Foudras, 1860)
Pearce: no records
First record: 1975
Number of records: many
Latest record: 2019

Longitarsus (Longitarsus) rutilus (Illiger, 1807)
Pearce: no records
First record: 1983
Number of records: 3
Latest record: 2013

Longitarsus (Longitarsus) strigicollis
Wollaston, 1864 [2012: *Longitarsus fowleri*]
Pearce: no records
First record: 1998
Number of records: many
Latest record: 2017

Longitarsus (Longitarsus) succineus
(Foudras, 1860)
Pearce: 4 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2019

Longitarsus (Longitarsus) suturellus
(Duftschmid, 1825)
Pearce: 5 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Longitarsus (Longitarsus) tabidus
(Fabricius, 1775)
Pearce: 3 localities
First subsequent record: 1982
Number of subsequent records: 9
Latest record: 2011

Longitarsus (Testergus) anchusae
(Paykull, 1799)
Pearce: 4 localities
First subsequent record: 1996
Number of subsequent records: 7
Latest record: 2019

Epitrix atropae Foudras, 1860
Pearce: no records
First record: 2006
Number of records: 2
Latest record: 2009

Epitrix pubescens (Koch, J.D.W., 1803)
Pearce: no records
First record: 1993
Number of records: many
Latest record: 2021

Phyllotreta astrachanica Lopatin, 1977
[2012: *Phyllotreta diademata*]
Pearce: no records
First record: 1926
Number of records: 5
Latest record: 2021

Phyllotreta atra (Fabricius, 1775)
Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Phyllotreta consobrina (Curtis, 1837)
Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: 6
Latest record: 2020

Phyllotreta cruciferae (Goeze, 1777)
Pearce: 7 localities
First subsequent record: 1981
Number of subsequent records: 3
Latest record: 2012

Phyllotreta exclamationis (Thunberg, 1784)

Pearce: 3 localities

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2019

Phyllotreta flexuosa (Illiger, 1794)

Pearce: no records

First record: 2008

Number of records: 2

Latest record: 2020

Phyllotreta nemorum (Linnaeus, 1758)

Pearce: 5 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Phyllotreta nigripes (Fabricius, 1775)

Pearce: 5 localities

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2021

Phyllotreta nodicornis (Marsham, 1802)

Pearce: 4 localities

First subsequent record: 1980

Number of subsequent records: 7

Latest record: 1988

Phyllotreta ochripes (Curtis, 1837)

Pearce: no records

First record: 1987

Number of records: 6

Latest record: 2020

Phyllotreta punctulata (Marsham, 1802)

Pearce: 1 locality

First subsequent record: 1969

Number of subsequent records: 1

Latest record: 1969

Phyllotreta tetrastigma (Comolli, 1837)

Pearce: 1 locality

First subsequent record: 1987

Number of subsequent records: many

Latest record: 2016

Phyllotreta undulata (Kutschera, 1860)

Pearce: 9 localities

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2021

Phyllotreta vittula (Redtenbacher, 1849)

Pearce: 3 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

93. Family NEMONYCHIDAE Bedel, 1882

Cimberis attelaboides (Fabricius, 1787)

Pearce: 1 locality

First subsequent record: 1980

Number of subsequent records: 3

Latest record: 2009

94. Family ANTHRIBIDAE Billberg, 1820

Anthribus fasciatus Forster, 1770

Pearce: 1 locality

First subsequent record: 1986

Number of subsequent records: 4

Latest record: 1997

Anthribus nebulosus Forster, 1770

Pearce: "Dorset" (A. Ford)

First subsequent record: 1986

Number of subsequent records: 5

Latest record: 2010

Platyrhinus resinosus (Scopoli, 1763)

Pearce: no records

First record: 2005

Number of records: 4

Latest record: 2016

Platystomos albinus (Linnaeus, 1758)

Pearce: 2 localities

First subsequent record: 1980

Number of subsequent records: 9

Latest record: 2018

Dissoleucas niveirostris (Fabricius, 1798)

Pearce: 1 locality (1893)

First subsequent record: 1984

Number of subsequent records: 2

Latest record: 1984

95. Family RHYNCHITIDAE Gistel, 1848

Involvulus icosandriae (Scopoli, 1763)
[2012: *Involvulus caeruleus*]

Pearce: 1 locality
First subsequent record: 1983
Number of subsequent records: 4
Latest record: 2001

Lasiorrhynchites cavifrons (Gyllenhal, 1833)

Pearce: 3 localities
First subsequent record: 1995
Number of subsequent records: 3
Latest record: 2014

Lasiorrhynchites comatus (Gyllenhal, 1833) [2012: *Lasiorrhynchites olivaceus*]

Pearce: 1 locality
First subsequent record: 1994
Number of subsequent records: 1
Latest record: 1994

Neocoenorrhinus (*Neocoenorrhinidius*) *interpunctatus* (Stephens, 1831)

Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: 3
Latest record: 1984

Neocoenorrhinus (*Neocoenorrhinus*) *germanicus* (Herbst, 1797)

Pearce: 4 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2019

Neocoenorrhinus (*Schoenitemnus*) *minutus* (Herbst, 1797)

Pearce: 4 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2019

Tatianaerhynchites aequatus (Linnaeus, 1767)

Pearce: 4 localities
First subsequent record: 1939
Number of subsequent records: many
Latest record: 2021

Temnocerus coeruleus (Fabricius, 1798)

Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: 8
Latest record: 2020

Temnocerus longiceps (Thomson, C.G., 1888)

Pearce: no records
First record: 1982
Number of records: 3
Latest record: 1999

Temnocerus nanus (Paykull, 1792)

Pearce: 2 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2019

Bytiscus betulae (Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2021

Caenorhinus mannerheimii (Hummel, 1823)

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2013

Deporaus betulae (Linnaeus, 1758)

Pearce: generally common
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2018

96. Family ATTELABIDAE Billberg, 1820

Apoderus coryli (Linnaeus, 1758)

Pearce: generally common
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Attelabus nitens (Scopoli, 1763)

Pearce: 6 localities
First subsequent record: 1981
Number of subsequent records: 1
Latest record: 1981

97. Family APIONIDAE Schönherr, 1823

Apion cruentatum Walton, 1844

Pearce: 2 localities
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2018

Apion frumentarium (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Apion haematodes Kirby, 1808

Pearce: 8 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Apion rubens Walton, 1837

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Apion rubiginosum Walton, 1837

Pearce: no records
First record: 1996
Number of records: 5
Latest record: 2015

Aizobius sedi (Germar, 1818)

Pearce: no records
First record: 1937
Number of records: 1
Latest record: 1937

Perapion affine (Kirby, 1808)

Pearce: no records
First record: 1953
Number of records: 4
Latest record: 2008

Perapion curtirostre (Germar, 1817)

Pearce: 7 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2020

Perapion hydrolapathi (Marsham, 1802)

Pearce: 2 localities
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2020

Perapion marchicum (Herbst, 1797)

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Perapion violaceum (Kirby, 1808)

Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2019

Pseudaplemonus limonii (Kirby, 1808)

Pearce: no records
First record: 1989
Number of records: 3
Latest record: 2018

Aspidapion (Aspidapion) radiolus (Marsham, 1802)

Pearce: 4 localities
First subsequent record: 1966
Number of subsequent records: many
Latest record: 2020

Aspidapion (Koestlinia) aeneum (Fabricius, 1775)

Pearce: 3 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2019

Acentrotypus brunnipes (Boheman, 1839)

Pearce: no records
First record: 1937
Number of records: 1
Latest record: 1937

Ceratapion (Acanephodus) onopordi (Kirby, 1808)

Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Ceratapion (Ceratapion) carduorum (Kirby, 1808)

Pearce: 8 localities, but many of these could refer to *gibbirostre*
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2018

Ceratapion (Ceratapion) gibbistrostre (Gyllenhal, 1813)

Pearce: no records, but see *carduorum*

First record: 1963

Number of records: many

Latest record: 2021

Diplapion confluens (Kirby, 1808)

Pearce: 3 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2020

Diplapion stolidum (Germar, 1817)

Pearce: no records

First record: 1968

Number of records: 7

Latest record: 1998

Omphalapion hookerorum (Kirby, 1808)

Pearce: 3 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2021

Exapion difficile (Herbst, 1797)

Pearce: no records

First record: 1981

Number of records: 5

Latest record: 2019

Exapion fuscirostre (Fabricius, 1775)

Pearce: 1 locality (1895)

First subsequent record: 2008

Number of subsequent records: 2

Latest record: 2008

Exapion genistae (Kirby, 1811)

Pearce: 2 localities

First subsequent record: 1981

Number of subsequent records: 1

Latest record: 1981

Exapion ulicis (Forster, 1771)

Pearce: generally common and widely distributed

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2021

Kalcapion semivittatum (Gyllenhal, 1833)

Pearce: no records

First record: 1980

Number of records: many

Latest record: 2019

Melanapion minimum (Herbst, 1797)

Pearce: 2 localities (1896, 1897)

No subsequent records

Squamapion atomarium (Kirby, 1808)

Pearce: no records

First record: 1941

Number of records: many

Latest record: 2015

Squamapion cineraceum (Wencker, 1864)

Pearce: 1 locality (1904)

First subsequent record: 2002

Number of subsequent records: 3

Latest record: 2021

Squamapion vicinum (Kirby, 1808)

Pearce: "Dorset" (A. Ford)

First subsequent record: 1984

Number of subsequent records: 1

Latest record: 1984

Taeniapion urticarium (Herbst, 1784)

Pearce: 4 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2020

Malvapion malvae (Fabricius, 1775)

Pearce: no records

First record: 1952

Number of records: many

Latest record: 2020

Pseudapion rufirostre (Fabricius, 1775)

Pearce: 1 locality

First subsequent record: 1976

Number of subsequent records: many

Latest record: 2020

Catapion curtisii (Stephens, 1831)

Pearce: 1 locality

First subsequent record: 1975

Number of subsequent records: 3

Latest record: 1983

Catapion pubescens (Kirby, 1811)

Pearce: 2 localities

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2018

Catapion seniculus (Kirby, 1808)

Pearce: 4 localities

First subsequent record: 1968

Number of subsequent records: many

Latest record: 2021

Cyanapion (Bothryorrhynchapion) afer
(Gyllenhal, 1833)

Pearce: 1 locality

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2018

Cyanapion (Bothryorrhynchapion)
gyllenhalii (Kirby, 1808)

Pearce: 2 localities

No subsequent records

Cyanapion (Cyanapion) spencii (Kirby,
1808)

Pearce: 1 locality

First subsequent record: 1954

Number of subsequent records: 9

Latest record: 2018

Eutrichapion (Cnemapion) vorax (Herbst,
1797)

Pearce: 1 locality

First subsequent record: 2010

Number of subsequent records: 1

Latest record: 2010

Eutrichapion (Eutrichapion) ervi (Kirby,
1808)

Pearce: 3 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2021

Eutrichapion (Eutrichapion) viciae
(Paykull, 1800)

Pearce: 1 locality

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2020

Eutrichapion (Psilocalymma) punctiger
(Paykull, 1792)

Pearce: "Dorset" (A. Ford)

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2016

Hemitrichapion reflexum (Gyllenhal,
1833)

Pearce: 1 locality (1893)

First subsequent record: 1954

Number of subsequent records: 4

Latest record: 1994

Hemitrichapion waltoni (Stephens, 1839)

Pearce: 2 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2020

Holotrichapion (Apiops) pisi (Fabricius,
1801)

Pearce: 4 localities

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2018

Holotrichapion (Holotrichapion) ononis
(Kirby, 1808)

Pearce: 1 locality

First subsequent record: 1966

Number of subsequent records: many

Latest record: 2020

Holotrichapion (Legaricapion) aethiops
(Herbst, 1797)

Pearce: 3 localities

First subsequent record: 1969

Number of subsequent records: many

Latest record: 2016

Ischnopterapion (Chlorapion) virens
(Herbst, 1797)

Pearce: 5 localities

First subsequent record: 1968

Number of subsequent records: many

Latest record: 2021

Ischnopterapion (Ischnopterapion) loti
(Kirby, 1808)

Pearce: 4 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2021

Ischnopterapion (Ischnopterapion)
modestum (Germar, 1817)

Pearce: no records

First record: 1980

Number of records: many

Latest record: 2020

Oxystoma cerdo (Gerstaecker, 1854)

Pearce: no records
First record: 2004
Number of records: 4
Latest record: 2020

Oxystoma cracca (Linnaeus, 1767)

Pearce: 1 locality
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2015

Oxystoma pomonae (Fabricius, 1798)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Oxystoma subulatum (Kirby, 1808)

Pearce: 2 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2018

Pirapion immune (Kirby, 1808)

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: 8
Latest record: 2019

Protopirapion atratum (Germar, 1817)

Pearce: 3 localities
First subsequent record: 1927
Number of subsequent records: many
Latest record: 2021

Stenopterapion (Cobosiotherium) scutellare (Kirby, 1811)

Pearce: 3 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2018

Stenopterapion (Stenopterapion) meliloti (Kirby, 1808)

Pearce: no records
First record: 1966
Number of records: 7
Latest record: 2018

Stenopterapion (Stenopterapion) tenue (Kirby, 1808)

Pearce: 2 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Synapion ebeninum (Kirby, 1808)

Pearce: 1 locality
First subsequent record: 1927
Number of subsequent records: many
Latest record: 2020

Betulapion simile (Kirby, 1811)

Pearce: no records
First record: 1985
Number of records: many
Latest record: 2020

Protapion apricans (Herbst, 1797)

Pearce: generally common
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2021

Protapion assimile (Kirby, 1808)

Pearce: 1 locality
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Protapion difforme (Germar, 1818)

Pearce: 4 localities
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2019

Protapion dissimile (Germar, 1817)

Pearce: 1 locality
First subsequent record: 2016
Number of subsequent records: 2
Latest record: 2016

Protapion filirostre (Kirby, 1808)

Pearce: no records
First record: 1966
Number of records: many
Latest record: 2019

Protapion fulvipes (Geoffroy in Fourcroy, 1785)

Pearce: 7 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2021

Protapion laevicolle (Kirby, 1811)

Pearce: 1 locality
First subsequent record: 1999
Number of subsequent records: many
Latest record: 2021

Protapion nigritarse (Kirby, 1808)

Pearce: 5 localities

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2019

Protapion ononidis (Gyllenhal, 1827)

Pearce: no records

First record: 1954

Number of records: many

Latest record: 2020

Protapion trifolii (Linnaeus, 1768)

Pearce: 6 localities

First subsequent record: 1939

Number of subsequent records: many

Latest record: 2021

Protapion varipes (Germar, 1817)

Pearce: 3 localities

First subsequent record: 2000

Number of subsequent records: 2

Latest record: 2000

98. Family NANOPHYIDAE Gistel, 1848

Dieckmanniellus gracilis (Redtenbacher, 1849)

Pearce: "Dorset" (A. Ford)

First subsequent record: 1964

Number of subsequent records: 4

Latest record: 2020

Nanophyes marmoratus (Goeze, 1777)

Pearce: 5 localities

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2019

99. Family DRYOPHTHORIDAE Schönherr, 1825

Sitophilus granarius (Linnaeus, 1758)

Pearce: 4 localities

No subsequent records

Sitophilus oryzae (Linnaeus, 1763)

Pearce: 4 localities

First subsequent record: 1999

Number of subsequent records: 1

Latest record: 1999

100. Family ERIRHINIDAE Schönherr, 1825

Grypus equiseti (Fabricius, 1775)

Pearce: 5 localities

First subsequent record: 1980

Number of subsequent records: many

Latest record: 2016

Notaris acridulus (Linnaeus, 1758)

Pearce: 4 localities

First subsequent record: 1978

Number of subsequent records: many

Latest record: 2021

Notaris scirpi (Fabricius, 1792)

Pearce: 2 localities

First subsequent record: 1938

Number of subsequent records: 9

Latest record: 2017

Thryogenes festucae (Herbst, 1795)

Pearce: 1 locality (1895)

First subsequent record: 1994

Number of subsequent records: 2

Latest record: 2019

Thryogenes fiorii Zumpt, 1928

Pearce: no records

First record: 2005

Number of records: 1

Latest record: 2005

Thryogenes nereis (Paykull, 1800)

Pearce: 2 localities

First subsequent record: 1992

Number of subsequent records: many

Latest record: 2021

Thryogenes scirrhosus (Gyllenhal, 1836)

Pearce: no records

First record: 2006

Number of records: 1

Latest record: 2006

Tournotaris bimaculata (Fabricius, 1787)

Pearce: 1 locality (1923)

First subsequent record: 2015

Number of subsequent records: 1

Latest record: 2015

Tanysphyrus lemnae (Paykull, 1792)

Pearce: 1 locality

First subsequent record: 1954

Number of subsequent records: 6

Latest record: 2019

101. Family RAYMONDIONYMIDAE
Reitter, 1913

Ferreria marqueti (Aubé, 1863)

Pearce: no records
First record: 2021
Number of records: 2
Latest record: 2021

102. Family CURCULIONIDAE
Latreille, 1802

Limnobaris dolorosa (Goeze, 1777)

Pearce: no records (but see *t-album*)
First record: 1981
Number of records: 7
Latest record: 2020

Limnobaris t-album (Linnaeus, 1758)

Pearce: 5 localities (but may include *dolorosa*)
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2020

Aulacobaris lepidii (Germar, 1824)

Pearce: 1 locality
First subsequent record: 2017
Number of subsequent records: 4
Latest record: 2021

Aulacobaris picicornis (Marsham, 1802)

Pearce: 1 locality
No subsequent records

Baris analis (Olivier, 1791)

Pearce: no records
First record: 1989
Number of records: many
Latest record: 2018

Melanobaris laticollis (Marsham, 1802)

Pearce: 1 locality (1886)
First subsequent record: 1981
Number of subsequent records: 2
Latest record: 1982

Acalyptus carpini (Fabricius, 1792)

Pearce: 2 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2020

Anoplus plantaris (Naezen, 1794)

Pearce: 2 localities (1911, 1919)
First subsequent record: 1974
Number of subsequent records: many
Latest record: 2015

Anoplus roboris Suffrian, 1840

Pearce: no records
First record: 1997
Number of records: 3
Latest record: 2001

Anthonomus bituberculatus Thomson, C.G., 1868

Pearce: no records
First record: 1977
Number of records: many
Latest record: 2020

Anthonomus brunnipennis Curtis, 1840

Pearce: 1 locality (1925)
First subsequent record: 1995
Number of subsequent records: 3
Latest record: 2014

Anthonomus chevrolati Desbrochers des Loges, 1868

Pearce: 1 locality
No subsequent records

Anthonomus pedicularius (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2019

Anthonomus pomorum (Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: 7
Latest record: 2009

Anthonomus rubi (Herbst, 1795)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Anthonomus rufus Gyllenhal, 1835

Pearce: no records
First record: 1986
Number of records: 2
Latest record: 1987

Anthonomus ulmi (De Geer, 1775)

Pearce: 3 localities
First subsequent record: 1981
Number of subsequent records: 9
Latest record: 2021

Cionus alauda (Herbst, 1784)

Pearce: 10 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Cionus hortulanus (Geoffroy in Fourcroy, 1785)

Pearce: 3 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Cionus nigratarsis Reitter, 1904

Pearce: 1 locality
No subsequent records

Cionus scrophulariae (Linnaeus, 1758)

Pearce: common
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Cionus tuberculosus (Scopoli, 1763)

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2019

Cleopus pulchellus (Herbst, 1795)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2006

Archarius pyrrhoceras (Marshall, 1802)

Pearce: 6 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Archarius salicivorus (Paykull, 1792)

Pearce: 6 localities
First subsequent record: 1967
Number of subsequent records: many
Latest record: 2021

Curculio betulae (Stephens, 1831)

Pearce: 2 localities (1881, 1894)
First subsequent record: 1993
Number of subsequent records: 4
Latest record: 2007

Curculio glandium Marshall, 1802

Pearce: 5 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2021

Curculio nucum Linnaeus, 1758

Pearce: 5 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2016

Curculio rubidus (Gyllenhal, 1835)

Pearce: 1 locality
First subsequent record: 2001
Number of subsequent records: 4
Latest record: 2010

Curculio venosus (Gravenhorst, 1807)

Pearce: 6 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2020

Curculio villosus Fabricius, 1781

Pearce: 5 localities
First subsequent record: 1981
Number of subsequent records: 5
Latest record: 1986

Dorytomus dejeani Faust, 1883

Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2020

Dorytomus filirostris (Gyllenhal, 1835)

Pearce: no records
First record: 2020
Number of records: 4
Latest record: 2021

Dorytomus longimanus (Forster, 1771)

Pearce: 1 locality
First subsequent record: 1983
Number of subsequent records: 7
Latest record: 2020

Dorytomus melanophthalmus (Paykull, 1792)

Pearce: 1 locality
First subsequent record: 1983
Number of subsequent records: 9
Latest record: 2020

Dorytomus rufatus (Bedel, 1888)

Pearce: 3 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Dorytomus taeniatus (Fabricius, 1781)

Pearce: 4 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2020

Dorytomus tortrix (Linnaeus, 1760)

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: 8
Latest record: 2015

Dorytomus tremulae (Fabricius, 1787)

Pearce: 1 locality
First subsequent record: 2010
Number of subsequent records: 1
Latest record: 2010

Ellescus bipunctatus (Linnaeus, 1758)

Pearce: 3 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2017

Cleopomiarus graminis (Gyllenhal, 1813)

Pearce: 1 locality
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2019

Cleopomiarus plantarum (Germar, 1824)

Pearce: 1 locality
First subsequent record: 1953
Number of subsequent records: 1
Latest record: 1953

Gymnetron beccabungae (Linnaeus, 1760)

Pearce: 1 locality
First subsequent record: 1989
Number of subsequent records: 2
Latest record: 2013

Gymnetron melanarium (Germar, 1821)

Pearce: 1 locality (1893)
First subsequent record: 1982
Number of subsequent records: 8
Latest record: 2021

Gymnetron veronicae (Germar, 1821)

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2020

Gymnetron villosulum Gyllenhal, 1838

Pearce: 3 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2006

Mecinus circulatus (Marshall, 1802)

Pearce: 5 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2019

Mecinus collaris Germar, 1821

Pearce: no records
First record: 2014
Number of records: 4
Latest record: 2015

Mecinus labilis (Herbst, 1795)

Pearce: 1 locality
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Mecinus pascuorum (Gyllenhal, 1813)

Pearce: 2 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Mecinus pyrae (Herbst, 1795)

Pearce: 8 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Miarus campanulae (Linnaeus, 1767)

Pearce: no records
First record: 1955
Number of records: many
Latest record: 2019

Rhinusa antirrhini (Paykull, 1800)

Pearce: 4 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2021

Rhinusa collina (Gyllenhal, 1813)

Pearce: 2 localities
No subsequent records

Isochnus sequensi (Stierlin, 1894)

Pearce: no records
First record: 2016
Number of records: 4
Latest record: 2017

Orchestes (Alyctus) rusci (Herbst, 1795)

Pearce: 2 localities
First subsequent record: 1967
Number of subsequent records: many
Latest record: 2020

Orchestes (Alyctus) testaceus (Müller, O.F., 1776)

Pearce: 1 locality
No subsequent records

Orchestes (Orchestes) alni (Linnaeus, 1758)

Pearce: 7 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2017

Orchestes (Orchestes) fagi (Linnaeus, 1758)

Pearce: 4 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2014

Orchestes (Orchestes) hortorum (Fabricius, 1792) [2012: *Orchestes signifer*]

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2018

Orchestes (Orchestes) jota (Fabricius, 1787)

Pearce: 2 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2021

Orchestes (Orchestes) pilosus (Fabricius, 1781)

Pearce: 4 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Orchestes (Orchestes) quercus (Linnaeus, 1758)

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2021

Pseudorchestes pratensis (Germar, 1821)

Pearce: 1 locality
No subsequent records

Rhamphus oxyacanthae (Marsham, 1802)

Pearce: no records
First record: 1980
Number of records: many
Latest record: 2019

Rhamphus pulicarius (Herbst, 1795)

Pearce: 4 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2020

Tachyerges salicis (Linnaeus, 1758)

Pearce: 5 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2017

Tachyerges stigma (Germar, 1821)

Pearce: 2 localities
First subsequent record: 1965
Number of subsequent records: many
Latest record: 2020

Smicronyx coecus (Reich, 1797)

Pearce: 1 locality
First subsequent record: 2015
Number of subsequent records: 2
Latest record: 2019

Smicronyx jungermanniae (Reich, 1797)

Pearce: 2 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2020

Smicronyx reichii (Gyllenhal, 1835)

Pearce: 1 locality
First subsequent record: 2000
Number of subsequent records: 6
Latest record: 2018

Pachytychius haematocephalus
(Gyllenhal, 1835)

Pearce: "Dorset" (A. Ford)
First subsequent record: 1947
Number of subsequent records: 2
Latest record: 2018

Orthochaetes insignis (Aubé, 1863)

Pearce: 2 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2018

Orthochaetes setiger (Beck, 1817)

Pearce: 5 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2019

Sibinia (Dichotychius) sodalis Germar,
1824

Pearce: 1 locality
No subsequent records

Sibinia (Sibinia) arenariae Stephens, 1831

Pearce: 3 localities
First subsequent record: 1964
Number of subsequent records: many
Latest record: 2016

Sibinia (Sibinia) primita (Herbst, 1795)

Pearce: 4 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2016

Sibinia (Sibinia) pyrrhodactyla (Marsham,
1802)

Pearce: 2 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2019

Tychius breviusculus Desbrochers des
Loges, 1873

Pearce: no records
First record: 1993
Number of records: many
Latest record: 2019

Tychius crassirostris Kirsch, 1871

Pearce: no records
First record: 1926
Number of records: 9
Latest record: 2020

Tychius junceus (Reich, 1797)

Pearce: 4 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2018

Tychius lineatulus Stephens, 1831

Pearce: 1 locality (1886)
First subsequent record: 1980
Number of subsequent records: 8
Latest record: 2016

Tychius meliloti Stephens, 1831

Pearce: 4 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Tychius picirostris (Fabricius, 1787)

Pearce: 4 localities
First subsequent record: 1966
Number of subsequent records: many
Latest record: 2021

Tychius pusillus Germar, 1842

Pearce: 3 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2021

Tychius schneideri (Herbst, 1795)

Pearce: 2 localities
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2006

Tychius squamulatus Gyllenhal, 1835

Pearce: 4 localities
First subsequent record: 1985
Number of subsequent records: many
Latest record: 2020

Tychius stephensi Schönherr, 1835

Pearce: 3 localities
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2020

Tychius tibialis Boheman, 1843

Pearce: 1 locality
First subsequent record: 1984
Number of subsequent records: many
Latest record: 2021

Bagous frit (Herbst, 1795)

Pearce: 1 locality
No subsequent records

Bagous limosus (Gyllenhal, 1827)

Pearce: no records
First record: 2005
Number of records: 5
Latest record: 2015

Bagous lutulosus (Gyllenhal, 1827)

Pearce: no records
First record: 2009
Number of records: 1
Latest record: 2009

Bagous tempestivus (Herbst, 1795)

Pearce: no records
First record: 1955
Number of records: 9
Latest record: 2018

Hydronomus alismatis (Marshall, 1802)

[2012: *Bagous alismatis*]
Pearce: 2 localities
First subsequent record: 1981
Number of subsequent records: 3
Latest record: 2006

Amalus scortillum (Herbst, 1795)

Pearce: 3 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Amalorrhynchus melanarius (Stephens, 1831)

Pearce: "Dorset" (A. Ford)
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Calosirus terminatus (Herbst, 1795)

Pearce: 3 localities
First subsequent record: 1966
Number of subsequent records: 8
Latest record: 2020

Ceutorhynchus alliariae (Brisout de Barneville, C., 1860)

Pearce: no records
First record: 1983
Number of records: many
Latest record: 2018

Ceutorhynchus assimilis (Paykull, 1792)

Pearce: 6 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2016

Ceutorhynchus atomus Boheman, 1845

Pearce: no records
First record: 2001
Number of records: 2
Latest record: 2012

Ceutorhynchus chalybaeus Germar, 1824

Pearce: 2 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2020

Ceutorhynchus cochleariae (Gyllenhal, 1813)

Pearce: 2 localities
First subsequent record: 1971
Number of subsequent records: many
Latest record: 2017

Ceutorhynchus constrictus (Marshall, 1802)

Pearce: 1 locality
First subsequent record: 1983
Number of subsequent records: 5
Latest record: 2011

Ceutorhynchus contractus (Marshall, 1802)

Pearce: 9 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2019

Ceutorhynchus erysimi (Fabricius, 1787)

Pearce: 4 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Ceutorhynchus obstrictus (Marsham, 1802)

Pearce: 6 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2021

Ceutorhynchus pallidactylus (Marsham, 1802)

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2020

Ceutorhynchus pyrrhorhynchus (Marsham, 1802)

Pearce: 3 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2020

Ceutorhynchus sulcicollis (Paykull, 1800)

Pearce: 1 locality
No subsequent records

Ceutorhynchus turbatus Schultze, 1903

Pearce: no records
First record: 1981
Number of records: 5
Latest record: 2013

Ceutorhynchus typhae (Herbst, 1795)

Pearce: 2 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Coeliodes rana (Fabricius, 1787)

Pearce: 5 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2019

Coeliodes ruber (Marsham, 1802)

Pearce: 5 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2019

Coeliodes transversealbofasciatus (Goeze, 1777)

Pearce: 2 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2016

Coelioidinus rubicundus (Herbst, 1795)

Pearce: 3 localities
First subsequent record: 1965
Number of subsequent records: many
Latest record: 2019

Datonychus melanostictus (Marsham, 1802)

Pearce: 2 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2021

Datonychus urticae (Boheman, 1845)

Pearce: 1 locality
First subsequent record: 1983
Number of subsequent records: 1
Latest record: 1983

Drupenatus nasturtii (Germar, 1824)

Pearce: 1 locality
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2020

Glocianus distinctus (Brisout de Barneville, C., 1870)

Pearce: 1 locality
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2021

Glocianus moelleri (Thomson, C.G., 1868)

Pearce: no records
First record: 2014
Number of records: 1
Latest record: 2014

Glocianus punctiger (Sahlberg, C.R., 1835)

Pearce: no records
First record: 1981
Number of records: many
Latest record: 2021

Hadroplontus litura (Fabricius, 1775)

Pearce: 2 localities
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2018

Hadroplontus trimaculatus (Fabricius, 1775)

Pearce: 2 localities

First subsequent record: 1980

Number of subsequent records: many

Latest record: 2015

Micrelus ericae (Gyllenhal, 1813)

Pearce: 4 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2019

Microplontus campestris (Gyllenhal, 1837)

Pearce: 5 localities

First subsequent record: 1980

Number of subsequent records: 7

Latest record: 2020

Microplontus melanostigma (Marsham, 1802) [2012: *Microplontus rugulosus*]

Pearce: 2 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2019

Microplontus triangulum (Boheman, 1845)

Pearce: no records

First record: 1983

Number of records: 1

Latest record: 1983

Mogulones asperifoliarum (Gyllenhal, 1813)

Pearce: 4 localities

First subsequent record: 1978

Number of subsequent records: many

Latest record: 2015

Mogulones geographicus (Goeze, 1777)

Pearce: 3 localities

First subsequent record: 1980

Number of subsequent records: 6

Latest record: 2011

Nedyus quadrimaculatus (Linnaeus, 1758)

Pearce: generally common

First subsequent record: 1966

Number of subsequent records: many

Latest record: 2021

Parethelcus pollinarius (Forster, 1771)

Pearce: 9 localities

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2021

Poophagus sisymbrii (Fabricius, 1777)

Pearce: 2 localities

First subsequent record: 1978

Number of subsequent records: many

Latest record: 2011

Sirocalodes depressicollis (Gyllenhal, 1813)

Pearce: 1 locality (1919)

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2017

Sirocalodes mixtus (Mulsant & Rey, 1859)

Pearce: no records

First record: 1983

Number of records: 2

Latest record: 2017

Sirocalodes quercicola (Paykull, 1792)

Pearce: no records

First record: 1985

Number of records: 3

Latest record: 1985

Stenocarus ruficornis (Stephens, 1831)

Pearce: 2 localities

First subsequent record: 1982

Number of subsequent records: 8

Latest record: 2021

Thamiocolus viduatus (Gyllenhal, 1813)

Pearce: 1 locality

No subsequent records

Trichosirocalus dawsoni (Brisout de Barneville, C., 1869)

Pearce: 2 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2020

Trichosirocalus horridus (Panzer, 1801)

Pearce: 3 localities

First subsequent record: 1978

Number of subsequent records: 3

Latest record: 2005

Trichosirocalus rufulus (Dufour, 1851)

Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: 8
Latest record: 2013

Trichosirocalus troglodytes (Fabricius, 1787)

Pearce: 8 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Zacladus exiguus (Olivier, 1807)

Pearce: 1 locality
First subsequent record: 1985
Number of subsequent records: 7
Latest record: 2019

Zacladus geranii (Paykull, 1800)

Pearce: 1 locality
First subsequent record: 1963
Number of subsequent records: 2
Latest record: 1981

Mononychus punctumalbum (Herbst, 1784)

Pearce: 2 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2020

Eubrychius velutus (Beck, 1817)

Pearce: 1 locality
No subsequent records

Neophytobius muricatus (Brisout de Barneville, C., 1867)

Pearce: no records
First record: 1983
Number of records: many
Latest record: 2020

Neophytobius quadrinodosus (Gyllenhal, 1813)

Pearce: 1 locality
First subsequent record: 1982
Number of subsequent records: many
Latest record: 2020

Pelenomus canaliculatus (Fåhraeus, 1843)

Pearce: 1 locality (1904)
First subsequent record: 2016
Number of subsequent records: 2
Latest record: 2020

Pelenomus commari (Panzer, 1795) [2012:

Pelenomus comari]
Pearce: no records
First record: 2010
Number of records: 2
Latest record: 2011

Pelenomus olssoni (Israelson, 1972)

Pearce: no records
First record: 2009
Number of records: 5
Latest record: 2020

Pelenomus quadricorniger (Colonnelli, 1986)

Pearce: no records
First record: 2016
Number of records: 2
Latest record: 2019

Pelenomus quadrituberculatus (Fabricius, 1787)

Pearce: 2 localities
First subsequent record: 1978
Number of subsequent records: many
Latest record: 2021

Pelenomus waltoni (Boheman, 1843)

Pearce: no records
First record: 1987
Number of records: many
Latest record: 2020

Phytobius leucogaster (Marsham, 1802)

Pearce: 1 locality
No subsequent records

Rhinoncus bruchoides (Herbst, 1784)

Pearce: 1 locality
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2010

Rhinoncus castor (Fabricius, 1792)

Pearce: 6 localities
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Rhinoncus inconspicuous (Herbst, 1795)

Pearce: 1 locality
First subsequent record: 1954
Number of subsequent records: many
Latest record: 2021

Rhinoncus pericarpus (Linnaeus, 1758)

Pearce: 9 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2021

Rhinoncus perpendicularis (Reich, 1797)

Pearce: 3 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2020

Tapinotus sellatus (Fabricius, 1794)

Pearce: no records
First record: 1981
Number of records: 1
Latest record: 1981

Cossonus parallelepipedus (Herbst, 1795)

Pearce: 1 locality
No subsequent records

Rhopalomesites tardii (Curtis, 1825)

Pearce: no records
First record: 1994
Number of records: 1
Latest record: 1994

Pselactus spadix (Herbst, 1795)

Pearce: no records
First record: 1964
Number of records: 3
Latest record: 2005

Pseudophloeophagus truncorum
(Stephens, 1831) [2012:

Pseudophloeophagus aeneopiceus]
Pearce: no records
First record: 1982
Number of records: 8
Latest record: 2018

Euophryum confine (Broun, 1881)

Pearce: no records
First record: 1980
Number of records: many
Latest record: 2021

Pentarthrum huttoni Wollaston, 1854

Pearce: 1 locality
First subsequent record: 2000
Number of subsequent records: 4
Latest record: 2008

Cryptorhynchus lapathi (Linnaeus, 1758)

Pearce: 3 localities
First subsequent record: 1934
Number of subsequent records: 7
Latest record: 2016

Acalles misellus Boheman, 1844

Pearce: 2 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2021

Acalles ptinoides (Marsham, 1802)

Pearce: 1 locality (1901)
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2020

Kyklioacalles roboris (Curtis, 1835)

Pearce: no records
First record: 1984
Number of records: many
Latest record: 2018

Gronops lunatus (Fabricius, 1775)

Pearce: 3 localities
First subsequent record: 1952
Number of subsequent records: many
Latest record: 2017

Graptus triguttatus (Fabricius, 1775)

Pearce: 1 locality
First subsequent record: 1967
Number of subsequent records: many
Latest record: 2020

Strophosoma (Neliocarus) faber (Herbst, 1784)

Pearce: 2 localities
First subsequent record: 1982
Number of subsequent records: 9
Latest record: 2021

Strophosoma (Neliocarus) nebulosum
(Stephens, 1831)

Pearce: 3 localities
First subsequent record: 1927
Number of subsequent records: many
Latest record: 2021

Strophosoma (Neliocarus) sus (Stephens, 1831)

Pearce: very common on heaths
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2020

Strophosoma (Strophosoma) capitatum
(De Geer, 1775)

Pearce: generally common
First subsequent record: 1954
Number of subsequent records: 4
Latest record: 2014

Strophosoma (Strophosoma) fulvicorne
(Walton, 1846)

Pearce: 2 localities
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2016

Strophosoma (Strophosoma)
melanogrammum (Forster, 1771)

Pearce: generally common
First subsequent record: 1935
Number of subsequent records: many
Latest record: 2021

Attactagenus plumbeus (Marsham, 1802)

Pearce: 8 localities
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2021

Philopodon plagiatum (Schaller, 1783)

Pearce: 7 localities
First subsequent record: 1932
Number of subsequent records: many
Latest record: 2021

Barynotus moerens (Fabricius, 1792)

Pearce: no records
First record: 1983
Number of records: 4
Latest record: 1987

Barynotus obscurus (Fabricius, 1775)

Pearce: generally common and widely distributed
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2019

Otiorhynchus (Aranihus) ligneus (Olivier, 1807)

Pearce: 4 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Otiorhynchus (Choilisanus) raucus
(Fabricius, 1777)

Pearce: 2 localities
No subsequent records

Otiorhynchus (Clypeotiorhynchus)
crataegi Germar, 1824

Pearce: no records
First record: 2014
Number of records: 1
Latest record: 2014

Otiorhynchus (Dorymerus) sulcatus
(Fabricius, 1775)

Pearce: generally common
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2020

Otiorhynchus (Metopiorrhynchus)
singularis (Linnaeus, 1767)

Pearce: generally common and widely distributed
First subsequent record: 1934
Number of subsequent records: many
Latest record: 2021

Otiorhynchus (Otiorhynchus) tenebricosus
(Herbst, 1784) [2012: *Otiorhynchus*
clavipes]

Pearce: generally common and widely distributed
First subsequent record: 1953
Number of subsequent records: many
Latest record: 2021

Otiorhynchus (Pendragon) desertus
Rosenhauer, 1847

Pearce: no records
First record: 1983
Number of records: 3
Latest record: 2004

Otiorhynchus (Pendragon) ovatus
(Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 1980
Number of subsequent records: many
Latest record: 2021

Otiorhynchus (Provadilus) rugifrons
(Gyllenhal, 1813)

Pearce: 6 localities
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2017

Otiorhynchus (Zadrehus) atroapterus (De Geer, 1775)

Pearce: 5 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Otiorhynchus (Zustalestus) rugosostriatus (Goeze, 1777)

Pearce: 8 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2019

Phyllobius (Dieletus) argentatus (Linnaeus, 1758)

Pearce: generally common

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Phyllobius (Metaphyllobius) glaucus (Scopoli, 1763)

Pearce: 5 localities

First subsequent record: 1968

Number of subsequent records: many

Latest record: 2021

Phyllobius (Metaphyllobius) pomaceus Gyllenhal, 1834

Pearce: 7 localities

First subsequent record: 1954

Number of subsequent records: many

Latest record: 2021

Phyllobius (Nemoicus) oblongus (Linnaeus, 1758)

Pearce: 6 localities

First subsequent record: 1980

Number of subsequent records: many

Latest record: 2012

Phyllobius (Parnemoicus) roboretanus Gredler, 1882

Pearce: generally common

First subsequent record: 1962

Number of subsequent records: many

Latest record: 2020

Phyllobius (Parnemoicus) viridicollis (Fabricius, 1792)

Pearce: 2 localities

First subsequent record: 1964

Number of subsequent records: many

Latest record: 2012

Phyllobius (Phyllobius) pyri (Linnaeus, 1758)

Pearce: 3 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2021

Phyllobius (Pterygorrhynchus) maculicornis Germar, 1824

Pearce: 4 localities

First subsequent record: 1980

Number of subsequent records: many

Latest record: 2021

Phyllobius (Subphyllobius) virideaeris (Laicharting, 1781)

Pearce: 4 localities

First subsequent record: 1962

Number of subsequent records: many

Latest record: 2021

Liophloeus tessulatus (Müller, O.F., 1776)

Pearce: 5 localities

First subsequent record: 1962

Number of subsequent records: many

Latest record: 2021

Pachyrhinus lethierryi (Desbrochers des Loges, 1875)

Pearce: no records

First record: 2011

Number of records: 1

Latest record: 2011

Polydrusus (Eudipnus) formosus (Mayer, 1779)

Pearce: 1 locality

First subsequent record: 2001

Number of subsequent records: many

Latest record: 2021

Polydrusus (Eudipnus) mollis (Strøm, 1768)

Pearce: 2 localities

First subsequent record: 2014

Number of subsequent records: 1

Latest record: 2014

Polydrusus (Eurodrusus) cervinus (Linnaeus, 1758)

Pearce: 5 localities

First subsequent record: 1935

Number of subsequent records: many

Latest record: 2021

Polydrusus (Eurodrusus) confluens

Stephens, 1831

Pearce: 4 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2021

Polydrusus (Eurodrusus) pulchellus

Stephens, 1831

Pearce: 1 locality

First subsequent record: 1995

Number of subsequent records: 6

Latest record: 2018

Polydrusus (Eustolus) flavipes (De Geer, 1775)

Pearce: 1 locality

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2021

Polydrusus (Eustolus) pterygomalis

Boheman, 1840

Pearce: 2 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2021

Polydrusus (Metallites) marginatus

Stephens, 1831

Pearce: 1 locality

First subsequent record: 2016

Number of subsequent records: 1

Latest record: 2016

Polydrusus (Polydrusus) tereticollis (De Geer, 1775)

Pearce: generally common

First subsequent record: 1978

Number of subsequent records: many

Latest record: 2016

Barypeithes sulcifrons (Boheman, 1842)

Pearce: 2 localities

First subsequent record: 1994

Number of subsequent records: 2

Latest record: 2008

Brachysomus echinatus (Bonsdorff, 1785)

Pearce: 1 locality

First subsequent record: 1982

Number of subsequent records: many

Latest record: 2020

Brachysomus hirtus (Boheman, 1845)

Pearce: 2 localities

No subsequent records

Exomias araneiformis (Schrank, 1781)

[2012: *Barypeithes araneiformis*]

Pearce: generally common

First subsequent record: 1978

Number of subsequent records: many

Latest record: 2021

Exomias pellucidus (Boheman, 1834)

[2012: *Barypeithes pellucidus*]

Pearce: 1 locality

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Sciaphilus asperatus (Bonsdorff, 1785)

Pearce: 3 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2021

Andrion regensteinense (Herbst, 1797)

Pearce: 4 localities

First subsequent record: 1927

Number of subsequent records: many

Latest record: 2021

Charagmus griseus (Fabricius, 1775)

Pearce: 1 locality

First subsequent record: 1979

Number of subsequent records: 1

Latest record: 1979

Coelositona cambricus (Stephens, 1831)

Pearce: 6 localities

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2019

Sitona ambiguus Gyllenhal, 1834

Pearce: 2 localities

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2020

Sitona cylindricollis Fåhræus, 1840

Pearce: 2 localities

First subsequent record: 1966

Number of subsequent records: many

Latest record: 2018

Sitona gemellatus Gyllenhal, 1834

Pearce: no records
First record: 1954
Number of records: many
Latest record: 2020

Sitona hispidulus (Fabricius, 1777)

Pearce: generally common and widely distributed
First subsequent record: 1936
Number of subsequent records: many
Latest record: 2021

Sitona humeralis Stephens, 1831

Pearce: 5 localities
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2019

Sitona lateralis Gyllenhal, 1834 [2012:

Sitona ononidis]

Pearce: 1 locality
First subsequent record: 1982
Number of subsequent records: 8
Latest record: 2020

Sitona lineatus (Linnaeus, 1758)

Pearce: generally common and widely distributed
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Sitona lineellus (Bonsdorff, 1785)

Pearce: no records
First record: 1982
Number of records: many
Latest record: 2020

Sitona macularius (Marsham, 1802)

Pearce: 1 locality
First subsequent record: 1994
Number of subsequent records: 4
Latest record: 2011

Sitona obsoletus (Gmelin, 1790) [2012:

Sitona lepidus]

Pearce: 6 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Sitona puncticollis Stephens, 1831

Pearce: 7 localities
First subsequent record: 1966
Number of subsequent records: 9
Latest record: 2008

Sitona striatellus Gyllenhal, 1834

Pearce: generally common
First subsequent record: 1937
Number of subsequent records: many
Latest record: 2020

Sitona sulcifrons (Thunberg, 1798)

Pearce: generally common
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2021

Sitona suturalis Stephens, 1831

Pearce: 2 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2020

Sitona waterhousei Walton, 1846

Pearce: 7 localities
First subsequent record: 1955
Number of subsequent records: many
Latest record: 2021

Tanymecus palliatus (Fabricius, 1787)

Pearce: 3 localities
First subsequent record: 1963
Number of subsequent records: many
Latest record: 2015

Caenopsis fissirostris (Walton, 1847)

Pearce: 1 locality
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2020

Caenopsis waltoni (Boheman, 1842)

Pearce: 2 localities
First subsequent record: 1977
Number of subsequent records: many
Latest record: 2021

Cathormiocerus aristatus (Gyllenhal, 1827)

Pearce: 1 locality (1896)
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2019

Cathormiocerus spinosus (Goeze, 1777)

Pearce: 1 locality
First subsequent record: 1968
Number of subsequent records: many
Latest record: 2021

Romualdius angustisetulus (Hansen, 1915)

Pearce: no records (but see *scaber*)

First record: 1991

Number of records: many

Latest record: 2021

Romualdius scaber (Linnaeus, 1758)

[2012: *Romualdius bifoveolatus*]

Pearce: 6 localities (but could include *angustisetulus*)

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2019

Trachyphloeus alternans Gyllenhal, 1834

Pearce: 5 localities

First subsequent record: 1964

Number of subsequent records: many

Latest record: 2019

Trachyphloeus digitalis (Gyllenhal, 1827)

Pearce: no records

First record: 2002

Number of records: 1

Latest record: 2002

Trachyphloeus scabriculus (Linnaeus, 1771)

Pearce: 1 locality

First subsequent record: 1979

Number of subsequent records: 9

Latest record: 2017

Trachyphloeus spinimanus Germar, 1824

Pearce: 1 locality

No subsequent records

Tropiphorus elevatus (Herbst, 1795)

Pearce: 2 localities

First subsequent record: 1984

Number of subsequent records: 6

Latest record: 2009

Brachypera zoilus (Scopoli, 1763) [2012: *Hypera zoilus*]

Pearce: 8 localities

First subsequent record: 1934

Number of subsequent records: many

Latest record: 2019

Hypera (Dapalinus) meles (Fabricius, 1792)

Pearce: no records

First record: 1979

Number of records: many

Latest record: 2015

Hypera (Eirinomorphus) conmaculata (Herbst, 1795) [2012: *Hypera pollux*]

Pearce: 3 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2020

Hypera (Eirinomorphus) rumicis (Linnaeus, 1758)

Pearce: generally common

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2021

Hypera (Hypera) melancholica (Fabricius, 1792) [2012: *Hypera fuscocinerea*]

Pearce: 2 localities

No subsequent records

Hypera (Hypera) miles (Paykull, 1792) [2012: *Hypera suspiciosa*]

Pearce: 1 locality

First subsequent record: 1979

Number of subsequent records: 5

Latest record: 2020

Hypera (Hypera) nigrirostris (Fabricius, 1775)

Pearce: generally common

First subsequent record: 1975

Number of subsequent records: many

Latest record: 2021

Hypera (Hypera) ononidis (Chevrolat, 1863)

Pearce: no records

First record: 1991

Number of records: 8

Latest record: 2020

Hypera (Hypera) plantaginis (De Geer, 1775)

Pearce: 6 localities

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2021

Hypera (Hypera) postica (Gyllenhal, 1813)

Pearce: 8 localities

First subsequent record: 1955

Number of subsequent records: many

Latest record: 2020

Hypera (Hypera) venusta (Fabricius, 1781)

Pearce: 6 localities

First subsequent record: 1963

Number of subsequent records: many

Latest record: 2019

Hypera (Kippenbergia) arator (Linnaeus, 1758)

Pearce: 8 localities

First subsequent record: 1982

Number of subsequent records: many

Latest record: 2021

Limobius mixtus (Boheman, 1834)

Pearce: 2 localities

No subsequent records

Cleonis pigra (Scopoli, 1763)

Pearce: 5 localities

First subsequent record: 1934

Number of subsequent records: 1

Latest record: 1934

Coniocleonis nebulosus (Linnaeus, 1758)

Pearce: 6 localities

First subsequent record: 1975

Number of subsequent records: 6

Latest record: 2003

Larinus carlinae (Olivier, 1807) [2012:

Larinus planus]

Pearce: 6 localities

First subsequent record: 1937

Number of subsequent records: many

Latest record: 2020

Lixus scabricollis Boheman, 1842

Pearce: no records

First record: 1998

Number of records: many

Latest record: 2020

Rhinocyllus conicus (Frölich, 1792)

Pearce: 8 localities

First subsequent record: 1953

Number of subsequent records: many

Latest record: 2019

Magdalis (Edo) ruficornis (Linnaeus, 1758)

Pearce: 2 localities

First subsequent record: 1981

Number of subsequent records: many

Latest record: 2008

Magdalis (Odontomagdalis) armigera (Geoffroy in Fourcroy, 1785)

Pearce: 2 localities

First subsequent record: 1979

Number of subsequent records: many

Latest record: 2021

Magdalis (Odontomagdalis) carbonaria (Linnaeus, 1758)

Pearce: no records

First record: 1988

Number of records: 4

Latest record: 2011

Magdalis (Panus) barbicornis (Latreille, 1804)

Pearce: no records

First record: 1985

Number of records: 1

Latest record: 1985

Magdalis (Porrothus) cerasi (Linnaeus, 1758)

Pearce: 2 localities

First subsequent record: 1980

Number of subsequent records: many

Latest record: 2019

Hylobius abietis (Linnaeus, 1758)

Pearce: fairly common and widely distributed

First subsequent record: 1945

Number of subsequent records: many

Latest record: 2019

Leiosoma deflexum (Panzer, 1795)

Pearce: 7 localities

First subsequent record: 1938

Number of subsequent records: many

Latest record: 2021

Leiosoma oblongulum Boheman, 1842

Pearce: 1 locality

First subsequent record: 2018

Number of subsequent records: 2

Latest record: 2018

Liparus coronatus (Goeze, 1777)

Pearce: 6 localities

First subsequent record: 1988

Number of subsequent records: many

Latest record: 2021

Syagrius intrudens Waterhouse, C.O., 1903

Pearce: no records
First record: 2014
Number of records: 2
Latest record: 2019

Pissodes castaneus (De Geer, 1775)

Pearce: 2 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2019

Pissodes pini (Linnaeus, 1758)

Pearce: no records
First record: 1997
Number of records: 9
Latest record: 2011

Trachodes hispidus (Linnaeus, 1758)

Pearce: no records
First record: 2001
Number of records: 1
Latest record: 2001

Orobitis cyanea (Linnaeus, 1758)

Pearce: 2 localities
First subsequent record: 2002
Number of subsequent records: many
Latest record: 2021

Platypus cylindrus (Fabricius, 1792)

Pearce: no records
First record: 1993
Number of records: many
Latest record: 2019

Hylastes angustatus (Herbst, 1794)

Pearce: no records
First record: 1979
Number of records: 7
Latest record: 2015

Hylastes ater (Paykull, 1800)

Pearce: 3 localities
First subsequent record: 1979
Number of subsequent records: many
Latest record: 2019

Hylastes attenuatus Erichson, 1836

Pearce: no records
First record: 1979
Number of records: 6
Latest record: 2015

Hylastes opacus Erichson, 1836

Pearce: 2 localities
First subsequent record: 1983
Number of subsequent records: many
Latest record: 2019

Hylurgops palliatus (Gyllenhal, 1813)

Pearce: 3 localities
First subsequent record: 2004
Number of subsequent records: many
Latest record: 2016

Hylastinus obscurus (Marsham, 1802)

Pearce: 1 locality
First subsequent record: 1987
Number of subsequent records: 4
Latest record: 2019

Hylesinus crenatus (Fabricius, 1787)

Pearce: 2 localities
First subsequent record: 1987
Number of subsequent records: 6
Latest record: 2019

Hylesinus taranio (Danthoine in Bernard, 1788)

Pearce: 2 localities
First subsequent record: 2005
Number of subsequent records: 3
Latest record: 2010

Hylesinus varius (Fabricius, 1775)

Pearce: 5 localities
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Hylesinus wachtli Reitter, 1887

Pearce: no records
First record: 2008
Number of records: 1
Latest record: 2008

Kissophagus vicinus (Comolli, 1837)

Pearce: 1 locality
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2020

Pteleobius vittatus (Fabricius, 1792)

Pearce: "Dorset" (A. Ford)
First subsequent record: 1975
Number of subsequent records: 4
Latest record: 1983

Tomicus minor (Hartig, 1834)

Pearce: no records
First record: 2001
Number of records: 2
Latest record: 2001

Tomicus piniperda (Linnaeus, 1758)

Pearce: 6 localities
First subsequent record: 1977
Number of subsequent records: many
Latest record: 2021

Phloeotribus rhododactylus (Marshall, 1802)

Pearce: 1 locality
First subsequent record: 1981
Number of subsequent records: many
Latest record: 2020

Pityophthorus pubescens (Marshall, 1802)

Pearce: 1 locality
First subsequent record: 1991
Number of subsequent records: 8
Latest record: 2020

Ernoporicus fagi (Fabricius, 1798)

Pearce: 1 locality (1825)
First subsequent record: 2019
Number of subsequent records: 2
Latest record: 2021

Trypophloeus binodulus (Ratzeburg, 1837)

Pearce: no records
First record: 1998
Number of records: 1
Latest record: 1998

Dryocoetes autographus (Ratzeburg, 1837)

Pearce: no records
First record: 1990
Number of records: 1
Latest record: 1990

Dryocoetes villosus (Fabricius, 1792)

Pearce: 1 locality
First subsequent record: 1975
Number of subsequent records: many
Latest record: 2021

Taphrorychus bicolor (Herbst, 1794)

Pearce: no records
First record: 2017
Number of records: 5
Latest record: 2019

Xylocleptes bispinus (Duftschmid, 1825)

Pearce: 1 locality
First subsequent record: 1986
Number of subsequent records: many
Latest record: 2016

Ips sexdentatus (Boerner, 1766)

Pearce: no records
First record: 2005
Number of records: 3
Latest record: 2009

Orthotomicus laricis (Fabricius, 1792)

Pearce: 1 locality
First subsequent record: 1954
Number of subsequent records: 3
Latest record: 2015

Pityogenes bidentatus (Herbst, 1784)

Pearce: 1 locality (1895)
First subsequent record: 1989
Number of subsequent records: 1
Latest record: 1989

Scolytus intricatus (Ratzeburg, 1837)

Pearce: 1 locality
First subsequent record: 1991
Number of subsequent records: many
Latest record: 2019

Scolytus mali (Beckstein in Beckstein & Scharfenberg, 1805)

Pearce: no records
First record: 1993
Number of records: 4
Latest record: 2016

Scolytus multistriatus (Marshall, 1802)

Pearce: 1 locality
First subsequent record: 1980
Number of subsequent records: 4
Latest record: 2019

Scolytus rugulosus (Müller, P.W.J., 1818)

Pearce: 1 locality
First subsequent record: 1970
Number of subsequent records: many
Latest record: 2021

Scolytus scolytus (Fabricius, 1775)

Pearce: 4 localities
First subsequent record: 1975
Number of subsequent records: 3
Latest record: 1983

Xyleborinus saxesenii (Ratzeburg, 1837)

Pearce: 1 locality

First subsequent record: 2004

Number of subsequent records: 4

Latest record: 2020

Xyleborus dryographus (Ratzeburg, 1837)

Pearce: no records

First record: 2004

Number of records: 1

Latest record: 2004

Xyleborus monographus (Fabricius, 1792)

Pearce: no records

First record: 2019

Number of records: 3

Latest record: 2020

Trypodendron domesticum (Linnaeus, 1758)

Pearce: no records

First record: 1981

Number of records: many

Latest record: 2019

Trypodendron lineatum (Olivier, 1795)

Pearce: no records

First record: 2008

Number of records: 1

Latest record: 2008

Trypodendron signatum (Fabricius, 1792)

Pearce: no records

First record: 1970

Number of records: 6

Latest record: 2020